



SEVIN* DUST

5%

REGISTERED TRADEMARK OF GORDON CORPORATION
KANSAS CITY, MISSOURI
MADE IN U.S.A.

CAUTION: This product is highly toxic to humans and animals. Avoid contact with skin and eyes. Do not breathe dust. Wash hands thoroughly after use. Do not eat or drink while using. Do not use in or near living quarters. Do not use on food crops.

MANUFACTURED BY
PBI - GORDON CORPORATION
KANSAS CITY, KANSAS

— DIRECTIONS —

Apply SEVIN DUST 5% when insects or their damage appear. Make applications to plants in the morning or evening when it is cool. Treat upper and lower surface of all foliage for complete coverage. Repeat as necessary, generally at 7-day intervals. For best results apply with a dust gun. Do not apply in excess.

VEGETABLE INSECT CONTROL

ASPARAGUS: To control Asparagus Beetle and Apache Cicada. Do not repeat application within three days. Do not treat within one day of harvest.

BEANS (Green Beans, Lima Beans, Snap Beans, Cowpeas, Black-eyed): To control Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetles, Flea Beetles, Japanese Beetle, Leafhoppers, Velvetbean Caterpillar, Western Bean Cutworm, Armyworms, Corn Earworm, Stink Bugs and Tarnish Plant Bugs.

SWEET CORN: To control Corn Earworm, Corn Rootworm adults, European Corn Borer, Fall Armyworm, Flea Beetles, Japanese Beetles, Sap Beetles, and Leafhoppers. For larvae in Whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear, and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are essential for effective control.

CUCUMBER, MELONS, PUMPKIN, SQUASH: To control Pickleworm, Melonworm, Cucumber Beetles, Flea Beetles, Leafhoppers and Squash Bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use on watermelons in Florida.

POTATO, TOMATO, EGGPLANT, PEPPER: To control Colorado Potato Beetle, Flea Beetles, Leafhoppers, European Corn Borer, Fall Armyworm, Lace Bugs, Tomato Fruit Worm, Tomato Hornworm, Tarnish Plant Bugs, Stink Bugs.

HEAD LETTUCE AND CARROTS: To control Flea Beetles, Harlequin Bug, Leafhoppers, Armyworms, Cabbage Caterpillars, Corn Earworm, Spittlebugs, Tarnish Plant Bugs and Stink Bugs. To control small Cabbage Loopers use on a 5 to 7 days schedule.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI: To control Flea Beetles, Harlequin Bugs, Armyworms, Cabbage Caterpillars, Corn Earworm. To control small Cabbage Loopers, use on a 5 to 7 day schedule.

SMALL FRUIT INSECT CONTROL

STRAWBERRIES: To control Meadow Spittlebugs, Strawberry Leaf Roller and Strawberry Weevil. Do not apply within one day of harvest.

GRASSHOPPERS ON THE ABOVE CROPS

Apply SEVIN DUST 5% as often as necessary on crops to control grasshoppers. On Asparagus follow cautions noted above.

NOTE

Allow 1 day between dusting and harvest of Strawberries. Three days on Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Head Lettuce and Kohlrabi. Do not use on Leaf Lettuce within 14 days of harvest. Injury to young tender foliage may occur if plants are wet or in the presence of high humidity of several days duration following treatment.

POULTRY (Chickens, Turkeys, Ducks, Geese, Game Birds, Pigeons)

Application of 5% SEVIN DUST to poultry for control of chicken mites and fleas is to be made as a supplement to litter treatment for control of these pests.

On birds: Northern fowl mite, chicken mite, lice, fleas 1 lb. 5% dust per 100 birds.

On litter: Northern fowl mite, chicken mite, lice, fleas, bedbugs 1 lb. 5% dust per 40 Sq. Ft.

Dust bath boxes: Northern fowl mite, chicken mite, lice and fleas 2 1/4 lbs. 5% dust per box for each 50 birds. DO NOT APPLY 7 days before slaughter. Do not treat nest litter. Avoid direct contamination of eggs. Tolerance 5 ppm. on meat and fat, 0 in eggs.

Do not repeat applications within 4 weeks.

DOGS AND CATS

Brown dog tick, fleas rub in skin and apply in sleeping quarters weekly. Do not treat kittens or puppies under 4 weeks old.

CAUTION

HARMFUL IF SWALLOWED OR INHALED. Do not take internally. Avoid breathing of dust. Skin Contact may be harmful. Avoid contact. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

E.P.A. REG. NO. 2217-572

611-1171

2217-572

12-29-1971

1/2

ACCEPTED

DEC 29 1971

UNDER THE FEDERAL INSTITUTE
 FOR THE STUDY OF
 FOREIGN BODILY SUBJECT
 TO ALL THE OBLIGATIONS

8217-572