

GORDON'S SEVIN® DUST 5%

A MULTI PURPOSE INSECTICIDE

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate)..... 5.0%


INERT INGREDIENTS..... 95.0%

TOTAL 100.0%

SEVIN® is the registered trademark of Rhone Poulenc for carbaryl insecticide.

KEEP OUT OF REACH OF CHILDREN


CAUTION



See side panel for Precautionary Statements and Statement of Practical Treatment

NET WEIGHT ONE POUND

APXXXXXX
 EPA REG. NO. 2217-572
 EPA EST. NO. 2217-KS-1
 Mfd. by PBI/GORDON CORPORATION
 KANSAS CITY, MISSOURI 64101



STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Avoid breathing vapors dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

- User Safety Requirements:**
- ◆ Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.
 - ◆ Change clothing as soon as possible after use.
 - ◆ Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
 - ◆ Do not allow children or pets to contact treated area until dust has settled.

Statement of Practical Treatment

IF SWALLOWED: Induce vomiting and seek medical attention immediately.

IF IN EYES OR ON SKIN: Flush eyes thoroughly with water; wash skin thoroughly with soap and water.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

NOTE TO PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates or cholinesterase inhibiting drugs.

ENVIRONMENTAL HAZARDS:

This pesticide is extremely toxic to aquatic and estuarine invertebrates. ~~For terrestrial uses, Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.~~ Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

STORAGE: ~~Do not contaminate water, food, or feed by storage or disposal.~~ Store in original container in a locked storage area inaccessible to children or animals. Keep in cool, dry place.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse container. Rinse thoroughly before discarding in trash.

BEST AVAILABLE COPY

- ◆ Apply Sevin® Dust 5% when insects or their damage appear.
- ◆ ~~Make applications to plants in the morning or evening when it is cool. Treat upper and lower surface of all foliage for complete coverage. Repeat as necessary, generally at 7 day intervals. For best results apply with a dust gun. Do not apply in excess.~~
- ◆ ~~Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust.~~
- ◆ Apply when air is calm to avoid drift and contact with eyes and skin.
- ◆ ~~Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering.~~
- ◆ ~~For plants above chest height, consider using a liquid carbaryl product.~~
- ◆ ~~Do not repeat application more than once a week.~~
- ◆ Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

INSECT CONTROL ON VEGETABLES

~~**Asparagus:** To control Asparagus Beetle and Apache Cicada. Do not repeat application within three days. Do not treat within one day of harvest.~~

Beans (Green Beans, Lima Beans, Snap Beans, Cowpeas, Black-eyed): To control Mexican bean beetle, bean leaf beetle, cucumber beetle, flea beetle, Japanese beetle, leafhoppers, velvetbean caterpillar, Western bean cutworm, armyworm, corn earworm, grasshoppers, stink bugs and tarnish plant bug.

Sweet Corn: To control corn earworm, corn rootworm adults, European corn borer, fall armyworm, flea beetle, Japanese beetle, sap beetle, grasshoppers, and leafhoppers. For larvae in Whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary.

For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are essential for effective control.

Cucumber, Melons, Pumpkin, Squash: To control pickleworm, melonworm, cucumber beetle, flea beetle, leafhoppers, grasshoppers, and squash bug. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use on watermelons in Florida.

Potato, Tomato, Eggplant, Pepper: To control Colorado potato beetle, flea beetle, leafhoppers, European corn borer, fall armyworm, lace bug, tomato fruitworm, tomato hornworm, grasshoppers, tarnish plant bug, stink bugs.

Head Lettuce and Carrots: To control flea beetle, harlequin bug, leafhoppers, armyworm, cabbage caterpillar, corn earworm, spittlebug, grasshoppers, tarnished plant bug and stink bugs. To control cabbage looper use on a 5 to 7 day schedule. Allow three (3) days between dusting and harvesting for head lettuce. Allow fourteen (14) days between dusting and harvest for leaf lettuce.

Cabbage, Broccoli, Brussels Sprouts, Cauliflower, Kohlrabi: To control flea beetle, harlequin bug, armyworm, cabbage caterpillar, grasshoppers, and corn earworm. To control cabbage looper use on a 5 to 7 day schedule. Allow three (3) days between dusting and harvesting.

Note for Vegetables: Injury to young tender foliage may occur if plants are wet or in the presence of high humidity for several days duration following treatment.

BEST AVAILABLE COPY

4/4

INSECT CONTROL ON SMALL FRUIT

Strawberries: To control meadow spittlebug, strawberry leaf roller and strawberry weevil. ~~Do not apply within one day of harvest. Allow 1 day between dusting and harvest of Strawberries.~~

~~TO CONTROL GRASSHOPPERS ON THE ABOVE CROPS~~

~~Apply Sevin® Dust 5% as often as necessary on crops to control grasshoppers. On Asparagus follow cautions noted above.~~

IMPORTANT NOTE: ~~Allow 1 day between dusting and harvest of Strawberries. Three days on Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Head Lettuce and Kahirabi. Do not use on Leaf Lettuce within 14 days of harvest, injury to young tender foliage may occur if plants are wet or in the presence of high humidity of several days duration following treatment.~~

~~POULTRY (Chickens, Turkeys, Ducks, Geese, Game Birds, Pigeons)~~

~~Application of 5% Sevin® Dust to poultry for control of chicken mites and fleas is to be made as a supplement to litter treatment for control of these pests:~~

~~On birds: Northern fowl mite, chicken mite, lice, fleas 1 pound 5% dust per 100 birds.~~

~~On litter: Northern fowl mite, chicken mite, lice, fleas, bedbugs 1 pound 5% dust per 40 square feet.~~

~~Dust bath boxes: Northern fowl mite, chicken mite, lice and fleas 2 1/2 pounds 5% dust per box for each 50 birds.~~

~~DO NOT APPLY 7 days before slaughter. Do not treat nest litter. Avoid direct contamination of eggs. Tolerance 5 ppm. on meat and fat, 0 in eggs.~~

~~ON DOGS AND CATS~~

~~Brown dog tick, fleas: rub in skin and apply in sleeping quarters weekly. Do not treat kitten and puppies under 4 weeks old. Do not use on pregnant dogs.~~

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tomadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of products or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

BEST AVAILABLE COPY