



TRIMEC **Weed & Feed 30** **24-4-8**

Twenty Pounds Covers up to 5,000 Sq. Ft.

KILLS: Dandelion, Chickweed, Knotweed, Plantains, Hen-bit Spurge and many other broadleaf weeds; some of which are listed on the back panel.

ACTIVE INGREDIENTS

***2,4-Dichlorophenoxyacetic acid
 ***2,4-Dimethyl-4-chlorophenoxyacetic acid
 ***2,4-Dimethyl-6-chlorophenoxyacetic acid
 ***2,4-Dimethyl-6-chlorophenoxyacetic acid
 INERT INGREDIENTS

0.001%
 0.001%
 0.001%
 0.001%
TOTAL 100.000%

THIS PRODUCT CONTAINS

***13.67% 2,4-Dichlorophenoxyacetic acid per lb. of 13.67% per acre.
 ***13.67% 2,4-Dimethyl-4-chlorophenoxyacetic acid per lb. of 13.67% per acre.
 ***13.67% 2,4-Dimethyl-6-chlorophenoxyacetic acid per lb. of 13.67% per acre.
 *Other Ingredients: ADAM, Methionine, L-Ascorbic

KEEP OUT OF REACH OF CHILDREN **CAUTION**

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water. For eye contact, wash eyes for 15 minutes and get medical attention.

See back panel for additional precautionary statements.

*TRIMEC® is a registered trademark of PBI GORDON CORPORATION.

NET WEIGHT 20 POUNDS

PBI GORDON CORPORATION
 KANSAS CITY, KANSAS 64101

BEST AVAILABLE COPY

Start! Read the entire label first.

Observe all cautions and follow directions carefully.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. Wash thoroughly after handling.

Environmental Hazards

Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate water used for irrigation or domestic purposes. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE

Store in original container and use only bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs. To prevent cross contamination, do not store near other fertilizers or herbicides, insecticides, fungicides or pesticides.

PRODUCT DISPOSAL

To dispose of bags which contain unused product, securely wrap bag in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL

To dispose of empty bag, fold and discard in trash.

USE PRECAUTIONS

Do not use this product where desirable flowers are present or allow the material to wind drift onto flowers, vegetable or ornamental shrubbery. Do not apply in fields undisturbed by roots of desirable trees or shrubs. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning. Do not scatter onto streams or irrigation waters.

INSTRUCTIONS

Grass and weeds 2 to 3 inches high. TRIMEC controls a wide range of broadleaf weeds, including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others. Apply 13.67% of 2,4-Dichlorophenoxyacetic acid per lb. of 13.67% per acre. For best results, apply when weeds are actively growing. A single application may be made in late spring or early summer. A second application may be made in late summer or early fall. Do not use on dormant or dead grass. Do not use on lawns during periods of drought or stress.

1. Use when weeds are 2 to 3 inches high.
2. Apply 13.67% of 2,4-Dichlorophenoxyacetic acid per lb. of 13.67% per acre.
3. Apply when weeds are actively growing and actively growing primarily in the morning when dew is on the grass. Grass should be cut prior to application. Spread evenly with water to hold the material and allow it to dry. Do not mow for 24 hours.
4. Do not use on dormant or dead grass. Do not use on lawns during periods of drought or stress.

Be patient, TRIMEC works slowly and gently and is friendly to your lawn.

SPREADER SETTINGS

Drift 0
 Load Capacity 5
 Discharge 25
 Speed 4

The above spreader settings are approximate. Check your spreader on 250 sq. ft. and adjust to apply at the rate of 1 lb. per 250 sq. ft.

FERTILIZER GUARANTEE

Total Nitrogen, not less than 24%
 Total Insoluble Nitrogen, not less than 5%
 Available Phosphoric Acid, not less than 4%
 Soluble Potash, not less than 8%

WEEDS			
BEDSTRAW	DANDELION	LESPEDEA	POISON IVY
BLACK MEDIC	DOCK	MALLOW	POISON OAK
BURNING	GROUND IVY	MORNING GLORY	PURSLANE
BURDOCK	HEAL-ALL	ORRIS	RAUNDEL
CHICKWEED	HENBIT	PIPPERCORN	SHEEP SORE
CHICKWEED	KNOTWEED	PIPPERCORN	SHEPHERD'S PURSE
CLOVER	LAMBSQUARTER	PLANTAIN	SPEEDWELL
			YARROW

NOTICE: Seller warrants that this product conforms to the ingredients statement on the label. Since conditions of use such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claim other than those stated on the label.

EPA REG. No. 277-579
 EPA EST. No. KS-1

975/184



*TRIMEC®

Weed & Feed 30 24-4-8

Twenty Pounds Covers up to 5,000 Sq. Ft.

KILLS: Dandelion, Chickweed, Knotweed, Plantains, Henbit, Spurge and many other broadleaf weeds; some of which are listed on the back panel.

ACTIVE INGREDIENTS:

*2,4-Dichlorophenoxyacetic acid	0.684%
**2-(2-methyl-4-chlorophenoxy) propionic acid	0.306%
***Dicamba (3,6-dichloro-o-anisic acid)	0.055%
INERT INGREDIENTS	98.945%
TOTAL	100.000%

THIS PRODUCT CONTAINS:

- *13.676 lbs. 2,4-Dichlorophenoxyacetic acid per ton or 1.191 lbs. per acre.
- **8.121 lb. 2-(2-methyl-4-chlorophenoxy) propionic acid per ton or 0.533 lb. per acre.
- ***1.291 lbs. 3,6-dichloro-o-anisic acid per ton or 0.112 lb. per acre.
- *Isomer Specific by AOAC Method 6.D01-5.

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water; for eyes flush with water for 15 minutes and get medical attention.

See back panel for additional precautionary statements.

*TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

NET WEIGHT 26 POUNDS



pbi/gordon
corporation

KANSAS CITY, KANSAS 66116

Stop! Read the entire label first.

Observe all cautions and follow directions carefully.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION. Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eye, or clothing. Wash thoroughly after handling.

Environmental Hazards

Do not apply directly to water. Do not apply when weather conditions favor target area. Do not contaminate water used for irrigation or domestic purposes. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs. To prevent cross-contamination, do not store near other fertilizers, or near herbicides, insecticides, fungicides, or seeds.

PRODUCT DISPOSAL: To dispose of bags which contain unused product, securely wrap bag in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Fold empty bag and discard in trash.

USE PRECAUTIONS

Do not use this product where desirable clovers are present or allow the material to "wind drift" onto flowers, vegetables or ornamental shrubbery. Do not apply in areas underlain by roots of desirable trees or shrubs. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning. Do not contaminate domestic or irrigation waters.

INSTRUCTIONS

Gordon's Weed & Feed 30 contains TRIMEC® herbicide. TRIMEC controls a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantains, henbit, spurge and many others commonly found in home lawns. In addition, a 24-4-8 fertilizer provides a generous supply of plant food needed for a normal season's growth. Possible retreatment may be necessary if (a) lawn is heavily over-run by weeds, or (b) adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days. Do not use on dandelion, carpetgrass, closely-mowed bentgrass putting greens or St. Augustine lawns.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see Vaso 4).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto non-target plants.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

Be patient, TRIMEC work, slowly and gently and is friendly to your lawn.

SPREADER SETTINGS

Scott	6
Lawn Beauty	5
Jackson	7 1/2
Sears	4

The above spreader settings are approximate. Check your spreader on 250 sq. ft. and adjust to apply at the rate of 1 lb. per 250 sq. ft.

FERTILIZER GUARANTEE

Total Nitrogen (not less than)	24%
Water Insoluble Nitrogen (not less than)	5%
Available Phosphoric Acid (not less than)	4%
Soluble Potash (not less than)	8%

WEEDS

BEDSTRAW	DANDELION	LESPEDEZA	POISON IVY	SPURGE
BLACK MEDIC	DOCK	MALLOW	POISON OAK	THISTLE
BUCKHORN	GROUND IVY	MORNING GLORY	PURSLANE	WILD CARROT
BURDOCK	HEAL-ALL	OXALIS	RAGWEED	WILD GARLIC
CHICORY	HENBIT	PEPPERCASS	SHEEP SORREL	WILD LETTUCE
CHICKWEED	KNOTWEED	PIGWEEED	SHEPHERD'S PURSE	WILD ONION
CLOVER	LAMBSQUARTER	PLANTAINS	SPEEDWELL	YARROW

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

EPA REG. No. 2217-579
EPA EST. No. -KS-1

575/184