	2217-579	·	4/30/98	Farma <b>A</b> mara	•	TIFICA		1/8
**************************************	Environme Wa	United Sta ental Prot ashington, D	ection Age C 20460	ncy [	Registra Amendm Other	tion	proval Expires 05-3 DPP Identifier Num	
		Applicati		icide - Section		<u> </u>		
1. Company/Product Num	ber 2217-579	•	<b>2.</b> E	PA Product Mana: Joanne 1.	•	3. Propo	sed Classification	
Company/Product (Nan Gordon's T		eed 30	? PM			$  \boxtimes  $	lone Res	stricted
5. Name and Address of A PBI/Gordon Corporat PO Box 014090 Kansas City, Missour	ion	IP Code)	(b)( to: EP	expedited Review i), my product is s A Reg. No.				· .
Check it till	s is a new address		Section	duct Name				
Amendment - Explain be Resubmission in respo	nse to Agency letter da	ted			ation.	,	<b>CATION</b> 0 1998	
registrants may presen labeling is enclosed. T 152.46, and no other cunderstand that it is a verthat if this notification is of FIFRA and I may be	his notification is changes have bed violation of 18 U.S s not consistent wi	consistent en made to i.C.Sec. 10 ith the term	with the provo the labeling 01 to willfully s of PR Notice	isions of PR No or the confident make any false e 95-2 and 40 C	otice 95-2 and tial statement statement to FR 152.46, th	d EPA regit of formul the EPA. In this product	ulations at 40 a of this prod I further under	CFR luct. I rstand
			Section	- 111				
1. Material This Product Will E Child-Resistant Packaging Yes* No * Certification must be submitted	Unit Packaging Yes No If "Yes" Unit Packaging wgt.	No. per container	Water Solubl Yes No If "Yes" Package wgt	No. per	2. Type of Co	c s		
3. Location of Net Contents Info	ormation tainer	4. Size(s) Re	tail Container	5.	Location of Label On Label On Labelng acc	Directions	oduct	
6. Manner in Which Label is Afi	fixed to Product	Lithogra Paper c Stencik	flued .	Other				
1. Contact Point (Complete item	ns directly below for ide	ntification of in	Section dividual to be cont		o process this an	plication.)		
Name	Martens		Title	of Regulatory S	1	Telephone No.	(include Area Cod	đe)
I certify that the statement I acknowledge that any kn both under applicable law.	's I have made on this t		ation achments thereto t may be punishab	are true, accurate an	d complete.	6. [	Date Application Received (Stamped)	
2. Signature	tenz_			f Regulatory Ser	vices		•••	
4. Typed Name () Craig	Martens		5. Date	pril 4, 1998		•	••••	

# GORDON'S TRIMEC® WEED & FEED 30 24-4-8

NOTIFICATION
APR 3 0 1998

## **ACTIVE INGREDIENTS:**

*2,4-dichlorophenoxyacetic acid	0.684%
**2-(2-methyl-4-chlorophenoxy) propionic acid	0.306%
***Dicamba: 3,6-dichloro-o-anisic acid	0.065%
INERT INGREDIENTS:	98.945%
TOTAL	100.000%

#### THIS PRODUCT CONTAINS:

- \*13.676 lbs. 2,4-Dichlorophenoxyacetic acid equivalent per ton
- \*\*6.121 lbs. 2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

# **KEEP OUT OF REACH OF CHILDREN**

# CAUTION

#### **Statement of Practical Treatment**

In case of contact, wash skin with plenty of soap and water.
For eyes, flush with water for 15 minutes and get medical attention.
See back panel for additional Precautionary Statements.

# NET WEIGHT (15, 16, 18 or 20) POUNDS

575 AP040498 EPA REG. NO. 2217-579 EPA EST. NO. 2217-KS-X MANUFACTURED BY:



Page 1 of 7

EPA Reg. No. 2217-579

<sup>\*\*\*1.291</sup> lbs. 3,6-dichloro-o-anisic acid equivalent per ton Isomer Specific by AOAC Methods.



# READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

#### PRECAUTIONARY STATEMENTS

#### Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

#### **ENVIRONMENTAL HAZARDS:**

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater. Do not apply when weather conditions favor drift from target area.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

### STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.

PESTICIDE DISPOSAL: Securely wrap bag in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

#### **USE PRECAUTIONS:**

Do not use on dichondra, carpetgrass, closely mowed bentgrass putting greens or St. Augustinegrass. Do not use this product where desirable clovers are present or allow the material to drift onto flowers, vegetables or ornamental shrubbery. Do not spread over exposed roots of trees and ornamentals. Do not exceed specified dosages for any area; be particularly careful within dripline of tree and other ornamental species. Apply when the air is calm. Application of this product to newly seeded areas should be delayed until the third mowing. During application, close spreader openings when turning. Do not contaminate domestic or irrigation waters.

#### **INSTRUCTIONS:**

Gordon's Trimec® Weed & Feed 30 contains Trimec® herbicide. Trimec® controls a wide range of lawn weeds, including dandelion, chickweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a (X-X-X analysis of fertilizer grade) fertilizer provides a supply of plant food. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days. Treated areas may be seeded 3 to 4 weeks after application.

Note to Supplemental Registrants: Guarantees for plant nutrients other than nitrogen, phosphorous, and potassium may be permitted or required by regulation by the \_\_\_\_. The guarantees for such other nutrients shall be expressed in the form of the element. The source (oxides, salts, chelates, etc.) of such other nutrients may be required to be stated and may be included on the label.

For state registrations in Kentucky, add the following paragraph:

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing (\*Number of pounds) of nitrogen per 1,000 square feet. Please consult your local extension office for questions regarding a regular fertilizer program suitable for your geographic region.

The maximum application rate to turf is 1.2 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

- 1.5 Mow lawn to normal height 1 to 2 days before application.
- 2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
- Apply when weeds are young and actively growing, preferably in the morning when the dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants.
- 4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

Spreader Settings	
Scott	6
Lawn Beauty	5
Jackson	71/2
Sears	4

The above spreader settings are approximate. (15 pound bag: Calibrate your spreader on 330 square feet and adjust to apply 1.0 pound of product per 330 square feet of turfgrass. (16 pound bag: Calibrate your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet) (18 pound bag: Calibrate your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet) (20 pound bag: Calibrate your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.)

24-4-8 GUARANTEED ANALYSIS	6: 1: ::
Total Nitrogen (N)	24%
Water Insoluble Nitrogen	5%
Available Phosphate (P2O5)	4%
Soluble Potash (K <sub>2</sub> O)	8%

Note to Supplemental Registrants: Fertilizer sources may vary in micronutrient composition, and iron may discolor or stain concrete. For fertilizers containing iron: Sweep all fertilizer from concrete surfaces immediately after application. Do not wash this product from concrete surfaces because fertilizers which contain iron may cause discoloration upon oxidation.

Bedstraw	Knotweed	∵ Ragweed
Black Medic	. Lambsquarters	Sheep Sorrel
Buckhorn	Lespedeza	Shepherdspurse
Burdock	Mallow	Speedwell
Chickweed	Morningglory	Spurge
Chicory	Oxalis	Thistle
Clover	Peppergrass	Wild Carrot
Dandelion	Pigweed	Wild Garlic
Dock	Plantain	Wild Lettuce
Ground Ivy	Poison Ivy	Wild Onion
Healall .	Poison Oak	Yarrow
Henbit	Pursiane	

#### LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

#### **APPENDIX**

# I. Previously Approved Fertilizer Grades:

◆ Fertilizer grades with nutrient contents expressed as the percentages of N,P₂O₅, and K₂O include the following:

10-6-4	16-3-5	20-10-0-(1)	23-7-7	25-4-8	28-3-8
10-16-20	16-6-9	20-10-5	24-3-7	26-3-3	29-3-4
10-18-10	18-3-5	21-5-7	24-4-8	26-3-5	30-3-4
12-5-5	18-6-12	22-6-6	25-3-3	26-4-6	<b>30</b> -3-5
15-5-10	20-4-6	22-7-7	25-3-5	28-3-3	30-3-10
16-2-3	20-4-8	22-10-10	25-3-8	28-3-7	30-4-8

The fertilizer grade guaranteed analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations. The following guaranteed analyses presentations have previously been submitted as notifications:

10-16-20 GUARANTEED ANALYSIS	
Total Nitrogen (N)	10%
6.0% Ammoniacal Nitrogen	
4.0% Urea Nitrogen	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> )	16%
Soluble Potash (K <sub>2</sub> O)	20%
Magnesium (Mg)	1.0%
0.1% Water Soluble Magnesium (Mg)	
Sulfur (S)	2.0%
2.0% Combined Sulfur (S) .	
Iron (Fe)	1.0%
Manganese (Mn)	0.5%
0.1% Water Soluble Manganese (Mn)	
Nutrient Sources: Urea, Ammonium Phosphate, Ammonium Sulfate, Muriate of I	Potash,
Ferrous Sulfate, Manganese Oxide, Manganese Sulfate, Sulfate of Potash Magnes	sia and
Magnesium oxide.	

Total Nitrogen (N)	29%
1.1% Ammoniacal Nitrogen	
27.9% Urea Nitrogen .	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> )	3%
Soluble Potash (K <sub>2</sub> O)	4%
Derived from: Polymer-coated urea, urea, diammonium phosphate, and muriate of p	otash.
*11.0% slowly available urea nitrogen from polymer coated urea.	
Chlorine (Cl), note more than	4.0%


Total Nitrogen (N)	10%
10.0% Ammoniacal Nitrogen	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> )	18%•
Soluble Potash (K <sub>2</sub> O)	10%
Sulfur (S)	2.0% _
5.0% Combined Sulfur (S)	•

26-4-6 GUARANTEED ANALYSIS		
Total Nitrogen (N)		26%
1.8% Ammoniacal Nitrogen		
24.4% Urea Nitrogen		,
Available Phosphoric Acid(P <sub>2</sub> O <sub>5</sub> )		4%
Soluble Potash (K <sub>2</sub> O)		6%
Boron (B)	0.02%	ŀ
Copper (Cu)	0.05%	
Iron (Fe)	1.00%	
Total Manganese (Mn)	0.06%	
Molybenum (Mo)	0.0005%	
Zinc (Zn)	0.05%	
Primary Plant Food Sources: Sulfur coated urea, urea, diammonium pho	sphate, mu	riate of
potash.		]
Secondary Plant Food Sources: Fertilizer grade borate, sulfates and		'
sulfates and oxides of iron, oxides of manganese, fertilizer grade molyl	bdate, and s	ulfates
and oxides of zinc.		j
*4.0% slowly available urea nitrogen from sulfur coated urea.		
Chlorine (CI), not more than		4.8%

28-3-7 GUARANTEED ANALYSIS Total Nitrogen (N)	28%
1.2% Ammoniacal Nitrogen	2070
<b>.</b>	
26.8% Urea Nitrogen	
Available Phosphoric Acid(P <sub>2</sub> O <sub>5</sub> )	3%
Soluble Potash (K <sub>2</sub> O)	7%
Primary Plant Food Sources: Polymer-coated urea, urea, diammonium phosphate,	nuriate
of potash .	
*7.0% slowly available urea nitrogen from polymer-coated urea.	
Chlorine (Cl), not more than	6.4%

#### II. Advertising claims that may be presented on the retail containers:

- ◆ (Fifteen, Sixteen, Eighteen or Twenty) Pounds Covers Up To 5,000 Square Feet.
- ♦ Controls dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf weeds; some of which are listed on the back panel
- ◆ An 18-6-12 fertilizer provides slow-release feeding to promote winter hardiness and drought tolerance in lawns.
- Feeds your lawn for up to 12 weeks.
- Kills over 30 varieties of weeds while feeding your lawn.
- Greens lawns and develops deep roots while controlling weeds.
- Bring out the best with the turf magic touch!
- · Kills dandelions, plantain, and other weeds.
- Feeds the lawn with two (2) types of nitrogen and is pelleted for easy application.
- Controls (Number) weeds! (Listing of weeds may include the following.)



			WE	EDS+	
A	nnual yellow sweetclover	Cockle		Musk thistle	Spiny amaranth
A	ster	Cocklebur	51	Mustard	Spiny cocklebur
	ustrian fieldcress	Common mullein		Narrowleaf plantain	Spiny sowthistle
B	edstraw	Creeping jenny	1	Narrowleaf vetch	Spotted catsear
	eggarticks	Cudweed		Nettle	Spotted spurge
		-Curly dock		Orange hawkweed	Spurweed
	indweed, field	Daisy, English		Oriental cocklebur	Stinging nettle
	ird vetch	Daisy fleabane	~	Oxalis	Strawberry, India mock
1	itter wintercress	Daisy, oxeye		Parsley-piert	Tall nettle
	ittercress, hairy	_Dandelion		Parsnip	Tall vervain
	itterweed	Dichondra		Pearlwort	Tansy ragwort
	lack-eyed Susan	Dogbane		Pennycress	Tansy mustard
	lack medic	Dogfennel		Pennywort	Tanweed
	lack mustard	Dollarweed	_	-Peppergrass	-Thistle
1 -	lackseed plantain	Elderberry		Pepperweed	Trailing crownvetch
	lessed thistle	False dandelion		-Pigweed	Tumble mustard
	loodflower milkweed	False flax		Pineywoods bedstraw	Tumble pigweed
	lue lettuce	False sunflower		Plains coreopsis (tickseed)	Velvetleaf
	lue vervain	Fiddleneck	_	Plantain	Venice mallow
1 -	racted plantain	Florida pusley		Poison ivy	Virginia buttonweed
	rassbuttons	Frenchweed		Poison oak	Virginia creeper
	ristly oxtongue	Galinsoga	_	Pokeweed	Virginia pepperweed
	roadleaf dock	Goathead		Poorjoe	Water pennywort
1	roadleaf plantain	Goldenrod		Prairie sunflower	Wavyleaf bullthistle
- 1	=	- Ground ivy		Prickly lettuce	Western clematis
Bı	uckhorn	Gumweed		Prickly sida	Western salsify
	uckhorn plantain	Hairy fleabane		Prostrate knotweed	White mustard
	ulbous buttercup	Hawkweed		Prostrate pigweed	Wild aster
	ull thistle	-Healail		Prostrate spurge	Wild buckwheat
1	ull nettle	Heartleaf drymary		Prostrate vervain	-Wild carrot
	urclover	Heathaster		Puncturevine	Wild four-o'clock
	urdock	Hedge bindweed		Purslane, common	_Wild garlic
Bı	urning nettle .	Hedge mustard		Ragweed	Wild geranium
	urweed	Hemp		Red sorrel	-Wild lettuce
Bu	uttercup ·	_Henbit		Redroot pigweed	Wild marigold
	uttonweed	Hoary cress		Redstem filaree	Wild mustard
	anada thistle	Hoary plantain		Rough cinquefoil	-Wild onion
Ca	arolina geranium	Hoary vervain		Rough fleabane	Wild parsnip
	arpetweed	Horse nettle		Russian pigweed	Wild radish
Ca	atchweed bedstraw	Jimsonweed		Russian thistle	Wild rape
C	atnip	Knawel		Scarlet pimpernel	Wild strawberry
		-Knotweed		Scotch thistle	Wild sweet potato
↓CI	hickweed, common	Kochia ·		Sheep sorrel	Wild yetch
CI	hickweed, mouseear	Lambsquarters		Shepherdspurse	Woodsorrel
		Lespedeza		Slender plantain	Woolly croton
		~ Mallow		Smallflower galinsoga	Woolly morningglory
	lover, crimson	Matchweed		Smooth dock	Woolly plantain ****
	lover, hop	Mexicanweed		Smooth pigweed	Wormseed
	lover, red	Milk vetch	•	Sorrel (sour dock)	-Yarrow •
		- Morningglory		Sowthistle	Yellow rocket
	lover, sweet	Mouseear hawkweed		Spanish needles	Yellowflower pepperweed
1	lover, white	Mugwort	. 🖘	Speedwell	and other broadlenf weeds