

# UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICEOF PREVENTION, PLATICIDES AND TOXIC SUBSTANCES

APR 2.6 2007

James L. Kunstman, Ph.D. Director of Regulatory Services PBI/Gordon Corporation Post Office Box 014090 Kansas City, Missouri 64101

RE: To Add Information Regarding the Contents and Levels of Metals Available on the Internet

EPA Registration Number: 2217-579 Date of Submission: March 21, 2007

# Dear Dr. Kunstman:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated March 21, 2007, for the product Gordon's Trimec Weed and Feed 30 24-8-8. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

> Sincerely, 1-

Linda Arrington Notifications & Minor Formulations Team Leader Registration Division (7505P) Office of Pesticide Programs

'ease read instructions on I	everse before completing form.	Form Approved. ON	18 No. 2070-0060. Approval Expires 2-28-95		
<b>≎</b> EPA	United States Environmental Protection Washington, DC 20460	Agency An	gistration  OPP Identifier Number  nendment  ner		
	Application for	Pesticide - Section I			
. Company/Product Num Gordon's Trime		2. EPA Product Manager Joanne I. Miller	3. Proposed Classification		
. Company/Product (Nar	ne)	PM#	None Restricted		
	2217-579	Product Manager—Tea			
PBI/Gordon Corpora Post Office Box 0140 Kansas City, Missour	90	(b)(i), my product is similar or EPA Reg. No.	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No.  Product Name		
Check if th	is is a new address	17044017141715			
	Sec	ction ~ II			
Amendment - Explain	below.	Final printed labels in res Agency letter dated	sponse to NOTIFICATION		
Resubmission in resp	onse to Agency letter dated		APR 2 6 2007		
Notification - Explain	pelow.	Other - Explain below.			
contents and levels of resections of the master lenclosed draft label. This notification is consistent abeling or Confidential State the EPA. I further understant	esticide Registration Notice (PRN) senetals in this product, is available on abel for this product. This referral is with the provisions of <u>PR Notice 98-10</u> and Expent of Formula of this product. I understand to that if this notification is not consistent with the orenforcement action and penalties under Sec	the Internet as <a href="http://www.aap">http://www.aap</a> presented as highlighted text of the regulations at 40 CFR 152.46, and that it is a violation of 18 U.S.C Sec. 1 to terms of PR Notice 98-10 and 40 CFR 152.46.	fco.org/metals.htm" to all three on pages 4, 14, and 26 of the		
	Ser	tion - III			
I. Material This Product Will	<del></del>				
Child-Resistant Packaging Yes* No Certification must	Unit Packaging Wate Yes  No. per if "Yes" Wate	Yes No No. per	pe of Container  Metal Plastic Glass Paper		
be submitted	Unit Packaging wgt. container Pack	age wgt. container	Other (Specify)		
Label Contents Ir  Label Cont  Manner in Which Label is A	ainer	On Lat	of Label Directions  peling accompanying product		
	Sec	tion - IV			
. Contact Point (Complete ite	ems directly below for identification of individua	I to be contacted, if necessary, to prod	cess this application )		
lame James L. F	Cunstman, Ph.D.	rector of Regulatory Services	Telephone No. (Include Area Code) 81C 46C-6292		
I certify that the stateme I acknowledge that any I both under applicable lat Signature	Certification  nts I have made on this form and all attachmen  nowingly false or misleading statement may be v.  3. Title	ts thereto are true, accurate and com, e punishable by fine or imprisonment i	6. Date Application Received (Stamped)		
		ctor of Regulatory Services			
. Typed Name James L. K	unstman, Ph.D.	March 21, 2007			

NOTIFICATION
APR 2 6 2007

# Master Label for GORDON'S TRIMEC® WEED & FEED 30 24-4-8

# Includes:

Sublabel 1	Gordon's Trimec® Weed & Feed 30 24-4-8
Sublabel 2	For use by supplemental distributor for Bayer Advanced <sup>TM</sup> Triple Action Lawn Fertilizer Plus Weed Control <sup>II</sup>
Sublabel 3	For use by supplemental distributor for Bayer Advanced <sup>TM</sup> Triple Action Winterizer Fertilizer Plus Weed Control <sup>II</sup>

575 AP032107 EPA REG. NO. 2217-579 EPA EST. NO. 2217-KS-X MANUFACTURED BY:



Telephone: 1-800-821-7925

# GORDON'S TRIMEC® WEED & FEED 30

24-4-8

Sublabel 1: For use for Gordon's Trimec® Weed & Feed 30 24-4-8, and supplemental registrations.

**ACTIVE INGREDIENTS:** 

2,4-dichlorophenoxyacetic acid	0.69%
(+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid	0.15%
Dicamba: 3,6-dichloro-o-anisic acid	0.06%
INERT INGREDIENTS:	<u>99.10%</u>
TOTAL	100.00%

# THIS PRODUCT CONTAINS:

13.8 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton 3.0 lbs. (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid equivalent per ton 1.2 lbs. 3,6-dichloro-o-anisic acid equivalent per ton Isomer Specific by AOAC Methods.

Contains the single isomer form of Mecoprop-p.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

# KEEP OUT OF REACH OF CHILDREN

# CAUTION

See back panel for Precautionary Statements.

NET WEIGHT: (15, 16, 18 or 20) POUNDS

575 AP032107 EPA REG. NO. 2217-579 EPA EST. NO. 2217-KS-X MANUFACTURED BY:

An Employee-Owned Company
1217 West 12th Street
Kansas Chy, Mesouri 64101

Telephone: 1-800-821-7925



# READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

# PRECAUTIONARY STATEMENTS

# Hazards to Humans & Domestic Animals

**CAUTION:** Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

# First Aid

If in eyes:	<ul> <li>Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If on skin or clothing:	<ul> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If swallowed:	<ul> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
for treatment.	ct container or label with you when calling a poison control center or doctor or going
You may also ca	all toll-free, 1-877-800-5556 for emergency medical treatment information.

# **ENVIRONMENTAL HAZARDS:**

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater. Do not apply when weather conditions favor drift from target area.

# **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

# STORAGE AND DISPOSAL

**STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.

# **DISPOSAL:**

If Empty—Do not reuse this container. Place in trash or offer for recycling if available.

If Partly Filled—Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

# **USE PRECAUTIONS:**

Do not use on dichondra, carpetgrass, closely mowed bentgrass putting greens or St. Augustinegrass. Do not use this product where desirable clovers are present or allow the material to drift onto flowers, vegetables or ornamental shrubbery. Do not spread over exposed roots of trees and ornamentals. Do not exceed specified dosages for any area; be particularly careful within dripline of tree and other ornamental species. Apply when the air is calm. Application of this product to newly seeded areas should be delayed until the third mowing. Do not contaminate domestic or irrigation waters.

# **INSTRUCTIONS:**

Gordon's Trimec® Weed & Feed 30 contains Trimec® herbicide. Trimec® controls a wide range of lawn weeds, including dandelion, chickweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a (X-X-X analysis of fertilizer grade) fertilizer provides a supply of plant food. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days. Treated areas may be seeded 3 to 4 weeks after application.

**Note to Supplemental Registrants:** Guarantees for plant nutrients other than nitrogen, phosphorous, and potassium may be permitted or required by regulation by the \_\_\_. The guarantees for such other nutrients shall be expressed in the form of the element. The source (oxides, salts, chelates, etc.) of such other nutrients may be required to be stated and may be included on the label.

# For state registrations in Kentucky, add the following paragraph:

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing (\*Number of pounds) of nitrogen per 1,000 square feet. Please consult your local extension office for questions regarding a regular fertilizer program suitable for your geographic region.

# Labeling Requirements:

- All fertilizer products distributed in Washington at a minimum, must have the following labeling statement: "This product has been registered with the Washington State Department of Agriculture. When applied as directed, this fertilizer meets the Washington standards for arsenic, cadmium, cobalt, mercury, molybdenum, lead, nickel, selenium, and zinc. You have the right to receive specific information about Washington standards from the distributor of this product."
- After July 1, 1999, the label must also state: "Information received by the Washington State
  Department of Agriculture regarding the components in this product is available on the internet at
  http://www.wa.gov/agr/."
- -OR- Information regarding the contents and levels of metals in this product, is available on the Internet as <a href="http://www.aapfco.org/metals.htm">http://www.aapfco.org/metals.htm</a>.

The maximum application rate to turf is 1.2 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

- 1. Mow lawn to normal height 1 to 2 days before application.
- 2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
- Apply when weeds are young and actively growing, preferably in the morning when the dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants.
- Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

- 1. Drop spreaders and rotary spreaders must be calibrated for accurate application of this product.
- 2. Close spreader opening (orifice).
- 3. Adjust spreader setting using the Spreader Suggestions chart

Application Rate = 3 lbs. of product per 1,000 square feet.

# SUGGESTED SPREADER SETTINGS:

Centrifugal/Rotary Spreaders	Application Rate, 3 lbs./1,000 sq.ft.
Frank's Broadcast	2 1/4
Scotts Speedy Green	5
Radio Flyer #0047	8 1/4
True Temper Rotary CB2000	3 ¾
Earthway Rotary #2050SU	14 ½
Drop Spreaders	Application Rate, 3 lbs./1,000 sq.ft.
Drop Spreaders Frank's Drop	Application Rate, 3 lbs./1,000 sq.ft.
Frank's Drop	8 1/4
Frank's Drop Greenview II	8 ¼ 5 ½

These spreader settings are approximate. Differences may occur due to the condition of spreader, speed at which applied and roughness of area treated. Check your spreader on 330 square feet and adjust to apply at the rate of 1 pound per 1,000 square feet.

# 4. CALIBRATION INSTRUCTIONS:

- For 3.0 pounds of product per 1,000 square feet of turfgrass: Check you spreader on 330 square feet and adjust to apply at the rate of 1 pound of product per 330 square feet.
- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check you spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check you spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 4.0 pounds of product per 1,000 square feet of turfgrass: Check you spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.
- 5. Fill spreader. Fill spreader on driveway or sidewalk to eliminate the chance of spillage on the lawn.
- 6. Apply the product to lawn. Refer to the patch charts below:

Path A for centrifugal/rotary spreaders.




Path B for drop spreaders.




- 7. Close the spreader openings when turning to change directions or when stopping.
- 8. If using a centrifugal/rotary spreader, avoid spreading product into adjacent landscape, flowers or vegetable plantings.

NOTE: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 square feet.

This \_\_\_ pound bag should cover approximately \_\_\_\_ square feet (\_\_\_ ft. x \_\_\_ ft.) Refer to the table and picture to determine the size of your lawn.





24-4-8 GUARANTEED ANALYSIS	
Total Nitrogen (N)	24%
Water Insoluble Nitrogen	5%
Available Phosphate (P2O5)	4%
Soluble Potash (K <sub>2</sub> O)	8%

**Note to Supplemental Registrants:** Fertilizer sources may vary in micronutrient composition, and iron may discolor or stain concrete. For fertilizers containing iron: Sweep all fertilizer from concrete surfaces immediately after application. Do not wash this product from concrete surfaces because fertilizers which contain iron may cause discoloration upon oxidation.

	- WEEDS -	
Bedstraw	Knotweed	Ragweed
Black medic	Lambsquarters	Sheep sorrel
Buckhorn	Lespedeza	Shepherdspurse
Burdock	Mallow	Speedwell
Chickweed	Morningglory	Spurge
Chicory	Oxalis	Thistle
Clover	Peppergrass	Wild carrot
Dandelion	Pigweed	Wild garlic
Dock	Plantain	Wild lettuce
Ground ivy	Poison ivy	Wild onion
Healall	Poison oak	Yarrow
Henbit	Purslane	

# LIMITED WARRANTY AND DISCLAIMER.

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once to the seller.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

# I. Previously Approved Fertilizer Grades:

• Fertilizer grades with nutrient contents expressed as the percentages of N,P<sub>2</sub>O<sub>5</sub>, and K<sub>2</sub>O include the following:

10-6-4	16-3-5	20-10-0-(1)	23-7-7	25-4-8	28-3-8	32-4-8
10-16-20	16-6-9	20-10-5	24-3-7	26-3-3	29-3-4	22-3-14
10-18-10	18-3-5	21-5-7	24-4-8	26-3-5	30-3-4	30-3-4
12-5-5	18-6-12	22-6-6	25-3-3	26-4-6	30-3-5	
15-5-10	20-4-6	22-7-7	25-3-5	28-3-3	30-3-10	
16-2-3	20-4-8	22-10-10	25-3-8	28-3-7	30-4-8	

The fertilizer grade guaranteed analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations. The following guaranteed analyses presentations have previously been submitted as notifications:

10-16-20 GUARANTEED ANALYSIS	
Total Nitrogen (N)	10%
6.0% Ammoniacal Nitrogen	
4.0% Urea Nitrogen	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> )	16%
Soluble Potash (K <sub>2</sub> O)	20%
Magnesium (Mg)	1.0%
0.1% Water Soluble Magnesium (Mg)	
Sulfur (S)	2.0%
2.0% Combined Sulfur (S)	
Iron (Fe)	1.0%
Manganese (Mn)	0.5%
0.1% Water Soluble Manganese (Mn)	
Nutrient Sources: Urea, Ammonium Phosphate, Ammonium Sulfate, Muriate of	Potash.
Ferrous Sulfate, Manganese Oxide, Manganese Sulfate, Sulfate of Potash Magne	sia and
Magnesium oxide.	

29-3-4 GUARANTEED ANALYSIS	
Total Nitrogen (N)	29%
1.1% Ammoniacal Nitrogen	
27.9% Urea Nitrogen	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> )	3%
Soluble Potash (K <sub>2</sub> O)	4%
Derived from: Polymer-coated urea, urea, diammonium phosphate, and muriate of p	otash.
*11.0% slowly available urea nitrogen from polymer coated urea.	
Chlorine (Cl), note more than.	4.0%

10-18-10 GUARANTEED ANALYSIS	
Total Nitrogen (N)	10%
10.0% Ammoniacal Nitrogen	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> )	18%
Soluble Potash (K <sub>2</sub> O)	10%
Sulfur (S)	2.0%
5.0% Combined Sulfur (S)	
Nutrient Sources: Ammonium Sulfate, Ammonium Phosphate and Muriate of Potash.	

Page 8 of 35

EPA Reg. No. 2217-579

26-4-6 GUARANTEED ANALYSIS	
Total Nitrogen (N)	26%
1.8% Ammoniacal Nitrogen	
24.4% Urea Nitrogen	
Available Phosphoric Acid(P <sub>2</sub> O <sub>5</sub> )	4%
Soluble Potash (K <sub>2</sub> O)	6%
Boron (B) 0.02%	
Copper (Cu)	
Iron (Fe)	
Total Manganese (Mn)	
Molybenum (Mo) 0.0005%	
Zinc (Zn)	
Primary Plant Food Sources: Sulfur coated urea, urea, diammonium phosphate, murpotash.	riate of
Secondary Plant Food Sources: Fertilizer grade borate, sulfates and oxides of	conner
sulfates and oxides of iron, oxides of manganese, fertilizer grade molybdate, and s	
and oxides of zinc.	,4114405
*4.0% slowly available urea nitrogen from sulfur coated urea.	
Chlorine (CI), not more than	4.8%

28-3-7 GUARANTEED ANALYSIS	
Total Nitrogen (N)	28%
1.2% Ammoniacal Nitrogen	
26.8% Urea Nitrogen	
Available Phosphoric Acid(P <sub>2</sub> O <sub>5</sub> )	3%
Soluble Potash (K <sub>2</sub> O)	7%
Primary Plant Food Sources: Polymer-coated urea, urea, diammonium phosphate, mu	riate of
potash	
*7.0% slowly available urea nitrogen from polymer-coated urea.	
Chlorine (Cl), not more than	6.4%

# II. Advertising claims that may be presented on the retail containers:

- (Fifteen, Sixteen, Eighteen or Twenty) Pounds Covers Up To 5,000 Square Feet.
- Controls/Kills dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf weeds; some of which are listed on the back panel
- ♦ An 18-6-12 fertilizer provides slow-release feeding to promote winter hardiness and drought tolerance in lawns.
- Feeds your lawn for up to 12 weeks.
- Controls/Kills over 30 varieties of weeds while feeding your lawn.
- · Greens lawns and develops deep roots while controlling weeds.
- Bring out the best with the turf magic touch!
- Controls/Kills dandelion, plantain, and other weeds.
- Feeds the lawn with two (2) types of nitrogen and is pelleted for easy application.
- ◆ Controls (Number) weeds! (Listing of weeds may include the following.)
- Kills dandelion, chickweed, clover and most common lawn weeds.
- · Triple Action Feeding works at every level.

Grass- Greener, Thicker lawn

Roots - Stronger Root System

Soil - Enriches & Fortifies Soil

Builds a better lawn – Guaranteed

- Gordon's Trimec Weed & Feed 30 kills unsightly weeds such as dandelion, chickweed and clover
  while it nourishes your lawn. Its Triple Action formula contains slow-release nitrogen which
  continuously feeds the grass, roots and soil for up to two full months [all winter long]. And it's
  guaranteed not to burn. The result is a greener, thicker lawn [this Fall and quicker green-up next
  Spring]. You're just one application away from a better lawn. That's the [company name] Better
  Lawn Guarantee.
- Triple Action Feeding works at every level for a Deep-Down Green!
- Why is Triple Action better? By feeding at all levels, your lawn receives the primary nutrients it
  needs to thrive. This ensures thicker, greener grass, stronger root system, and richer soil the
  key to building a lush lawn that lasts.
- Grass: Builds thicker, greener grass and enhances overall appearance.
- Roots: Builds stronger, deeper root systems which transport nutrients from the soil to the grass to help lawn withstand heat-stress and drought.
- Soil: Enriches and fortifies your soil. Rich, fertile soil provides a foundation for establishing roots and replenishes vital primary nutrients.
- For questions or comments, call toll-free 1-877-bayerAL (1-877-229-3725)
- Made in the U.S.A.
- Kills weeds and builds a thick, green Fall lawn.
- Nourishes all Winter for a quick Spring green-up.



- Controls most annual broadleaf weeds
- Premium slow release fertilizer stimulates grass growth to quickly conceal where weeds have died.
  - NITROGEN for Dense Green Lawns
  - o PHOSPHORUS Improves Root Growth
  - o POTASSIUM Vital for Growth and Drought Resistance

		WEEDS -	
Annual yellow sweetclover	Clover, white	Mustard	Spiny cocklebur
Aster	Cockle	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Cocklebur	Narrowleaf vetch	Spotted catsear
Bedstraw	Common mullein	Nettle	Spotted spurge
Beggarticks	Creeping Charlie	Orange hawkweed	Spurweed
Betony, Florida	Creeping jenny	Oriental cocklebur	Stinging nettle
Bindweed, field	Cudweed	Oxalis	Strawberry, India mock
Bird vetch	Curly dock	Parsley-piert	Tall nettle
Bitter wintercress	Daisy, English	Parsnip	Tall vervain
Bittercress, hairy	Daisy fleabane	Pearlwort	Tansy ragwort
Bitterweed	Daisy, oxeye	Pennycress	Tansy mustard
Black-eyed Susan	Dandelion	Pepperweed	Tanweed
Black medic	Dichondra	Pigweed	Thistle
Black mustard	Dogfennel	Pineywoods bedstraw	Trailing crownvetch
Blackseed plantain	Dollarweed	Plains coreopsis	Tumble mustard
Blessed thistle	False dandelion	Plantain	Tumble pigweed
Bloodflower milkweed	Falseflax	Poison ivy	Velvetleaf
Blue lettuce	False sunflower	Poison oak	Venice mallow
Blue vervain	Fiddleneck	Pokeweed	Virginia buttonweed
Bracted plantain	Florida pusley	Poorjoe	Virginia creeper
Bristly oxtongue	Frenchweed	Prairie sunflower	Virginia pepperweed
Broadleaf dock	Galinsoga	Prickly lettuce	Wavyleaf bullthistle
Broadleaf plantain	Goldenrod	Prickly sida	Western clematis
Broomweed	Ground ivy	Prostrate knotweed	Western salsify
Buckhorn	Gumweed	Prostrate pigweed	White mustard
Buckhorn plantain	Hairy fleabane	Prostrate spurge	Wild aster
Bulbous buttercup	Hawkweed	Prostrate vervain	Wild buckwheat
Bull thistle	Healall	Puncturevine	Wild carrot
Bullnettle	Heartleaf drymary	Purslane, common	Wild four-o'clock
Burclover	Heath aster	Ragweed	Wild garlic
Burdock	Hedge bindweed	Red sorrel	Wild geranium
Burning nettle	Hedge mustard	Redroot pigweed	Wild lettuce
Burweed	Henbit	Redstem filaree	Wild marigold
Buttercup	Hoary cress	Rough cinquefoil	Wild mustard
Buttonweed	Hoary plantain	Rough fleabane	Wild onion
Canada thistle	Hoary vervain	Russian pigweed	Wild parsnip
Carolina geranium	Horsenettle	Russian thistle	Wild radish
Carpetweed	Jimsonweed	Scarlet pimpernel	Wild rape
Catchweed bedstraw	Knawel	Scotch thistle	Wild strawberry
Catnip	Knotweed	Sheep sorrel	Wild sweet potato
Catsear	Kochia	Shepherdspurse	Wild vetch
Chickweed, common	Lambsquarters	Slender plantain	Woodsorrel
Chickweed, mouseear	Lespedeza	Smallflower galinsoga	Woolly croton
Chicory	Mallow	Smooth dock	Woolly morningglory
Cinquefoil	Matchweed	Smooth pigweed	Woolly plantain
Clover, crimson	Mexicanweed	Sorrel	Wormseed
Clover, hop	Milk vetch	Sowthistle	Yarrow
Clover, red	Morningglory	Spanishneedles	Yellow rocket
Clover, strawberry	Mouseear hawkweed	Speedwell	Yellowflower pepperweed
Clover, sweet	Mugwort	Spiny amaranth	and other broadleaf weeds
	Musk thistle	-	_

# GORDON'S TRIMEC® WEED & FEED 30 24-4-8

Sublabel 2: For use by supplemental distributor for *Bayer Advanced Triple Action Lawn Fertilizer Plus Weed Control* "

# **ACTIVE INGREDIENTS:**

2,4-D (CAS No. 94-75-7)	0.69%
Mecoprop-p (CAS No. 16484-77-8)	0.15%
Dicamba (CAS No. 1918-00-9)	0.06%
INERT INGREDIENTS:	<u>99.10%</u>
TOTAL.	100.00%

### THIS PRODUCT CONTAINS:

13.8 lbs. 2.4-D acid equivalent per ton
3.0 lbs. Mecoprop-p acid equivalent per ton
1.2 lbs. Dicamba acid equivalent per ton
Isomer Specific by AOAC Methods.
Contains the single isomer form of Mecoprop-p.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

# KEEP OUT OF REACH OF CHILDREN

# CAUTION

See back panel for Precautionary Statements.

NET WEIGHT: (15, 16, 18 or 20) POUNDS

575 AP032107 EPA REG. NO. 2217-579 EPA EST. NO. 2217-KS-X MANUFACTURED BY:

An Employee-Owned Company 1217 West 12th Street Kursus City, Misseun 84101

Telephone: 1-800-821-7925

# PRECAUTIONARY STATEMENTS

# Hazards to Humans & Domestic Animals

**CAUTION:** Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

# First Aid

If in eyes:	<ul> <li>Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If on skin or clothing:	<ul> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
if swallowed:	<ul> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
for treatment.	ct container or label with you when calling a poison control center or doctor or going all toll-free, 1-877-800-5556 for emergency medical treatment information.

# **ENVIRONMENTAL HAZARDS:**

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater. Do not apply when weather conditions favor drift from target area.

# **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until dust has settled.

# STORAGE AND DISPOSAL

**STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.

### DISPOSAL

If Empty—Do not reuse this container. Place in trash or offer for recycling if available.

If Partly Filled—Call your local solid waste agency or toll free number (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.

# **USE PRECAUTIONS:**

Do not use on dichondra, carpetgrass, closely mowed bentgrass putting greens or St. Augustinegrass. Do not use this product where desirable clovers are present or allow the material to drift onto flowers, vegetables or ornamental shrubbery. Do not spread over exposed roots of trees and ornamentals. Do not exceed specified dosages for any area; be particularly careful within dripline of tree and other ornamental species. Apply when the air is calm. Application of this product to newly seeded areas should be delayed until the third mowing. Do not contaminate domestic or irrigation waters.

# INSTRUCTIONS:

Gordon's Trimec® Weed & Feed 30 contains Trimec® herbicide. Trimec® controls a wide range of lawn weeds, including dandelion, chickweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a (X-X-X analysis of fertilizer grade) fertilizer provides a supply of plant food. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days. Treated areas may be seeded 3 to 4 weeks after application.

**Note to Supplemental Registrants:** Guarantees for plant nutrients other than nitrogen, phosphorous, and potassium may be permitted or required by regulation by the \_\_\_. The guarantees for such other nutrients shall be expressed in the form of the element. The source (oxides, salts, chelates, etc.) of such other nutrients may be required to be stated and may be included on the label.

# For state registrations in Kentucky, add the following paragraph:

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing (\*Number of pounds) of nitrogen per 1,000 square feet. Please consult your local extension office for questions regarding a regular fertilizer program suitable for your geographic region.

# Labeling Requirements:

- All fertilizer products distributed in Washington at a minimum, must have the following labeling statement: "This product has been registered with the Washington State Department of Agriculture. When applied as directed, this fertilizer meets the Washington standards for arsenic, cadmium, cobalt, mercury, molybdenum, lead, nickel, selenium, and zinc. You have the right to receive specific information about Washington standards from the distributor of this product."
- After July 1, 1999, the label must also state: "Information received by the Washington State
  Department of Agriculture regarding the components in this product is available on the internet at
  <a href="http://www.wa.gov/agr/">http://www.wa.gov/agr/</a>."
- -OR- Information regarding the contents and levels of metals in this product is available on the internet at http://www.regulatory-info-ba.com
- -OR- Information regarding the contents and levels of metals in this product, is available on the Internet as <a href="http://www.aapfco.org/metals.htm">http://www.aapfco.org/metals.htm</a>.

The maximum application rate to turf is 1.2 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

- 1. Mow lawn to normal height 1 to 2 days before application.
- 2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
- Apply when weeds are young and actively growing, preferably in the morning when the dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants.
- Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

- WEEDS -				
Annual yellow sweetclover	Cockle	Musk thistle	Spiny amaranth	
Aster	Cocklebur	Mustard	Spiny cocklebur	
Austrian fieldcress	Common mullein	Narrowleaf plantain	Spiny sowthistle	
Bedstraw	Creeping Charlie	Narrowleaf vetch	Spotted catsear	
Beggarticks	Creeping jenny	Nettle	Spotted spurge	
Betony, Florida	Cudweed	Orange hawkweed	Spurweed	
Bindweed, field	Curly dock	Oriental cocklebur	Stinging nettle	
Bird vetch	Daisy, English	Oxalis	Strawberry, India mock	
Bitter wintercress	Daisy fleabane	Parsley-piert	Tall nettle	
Bittercress, hairy	Daisy, oxeye	Parsnip	Tall vervain	
Bitterweed	Dandelion	Pearlwort	Tansy ragwort	
Black-eyed Susan	Dichondra	Pennycress	Tansy mustard	
Black medic	Dogbane	Pennywort	Tanweed	
Black mustard	Dogfennel	Peppergrass (Peppergrass)	Thistle	
Blackseed plantain	Dollarweed	Pepperweed	Trailing crownvetch	
Blessed thistle	Elderberry	Pigweed	Tumble mustard	
Bloodflower milkweed	False dandelion	Pineywoods bedstraw	Tumble pigweed	
Blue lettuce	False flax	Plains coreopsis (tickseed)	Velvetleaf	
Blue vervain	False sunflower	Plantain	Venice mallow	
Bracted plantain	Fiddleneck	Poison ivy	Virginia buttonweed	
Bristly oxtongue	Florida pusley	Poison oak	Virginia creeper	
Broadleaf dock	Frenchweed	Pokeweed	Virginia pepperweed	
Broadleaf plantain	Galinsoga	Poorjoe	Water pennywort	
Broomweed	Goldenrod	Prairie sunflower	Wavyleaf bullthistle	
Buckhorn	Ground ivy	Prickly lettuce	Western clematis	
Buckhorn plantain	Gumweed	Prickly sida	Western salsify	
Bulbous buttercup	Hairy fleabane	Prostrate knotweed	White mustard	
Bull thistle	Hawkweed	Prostrate pigweed	Wild aster	
Bullnettle	Healall	Prostrate spurge	Wild buckwheat	
Burclover	Heartleaf drymary	Prostrate vervain	Wild carrot	
Burdock	Heathaster	Puncturevine	Wild four-o'clock	
Burning nettle	Hedge bindweed	Purslane, common	Wild garlic	
Burweed	Hedge mustard	Ragweed	Wild geranium	
Buttercup	Henbit	Red sorrel	Wild lettuce	
Buttonweed	Hoary cress	Redroot pigweed	Wild marigold	
Canada thistle	Hoary plantain	Redstem filaree	Wild mustard	
Carolina geranium	Hoary vervain	Rough cinquefoil	Wild onion	
Carpetweed	Horsenettle	Rough fleabane	Wild parsnip	
Catchweed bedstraw	Jimsonweed	Russian pigweed	Wild radish	
Catnip	Knawel	Russian thistle	Wild rape	
Catsear	Knotweed	Scarlet pimpernel	Wild strawberry	
Chickweed, common	Kochia	Scotch thistle	Wild sweet potato	
Chickweed, mouseear	Lambsquarters	Sheep sorrel	Wild vetch	
Chicory	Lespedeza	Shepherdspurse	Woodsorrel	
Cinquefoil	Mallow	Slender plantain	Woolly croton	
Clover, crimson	Matchweed	Smallflower galinsoga	Woolly morningglory	
Clover, hop	Mexicanweed	Smooth dock	Woolly plantain	
Clover, red	Milk vetch	Smooth pigweed	Wormseed	
Clover, strawberry	Morningglory	Sorrel	Yarrow	
Clover, sweet	Mouseear hawkweed	Sowthistle	Yellow rocket	
Clover, white	Mugwort	Spanishneedles	Yellowflower pepperweed	
·	Ç ·	Speedwell	and other broadleaf weeds	

# SPREADERS AND CALIBRATION:

- 1. Drop spreaders and rotary spreaders must be calibrated for uniform application of this product.
- 2. Close spreader opening (orifice).
- 3. Adjust spreader setting using the Spreader Suggestions chart

Application Rate = 3 lbs. of product per 1,000 square feet.

# SUGGESTED SPREADER SETTINGS:


Centrifugal/Rotary Spreaders	Application Rate, 3 lbs./1,000 sq.ft.
Frank's Broadcast	2 1/4
Scotts Speedy Green	5
Radio Flyer #0047	8 ¼
True Temper Rotary CB2000	3 ¾
Earthway Rotary #2050SU	14 ½
Drop Spreaders	Application Rate, 3 lbs./1,000 sq.ft.
Frank's Drop	8 1/4
Greenview II	5 ½
True Temper CD20	4 - 4 1/2
Scotts Accugreen AG1	5 ½
Earthway Drop #7300SU	13 ½

These spreader settings are approximate. Differences may occur due to the condition of spreader, speed at which applied and roughness of area treated. Check your spreader on 330 square feet and adjust to apply at the rate of 1 pound per 1,000 square feet.


# 4. CALIBRATION INSTRUCTIONS:

- For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 330 square feet and adjust to apply at the rate of 1 pound of product per 330 square feet.
- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.
- 5. Fill spreader. Fill spreader on driveway or sidewalk to eliminate the chance of spillage on the lawn,
- 6. Apply the product to lawn. Refer to the patch charts below:

Path A for centrifugal/rotary spreaders.




Path B for drop spreaders.




- 7. Close the spreader opening when turning to change directions or when stopping.
- 8. If using a centrifugal/rotary spreader, avoid spreading product into adjacent landscape, flowers or vegetable plantings.

NOTE: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 square feet.

This \_\_\_ pound bag should cover approximately \_\_\_\_ square feet (\_\_\_ ft. x \_\_\_ ft.) Refer to the table and picture to determine the size of your lawn.





24-4-8 GUARANTEED ANALYSIS	
Total Nitrogen (N)	24%
Water Insoluble Nitrogen	5%
Available Phosphate (P2O5)	4%
Soluble Potash (K <sub>2</sub> O)	8%

**Note to Supplemental Registrants:** Fertilizer sources may vary in micronutrient composition, and iron may discolor or stain concrete. For fertilizers containing iron: Sweep all fertilizer from concrete surfaces immediately after application. Do not wash this product from concrete surfaces because fertilizers which contain iron may cause discoloration upon oxidation.

- WEEDS -				
Bedstraw	Knotweed	Ragweed		
Black medic	Lambsquarters	Sheep sorrel		
Buckhorn	Lespedeza	Shepherdspurse		
Burdock	Mallow	Speedwell		
Chickweed	Morningglory	Spurge		
Chicory	Oxalis	Thistle		
Clover	Peppergrass	Wild carrot		
Dandelion	Pigweed	Wild garlic		
Dock	Plantain	Wild lettuce		
Ground ivy	Poison ivy	Wild onion		
Healall	Poison oak	Yarrow		
Henbit	Purslane			

# LIMITED WARRANTY AND DISCLAIMER.

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once to the seller.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

# Sublabel 2 - For Supplemental Registrations to Bayer Advanced LLC. Distributor Company Number 72155

Distributor Product Label:

Bayer Advanced Lawn<sup>TM</sup> Triple Action Lawn Fertilizer Plus Weed Control<sup>il</sup>

# PDF File of proposed container label follows:




NET WT 16 LBS (7.2kg)

KEEP DUT OF REACH OF CHILDREN CAUTION Serback point for CAUTION Serback point for CAUTION Serback of CAUTION SERBACK OF CAUTION OF C

THIS PRODUCT CONTRIPUT:
128 hs. 240 aced on inderthes for
30 hs. Mosaprop and appealed parting
14 hs. Throathe aced on indept per for

# Covers 5,000 Seems ---



. . . . . . .

)

# **APPENDIX**

# II. Previously Approved Fertilizer Grades:

◆ Fertilizer grades with nutrient contents expressed as the percentages of N,P₂O₅, and K₂O include the following:

10-6-4	16-3-5	20-10-0-(1)	23-7-7	25-4-8	28-3-8	32-4-8
10-16-20	16-6-9	20-10-5	24-3-7	26-3-3	29-3-4	
10-18-10	18-3-5	21-5-7	24-4-8	26-3-5	30-3-4	
12-5-5	18-6-12	22-6-6	25-3-3	26-4-6	30-3-5	
15-5-10	20-4-6	22-7-7	25-3-5	28-3-3	30-3-10	
16-2-3	20-4-8	22-10-10	25-3-8	28-3-7	30-4-8	

The fertilizer grade guaranteed analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations. The following guaranteed analyses presentations have previously been submitted as notifications:

10-16-20 GUARANTEED ANALYSIS		
Total Nitrogen (N)	10%	
6.0% Ammoniacal Nitrogen		
4.0% Urea Nitrogen		
Available Phosphate (P <sub>2</sub> O <sub>5</sub> )	16%	
Soluble Potash (K <sub>2</sub> O)	20%	
Magnesium (Mg)	1.0%	
0.1% Water Soluble Magnesium (Mg)		
Sulfur (S)	2.0%	
2.0% Combined Sulfur (S)		
Iron (Fe)	1.0%	
Manganese (Mn)	0.5%	
0.1% Water Soluble Manganese (Mn)		
Nutrient Sources: Urea, Ammonium Phosphate, Ammonium Sulfate, Muriate of	Potash,	
Ferrous Sulfate, Manganese Oxide, Manganese Sulfate, Sulfate of Potash Magne	sia and	
Magnesium oxide.		

29-3-4 GUARANTEED ANALYSIS	
Total Nitrogen (N)	29%
1.1% Ammoniacal Nitrogen	
27.9% Urea Nitrogen	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> )	3%
Soluble Potash (K <sub>2</sub> O)	4%
Derived from: Polymer-coated urea, urea, diammonium phosphate, and muriate of p	otash.
*11.0% slowly available urea nitrogen from polymer coated urea.	
Chlorine (Cl), note more than	4.0%

10-18-10 GUARANTEED ANALYSIS	
Total Nitrogen (N)	10%
10.0% Ammoniacal Nitrogen	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> )	18%
Soluble Potash (K <sub>2</sub> O)	10%
Sulfur (S)	2.0%
5.0% Combined Sulfur (S)	
Nutrient Sources: Ammonium Sulfate, Ammonium Phosphate and Muriate of Potash.	

26-4-6 GUARANTEED ANALYSIS	
Total Nitrogen (N)	26%
1.8% Ammoniacal Nitrogen	
24.4% Urea Nitrogen	
Available Phosphoric Acid(P <sub>2</sub> O <sub>5</sub> )	4%
Soluble Potash (K <sub>2</sub> O)	6%
Boron (B) 0.02%	
Copper (Cu)	
Iron (Fe)	
Total Manganese (Mn)	
Molybenum (Mo)	
Zinc (Zn)	
Primary Plant Food Sources: Sulfur coated urea, urea, diammonium phosphate, mu potash.	riate of
Secondary Plant Food Sources: Fertilizer grade borate, sulfates and oxides of sulfates and oxides of iron, oxides of manganese, fertilizer grade molybdate, and and oxides of zinc.	
*4.0% slowly available urea nitrogen from sulfur coated urea.	
Chlorine (Cl), not more than	4.8%

28-3-7 GUARANTEED ANALYSIS	
Total Nitrogen (N)	28%
1.2% Ammoniacal Nitrogen	
26.8% Urea Nitrogen	
Available Phosphoric Acid(P <sub>2</sub> O <sub>5</sub> )	3%
Soluble Potash (K <sub>2</sub> O)	7%
Primary Plant Food Sources: Polymer-coated urea, urea, diammonium phosphate, mu potash	riate of
*7.0% slowly available urea nitrogen from polymer-coated urea.	
Chlorine (Cl), not more than	6.4%

# III. Advertising claims that may be presented on the retail containers:

- (Fifteen, Sixteen, Eighteen or Twenty) Pounds Covers Up To 5,000 Square Feet.
- Controls/Kills dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf weeds; some of which are listed on the back panel
- ◆ An 18-6-12 fertilizer provides slow-release feeding to promote winter hardiness and drought tolerance in lawns.
- Feeds your lawn for up to 12 weeks.
- Controls/Kills over 30 varieties of weeds while feeding your lawn.
- Greens lawns and develops deep roots while controlling weeds.
- Bring out the best with the turf magic touch!
- ◆ Controls/Kills dandelion, plantain, and other weeds.
- Feeds the lawn with two (2) types of nitrogen and is pelleted for easy application.
- Controls (Number) weeds! (Listing of weeds may include the following.)

- · Kills dandelion, chickweed, clover and most common lawn weeds.
- Triple Action Feeding works at every level.

Grass- Greener, Thicker lawn

Roots - Stronger Root System

Soil - Enriches & Fortifies Soil

- Builds a better lawn Guaranteed
- Gordon's Trimec Weed & Feed 30 kills unsightly weeds such as dandelion, chickweed and clover
  while it nourishes your lawn. Its Triple Action formula contains slow-release nitrogen which
  continuously feeds the grass, roots and soil for up to two full months [all winter long]. And it's
  guaranteed not to burn. The result is a greener, thicker lawn [this Fall and quicker green-up next
  Spring]. You're just one application away from a better lawn. That's the [company name] Better
  Lawn Guarantee.
- Triple Action Feeding works at every level for a Deep-Down Green!
- Why is Triple Action better? By feeding at all levels, your lawn receives the primary nutrients it needs to thrive. This ensures thicker, greener grass, stronger root system, and richer soil the key to building a lush lawn that lasts.
- Grass: Builds thicker, greener grass and enhances overall appearance.
- Roots: Builds stronger, deeper root systems which transport nutrients from the soil to the grass to help lawn withstand heat-stress and drought.
- Soil: Enriches and fortifies your soil. Rich, fertile soil provides a foundation for establishing roots and replenishes vital primary nutrients.
- For questions or comments, call toll-free 1-877-bayerAL (1-877-229-3725)
- · Made in the U.S.A.
- Kills weeds and builds a thick, green Fall lawn.
- Nourishes all Winter for a quick Spring green-up.



# GORDON'S TRIMEC® WEED & FEED 30 24-4-8

Sublabel 3: For use by supplemental distributor for Bayer Advanced Triple Action Winterizer Fertilizer Plus Weed Control "

# **ACTIVE INGREDIENTS:**

2,4-D (CAS No. 94-75-7)	0.69%
Mecoprop-p (CAS No. 16484-77-8)	0.15%
Dicamba (CAS No. 1918-00-9)	0.06%
INERT INGREDIENTS:	99.10%
TOTAL	100.00%

# THIS PRODUCT CONTAINS:

13.8 lbs. 2,4-D acid equivalent per ton
3.0 lbs. Mecoprop-p acid equivalent per ton
1.2 lbs. Dicamba acid equivalent per ton
Isomer Specific by AOAC Methods.
Contains the single isomer form of Mecoprop-p.
TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

# **KEEP OUT OF REACH OF CHILDREN**

# CAUTION

See back panel for Precautionary Statements.

NET WEIGHT: (15, 16, 18 or 20) POUNDS

575 AP032107 EPA REG. NO. 2217-579 EPA EST. NO. 2217-KS-X MANUFACTURED BY:

An Employee Owned Company
1217 West 12th Street
Kansao City, Miss.cum 64101

Telephone: 1-800-821-7925



# READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

# PRECAUTIONARY STATEMENTS

# **Hazards to Humans & Domestic Animals**

**CAUTION:** Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

# First Aid

	وي منزوا من الموادل المراجع الموادية في الموادية من الموادية الموادية الموادية الموادية والموادية الموادية المو			
If in eyes:	<ul> <li>Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>			
If on skin or	Take off contaminated clothing.			
clothing:	Rinse skin immediately with plenty of water for 15 - 20 minutes.			
	Call a poison control center or doctor for treatment advice.			
if swallowed:	Call a poison control center or doctor immediately for treatment advice.			
•	Have person sip a glass of water if able to swallow.			
	<ul> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> </ul>			
	Do not give anything by mouth to an unconscious person.			
Have the product container or label with you when calling a poison control center or doctor or going for treatment.				
You may also ca	ill toll-free, 1-877-800-5556 for emergency medical treatment information.			

# **ENVIRONMENTAL HAZARDS:**

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater. Do not apply when weather conditions favor drift from target area.

# **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until dust has settled.

# STORAGE AND DISPOSAL

**STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.

# **DISPOSAL:**

If Empty—Do not reuse this container. Place in trash or offer for recycling if available.

**If Partly Filled**—Call your local solid waste agency or toll free number (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.

# **USE PRECAUTIONS:**

Do not use on dichondra, carpetgrass, closely mowed bentgrass putting greens or St. Augustinegrass. Do not use this product where desirable clovers are present or allow the material to drift onto flowers, vegetables or ornamental shrubbery. Do not spread over exposed roots of trees and ornamentals. Do not exceed specified dosages for any area; be particularly careful within dripline of tree and other ornamental species. Apply when the air is calm. Application of this product to newly seeded areas should be delayed until the third mowing. Do not contaminate domestic or irrigation waters.

# **INSTRUCTIONS:**

Gordon's Trimec<sup>∞</sup> Weed & Feed 30 contains Trimec<sup>∞</sup> herbicide. Trimec<sup>∞</sup> controls a wide range of lawn weeds, including dandelion, chickweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a (X-X-X analysis of fertilizer grade) fertilizer provides a supply of plant food. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days. Treated areas may be seeded 3 to 4 weeks after application.

Note to Supplemental Registrants: Guarantees for plant nutrients other than nitrogen, phosphorous, and potassium may be permitted or required by regulation by the \_\_\_\_. The guarantees for such other nutrients shall be expressed in the form of the element. The source (oxides, salts, chelates, etc.) of such other nutrients may be required to be stated and may be included on the label.

# For state registrations in Kentucky, add the following paragraph:

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing (\*Number of pounds) of nitrogen per 1,000 square feet. Please consult your local extension office for questions regarding a regular fertilizer program suitable for your geographic region.

# Labeling Requirements:

- All fertilizer products distributed in Washington at a minimum, must have the following labeling statement: "This product has been registered with the Washington State Department of Agriculture. When applied as directed, this fertilizer meets the Washington standards for arsenic, cadmium, cobalt, mercury, molybdenum, lead, nickel, selenium, and zinc. You have the right to receive specific information about Washington standards from the distributor of this product."
- After July 1, 1999, the label must also state: "Information received by the Washington State
  Department of Agriculture regarding the components in this product is available on the internet at
  <a href="http://www.wa.qov/agr/">http://www.wa.qov/agr/</a>."
- -OR- Information regarding the contents and levels of metals in this product is available on the internet at <a href="http://www.regulatory-info-ba.com">http://www.regulatory-info-ba.com</a>
- -OR- Information regarding the contents and levels of metals in this product, is available on the Internet as <a href="http://www.aapfco.org/metals.htm">http://www.aapfco.org/metals.htm</a>.

The maximum application rate to turf is 1.2 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

- Mow lawn to normal height 1 to 2 days before application.
- 2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
- Apply when weeds are young and actively growing, preferably in the morning when the dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants.
- 4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

	- WEEDS -				
Annual yellow sweetclover	Cockle	Musk thistle	Spiny amaranth		
Aster	Cocklebur	Mustard	Spiny cocklebur		
Austrian fieldcress	Common mullein	Narrowleaf plantain	Spiny sowthistle		
Bedstraw	Creeping Charlie	Narrowleaf vetch	Spotted catsear		
Beggarticks	Creeping jenny	Nettle	Spotted spurge		
Betony, Florida	Cudweed	Orange hawkweed	Spurweed		
Bindweed, field	Curly dock	Oriental cocklebur	Stinging nettle		
Bird vetch	Daisy, English	Oxalis	Strawberry, India mock		
Bitter wintercress	Daisy fleabane	Parsley-piert	Tall nettle		
Bittercress, hairy	Daisy, oxeye	Parsnip	Tall vervain		
Bitterweed	Dandelion	Pearlwort	Tansy ragwort		
Black-eyed Susan	Dichondra	Pennycress	Tansy mustard		
Black medic	Dogbane	Pennywort	Tanweed		
Black mustard	Dogfennel	Peppergrass (Peppergrass)	Thistle		
Blackseed plantain	Dollarweed	Pepperweed	Trailing crownvetch		
Blessed thistle	Elderberry	Pigweed	Tumble mustard		
Bloodflower milkweed	False dandelion	Pineywoods bedstraw	Tumble pigweed		
Blue lettuce	False flax	Plains coreopsis (tickseed)	Velvetleaf		
Blue vervain	False sunflower	Plantain	Venice mallow		
Bracted plantain	Fiddleneck	Poison ivy	Virginia buttonweed		
Bristly oxtongue	Florida pusley	Poison oak	Virginia creeper		
Broadleaf dock	Frenchweed	Pokeweed	Virginia pepperweed		
Broadleaf plantain	Galinsoga	Poorjoe	Water pennywort		
Broomweed	Goldenrod	Prairie sunflower	Wavyleaf bullthistle		
Buckhorn	Ground ivy	Prickly lettuce	Western clematis		
Buckhorn plantain	Gumweed	Prickly sida	Western salsify		
Bulbous buttercup	Hairy fleabane	Prostrate knotweed	White mustard		
Bull thistle	Hawkweed	Prostrate pigweed	Wild aster		
Bullnettle	Healall	Prostrate spurge	Wild buckwheat		
Burclover	Heartleaf drymary	Prostrate vervain	Wild carrot		
Burdock	Heathaster	Puncturevine	Wild four-o'clock		
Burning nettle	Hedge bindweed	Purslane, common	Wild garlic		
Burweed	Hedge mustard	Ragweed	Wild geranium		
Buttercup	Henbit	Red sorrel	Wild lettuce		
Buttonweed	Hoary cress	Redroot pigweed	Wild marigold		
Canada thistle	Hoary plantain	Redstem filaree	Wild mustard		
Carolina geranium	Hoary vervain	Rough cinquefoil	Wild onion		
Carpetweed	Horsenettle	Rough fleabane	Wild parsnip		
Catchweed bedstraw	Jimsonweed	Russian pigweed	Wild radish		
Catnip	Knawel	Russian thistle	Wild rape		
Catsear	Knotweed	Scarlet pimpernel	Wild strawberry		
Chickweed, common	Kochia	Scotch thistle	Wild sweet potato		
Chickweed, mouseear	Lambsquarters	Sheep sorrel	Wild vetch		
Chicory	Lespedeza	Shepherdspurse	Woodsorrel		
Cinquefoil	Mallow	Slender plantain	Woolly croton		
Clover, crimson	Matchweed	Smallflower galinsoga	Woolly morningglory		
Clover, hop	Mexicanweed	Smooth dock	Woolly plantain		
Clover, red	Milk vetch	Smooth pigweed	Wormseed		
Clover, red Clover, strawberry	Morningglory	Sorrel	Yarrow		
Clover, strawberry Clover, sweet	Mouseear hawkweed	Sowthistle			
			Yellow rocket		
Clover, white	Mugwort	Spanishneedles	Yellowflower pepperweed		
	<del></del>	Speedwell	and other broadleaf weeds		

# SPREADERS AND CALIBRATION:

- 1. Drop spreaders and rotary spreaders must be calibrated for uniform application of this product.
- 2. Close spreader opening (orifice).
- 3. Adjust spreader setting using the Spreader Suggestions chart

Application Rate = 3 lbs. of product per 1,000 square feet.

# SUGGESTED SPREADER SETTINGS:

Centrifugal/Rotary Spreaders	Application Rate, 3 lbs./1,000 sq.ft.
Frank's Broadcast	2 1/4
Scotts Speedy Green	5
Radio Flyer #0047	8 1/4
True Temper Rotary CB2000	3 ¾
Earthway Rotary #2050SU	14 ½
Drop Spreaders	Application Rate, 3 lbs./1,000 sq.ft.
Frank's Drop	8 1/4
Greenview II	5 1/2
True Temper CD20	4 - 4 1/2
Scotts Accugreen AG1	5 1/2
Earthway Drop #7300SU	13 ½

These spreader settings are approximate. Differences may occur due to the condition of spreader, speed at which applied and roughness of area treated. Check your spreader on 330 square feet and adjust to apply at the rate of 1 pound per 1,000 square feet.


# 4. CALIBRATION INSTRUCTIONS:

- For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 330 square feet and adjust to apply at the rate of 1 pound of product per 330 square feet.
- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.
- 5. Fill spreader. Fill spreader on driveway or sidewalk to eliminate the chance of spillage on the lawn.
- 6. Apply the product to lawn. Refer to the patch charts below:

Path A for centrifugal/rotary spreaders.




Path B for drop spreaders.




- 7. Close the spreader openings when turning to change directions or when stopping.
- 8. If using a centrifugal/rotary spreader, avoid spreading product into adjacent landscape, flowers or vegetable plantings.

NOTE: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 square feet.

This \_\_\_ pound bag should cover approximately \_\_\_\_ square feet (\_\_\_ ft. x \_\_\_ ft.) Refer to the table and picture to determine the size of your lawn.





24-4-8 GUARANTEED ANALYSIS		
Total Nitrogen (N)	24%	
Water Insoluble Nitrogen	5%	
Available Phosphate (P2O5)	4%	
Soluble Potash (K <sub>2</sub> O)	8%	

**Note to Supplemental Registrants:** Fertilizer sources may vary in micronutrient composition, and iron may discolor or stain concrete. For fertilizers containing iron: Sweep all fertilizer from concrete surfaces immediately after application. Do not wash this product from concrete surfaces because fertilizers which contain iron may cause discoloration upon oxidation.

- WEEDS -			
Bedstraw	Knotweed	Ragweed	
Black medic	Lambsquarters	Sheep sorrel	
Buckhorn	Lespedeza	Shepherdspurse	
Burdock	Mallow	Speedwell	
Chickweed	Morningglory	Spurge	
Chicory	Oxalis	Thistle	
Clover	Peppergrass	Wild carrot	
Dandelion	Pigweed	Wild garlic	
Dock	Plantain	Wild lettuce	
Ground ivy	Poison ivy	Wild onion	
Healall	Poison oak	Yarrow	
Henbit	Purslane		

# LIMITED WARRANTY AND DISCLAIMER.

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once to the seller.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

# Sublabel 3 - For Supplemental Registrations to Bayer Advanced LLC, Distributor Company Number 72155

Distributor Product Label:

Bayer Advanced TM Triple Action Winterizer Fertilizer plus Weed Control<sup>11</sup>

# PDF File of proposed container label follows:



NET WT 18 LBS (8.1kg)

KEEP OUT OF REACH OF CHILDREN CAUTION See heat proof for procountries; statements



THE PRODUCT CONTAINS: A9 - 04 USB line, 3 fed a city op a milest product of the 12 feb. March pop a cit in prime large per ten 12 feb. (Countain a cit one insulation per ten insulation per ten insulation special to y a OM. (Methods. Communities angle somer insulation per ten ins

# Covers 5,000 Square Feet



1

1

١

1

1

1

ĺ

Winters Wood Commo!

This product left is employ come with contracted in a thirteen which it is not there where the common common with the common which contracted product in the common common which contracted product in common common which contracted product is common the contract. The contract is conserved their forest the forest half and one that of contracted the contracted product is contracted to contract in the contracted product in the contracted contracted to contract in the contracted contracted contracted to contract in the contracted contracted

READY-TO-SPREAD CRANULES

Imple Action Feeding works at every level for a Deep Down Green! 134 With h Bryko Ariban Lacker? By briding an all levels your laws marriers the princip named it made he through the community that of, growing game supergram most systems, and define will—the day to healthy clear that have





Carrie additions promy gamen interpresent

Bullik tampet desper kat synoto which purspers may an have de godin the great solephysist level safetand (per great and despite midmam lintika som vol. Helt i nji k och prantse e kombotur n suchfalingens and nybodow viad prantse en men



CONTRACTOR OF CONTRACTOR

# ENRECTIONS FOR USE

It is a valution of Federal law to use this product in a manner more state with its labeling

# READ THE ENTIRE LABEL FIRST.

YEDS:	* <b>₹</b>					
APR Will LINESSON	ER areti	Constant	L 1 Later L 1 K	3 30	14 - ACT HER D.	
LEE KEEL	2122 2 2 4m. 22*******	E CAMPE E CAMPH	4 TE - 115 B	to a the	34.00	
Giffyr -	Chilley restrey his	3.0	SEL SHORTER	A. P. Salvana	rate legal.	
्राचित्र के बीतिया संक्रिकेट स्थापन	Table 1 (MAC)	. (1) (1) (2) . (1) (2) (3) . (1) (4) (4)	ja Tiga (, Saba)) 15 na distribution	० सर्वे । इस्ते ० म्योचन इस्त्राज्ञ ० वेदर विशेषक	Total State of the Control of the Co	
n K	.J.	ALS ES HATS	4. For	e service and	201 BF 224	
1111 401	ं असे " चला , श्राप्त द	e is	न । के कहा न है कि कर	1	- 23 - 24 s - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1	
1. 1. February	, and (a)	me and the state of the state o	Treasure Leaves	Charte Charte		
ii bildune	81 1949 11 Mars	απι. 7 ε απη: Α-χε. χηνι <b>α</b> νι	15ki	K wants	Congression of the Congression o	
1124	100	- AT 188	447 E	tart merte	1 1 1000	
i ja	1200 € 2000 1000 € 2000 1000 € 1000	Carrier Carrier	The party of the state of the s	TALEST SAFE	The state of the s	
ine		F. 100	Cres Sunsee	1 2	¥.3	
parameter (1918) 1	10 May 1	720-8 57-4781 57-4781	NATE OF THE PARTY	ET LIST BY	i marija i marijani 1 marija i marijani 1 marija i marijani	
Late &	( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( )	2 term	Carried Comments	5	ACT WEST	
4 2 % 4 1 - 2 4 (m.) 10 kg	in the same	200 annes 20 annes 10 annes	Mar.	<b>本体:子子等 1</b> 3	A STATE OF STATE OF	
45.4	io alani Sai	an off Budson	्र केट्र वे <b>ट्रिका</b> २ केट्र हे सिक्क	T. CALL	***	

SPREADER SETTINGS	3
Contribution from T	Application of Page
A STATE OF THE STA	15 to 1200scal
CHATRO ROSSAN	3
THE RESIDENCE HEAT PARTY AND ADDRESS.	:
BELLEVAL BOOMS	:
HELEFIE & BOOKEN ASSE	2
AUCAL? PAREASMAN DAUGHOU	1.
THE MEDICAL MACHINE	1
THE STREET BY COM	4
Comp Salaucis III	Applicable   Rate
	No tage
CURYCUME.	ı .
報的可能: (7 DM P )(433)	,
COTALICATED MINESO	44
<b>公司在从公司通過 (10)</b>	Ļ
ECTES DEEP (ANY TICE PER TH)	111



# USE PRECAUTIONS

USE PRECAUTIONS
De les une sonicidante authory un
les et une é l'empare, parier yeurs de
l'équatempais. De dataine s'ét parier
l'équatempais. De dataine s'étaine s'étaine de l'empartie d'étaine s'étaine auté un sone de parier l'étaine s'étaine auté un sone de parier l'étaine s'étaine s'étaine s'étaine s'étaine s'étaine auté un sone de parier l'étaine s'étaine s'étaine

UNCORNING MINISTER

of based cut approach in just to mean the life. See and 1. Show here to recreate the glot to a serve to be seen as placed as a long to be seen as placed as a long to the large and to the large and large

্যালনামা হাত্ৰা বিষয় হৈছে ক্ষমান্ত না চুক্তাক ব বিভাগৰ ভ কবিন্ত ক্ষমান্ত না কিছিল কৰিছে ই বাহন কৰিছে বাহুকি ক্ষম ব বিশ্বাস্থা চয়ত ত ক্ষমান্ত কৰিছে বাহুকি কৰাৰ কৈ ক্ষমান্ত

# PRECAUTIONARY STATEMENTS AND ENVIRONMENTAL HAZARDS

# PRECAUTIONARY STATEMENTS Hazarda to Humans and Demonic Animais CABTION

ener i i um ការបានក្រុម ការបានិការបានការបានការបានការបានការបានការបានការបានការបានការបានការបានការ

ும் நிறு இது நிறுந்திருள்ளது. இது இருவின் நிறிந்த நார்ந்து நிருந்து நிறுந்திருள்ளது. இது நார்ந்திருந்து இது இருக்கு FIRST AID +


\* Transcondition ( ) 本面的版 ・ 「中午」 「CLUSTON ( ) 」」 ・ 「CLUSTON ( ) 」」 ・ 「CLUSTON ( ) 「CLUSTON

The state of the properties of the state of

# STORAGE AND DISPOSAL

関連機能 (A delta winda yorana and none atthem (カンカル) (A delta スポーティス delta tipe and and (-and ) ( an (A delta スポーティス delta tipe and and (-and ) ( an (A delta ) ( A delta della della della delta delta delta della della della della della della della della dell

- 東京の大学 (東京の大学) (東京の



# GUARANTEED ANALYSIS: 22-3-14

The loss specified to the control of AND ALL TO BE AND THE AND T

file to a country of Glorian in the STA (Fig. 1) and the state of Glorian in t NOTED HOST A THREE TO

Berga 🏶 se nast tasse at 1 4



ļ

1

ţ

14.1128.3514.1

# **APPENDIX**

# III. Previously Approved Fertilizer Grades:

• Fertilizer grades with nutrient contents expressed as the percentages of N,P<sub>2</sub>O<sub>5</sub>, and K<sub>2</sub>O include the following:

10-6-4	16-3-5	20-10-0-(1)	23-7-7	25-4-8	28-3-8	32-4-8
10-16-20	16-6-9	20-10-5	24-3-7	26-3-3	29-3-4	
10-18-10	18-3-5	21-5-7	24-4-8	26-3-5	30-3-4	
12-5-5	18-6-12	22-6-6	25-3-3	26-4-6	30-3-5	
15-5-10	20-4-6	22-7-7	25-3-5	28-3-3	30-3-10	
16-2-3	20-4-8	22-10-10	25-3-8	28-3-7	30-4-8	

The fertilizer grade guaranteed analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations. The following guaranteed analyses presentations have previously been submitted as notifications:

10-16-20 GUARANTEED ANALYSIS	
Total Nitrogen (N)	10%
6.0% Ammoniacal Nitrogen	
4.0% Urea Nitrogen	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> )	16%
Soluble Potash (K <sub>2</sub> O)	20%
Magnesium (Mg)	1.0%
0.1% Water Soluble Magnesium (Mg)	
Sulfur (S)	2.0%
2.0% Combined Sulfur (S)	
Iron (Fe)	1.0%
Manganese (Mn)	0.5%
0.1% Water Soluble Manganese (Mn)	
Nutrient Sources: Urea, Ammonium Phosphate, Ammonium Sulfate, Muriate of	Potash,
Ferrous Sulfate, Manganese Oxide, Manganese Sulfate, Sulfate of Potash Magne	sia and
Magnesium oxide.	

29-3-4 GUARANTEED ANALYSIS		
Total Nitrogen (N)	29%	
1.1% Ammoniacal Nitrogen		
27.9% Urea Nitrogen		
Available Phosphate (P <sub>2</sub> O <sub>5</sub> )	3%	
Soluble Potash (K <sub>2</sub> O)	4%	
Derived from: Polymer-coated urea, urea, diammonium phosphate, and muriate of	ootash.	
*11.0% slowly available urea nitrogen from polymer coated urea.		
Chlorine (Cl), note more than	4.0%	

10-18-10 GUARANTEED ANALYSIS		
Total Nitrogen (N)	10%	
10.0% Ammoniacal Nitrogen		
Available Phosphate (P <sub>2</sub> O <sub>5</sub> )	18%	
Soluble Potash (K <sub>2</sub> O)	10%	
Sulfur (S)	2.0%	
5.0% Combined Sulfur (S)		
Nutrient Sources: Ammonium Sulfate, Ammonium Phosphate and Muriate of Potash.		

26-4-6 GUARANTEED ANALYSIS	
Total Nitrogen (N)	26%
1.8% Ammoniacal Nitrogen	
24.4% Urea Nitrogen	
Available Phosphoric Acid(P <sub>2</sub> O <sub>5</sub> )	4%
Soluble Potash (K <sub>2</sub> O)	6%
Boron (B) 0.02%	
Copper (Cu)	
Iron (Fe)	
Total Manganese (Mn)	
Molybenum (Mo)	
Zinc (Zn) 0.05%	
Primary Plant Food Sources: Sulfur coated urea, urea, diammonium phosphate, mur potash.	riate of
Secondary Plant Food Sources: Fertilizer grade borate, sulfates and oxides of could sulfates and oxides of iron, oxides of manganese, fertilizer grade molybdate, and sund oxides of zinc.	
*4.0% slowly available urea nitrogen from sulfur coated urea.	
Chlorine (Cl), not more than	4.8%

28-3-7 GUARANTEED ANALYSIS	
Total Nitrogen (N)	28%
1.2% Ammoniacal Nitrogen	
26.8% Urea Nitrogen	
Available Phosphoric Acid(P <sub>2</sub> O <sub>5</sub> )	3%
Soluble Potash (K <sub>2</sub> O)	7%
Primary Plant Food Sources: Polymer-coated urea, urea, diammonium phosphate, mu potash	riate of
*7.0% slowly available urea nitrogen from polymer-coated urea.	
Chlorine (Cl), not more than	6.4%

# IV. Advertising claims that may be presented on the retail containers:

- (Fifteen, Sixteen, Eighteen or Twenty) Pounds Covers Up To 5,000 Square Feet.
- Controls/Kills dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf weeds; some of which are listed on the back panel
- ◆ An 18-6-12 fertilizer provides slow-release feeding to promote winter hardiness and drought tolerance in lawns.
- Feeds your lawn for up to 12 weeks.
- Controls/Kills over 30 varieties of weeds while feeding your lawn.
- Greens lawns and develops deep roots while controlling weeds.
- Bring out the best with the turf magic touch!
- Controls/Kills dandelion, plantain, and other weeds.
- Feeds the lawn with two (2) types of nitrogen and is pelleted for easy application.
- Controls (Number) weeds! (Listing of weeds may include the following.)

- · Triple Action Feeding works at every level.
  - Grass- Greener, Thicker lawn

Roots – Stronger Root System

Soil - Enriches & Fortifies Soil

- Builds a better lawn Guaranteed
- Gordon's Trimec Weed & Feed 30 kills unsightly weeds such as dandelion, chickweed and clover
  while it nourishes your lawn. Its Triple Action formula contains slow-release nitrogen which
  continuously feeds the grass, roots and soil for up to two full months [all winter long]. And it's
  guaranteed not to burn. The result is a greener, thicker lawn [this Fall and quicker green-up next
  Spring]. You're just one application away from a better lawn. That's the [company name] Better
  Lawn Guarantee.
- Triple Action Feeding works at every level for a Deep-Down Green!
- Why is Triple Action better? By feeding at all levels, your lawn receives the primary nutrients it
  needs to thrive. This ensures thicker, greener grass, stronger root system, and richer soil the
  key to building a lush lawn that lasts.
- Grass: Builds thicker, greener grass and enhances overall appearance.
- Roots: Builds stronger, deeper root systems which transport nutrients from the soil to the grass to help lawn withstand heat-stress and drought.
- Soil: Enriches and fortifies your soil. Rich, fertile soil provides a foundation for establishing roots and replenishes vital primary nutrients.
- For questions or comments, call toll-free 1-877-bayerAL (1-877-229-3725)
- Made in the U.S.A.
- · Kills weeds and builds a thick, green Fall lawn.
- Nourishes all Winter for a guick Spring green-up.

