

2217-579 3-19-2010

1/31



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs  
Registration Division (7505P)  
Ariel Rios Building  
1200 Pennsylvania Ave., NW  
Washington, D.C. 20460

NOTICE OF PESTICIDE:  
Registration  
 Reregistration  
(under FIFRA, as amended)

EPA Reg. Number:  
2217-579

Date of Issuance:  
MAR 19 2010

Term of Issuance:  
Conditional

Name of Pesticide Product:  
**Gordon's Trimec® Weed & Feed 30**

Name and Address of Registrant (include ZIP Code):

PBI/Gordon Corporation  
P.O. Box 014090  
Kansas City, Missouri 64101

~~Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA Registration Number.~~

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is reregistered in accordance with FIFRA sec. 4(g)(2)(C) provided that you:

- 1. Submit and or cite all data required for registration/re-registration review of your product when this Agency requires registrants of similar products to submit data.

Comments continued on pages 2 and 3.

Signature of Approving Official:

*Kathryn V. Montague*  
Kathryn V. Montague, Product Manager 23, HB, Registration Division

Date:

MAR 19 2010

- 2. Revise the chemical expression for dicamba in the Ingredient Statement to read: 3,6-dichloro-2-methoxybenzoic acid and add CAS Number 1982-69-0 as a footnote to the name of the a.i.; Sodium Salt of Dicamba. (Comply with PR Notice 97-5).

Also, Correct the chemical expression for MCPP to read:  
 "(+)-(R)-2-(4-chloro-2-methylphenoxy)propanoic acid"

- 3. On page 2, revise the precautions following the signal word to read: "Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wear long-sleeved shirt, long pants, shoes, socks and chemical resistant gloves made of any waterproof material. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet."
- 4. Revise the First Aid statements by reordering them into the following order: "IF ON SKIN OR CLOTHING", (Next, add: "IF INHALED", "Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, Preferable mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice."), "IF IN EYES" and last, "IF SWALLOWED".
- 5. On page 3, add the ground water warning to the alternate "Environmental Hazards" statement. It reads: "This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater."
- 6. The above comments apply to the sub-labels received November 11, 2008, submitted under the application on EPA Form 8570-1, dated November 5, 2008, to reregister this product.
- 7. On pages 4, 14 and 23, delete the "Suggested Spreader Settings" for the listed spreaders in the charts. On pages 5, 15 and 24, either delete the charts listing "Weeds" or connect these charts to the "Directions of Use". The same weeds are listed later in the labeling.
- 8. On pages 3,12, and 21, delete the word "General" or replace it with the word "Use" in the section head under "Directions for Use".

Page 3  
EPA Reg. No. 2217-579

8. On pages 3, 12 and 21, replace the word "since" with "because".
9. You must submit one copy of the final printed labeling before you release the product for shipment. Product shipped after 12 months from the date of this **Notice of Re-registration** must bear revised labeling reflecting the above comments.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product under the enclosed stamped copy of the label constitutes acceptance of these conditions. This labeling supersedes all previously accepted labeling. If you have any questions, you may call the Product Manager for this pesticide product registration.

---

Enclosure



5/31

**SUBLABEL 1: For use for Gordon's Trimec® Weed & Feed 30 24-4-8, and supplemental registrations.**

# **GORDON'S TRIMEC® WEED & FEED 30 24-4-8**

## **EPA Reg. No. 2217-579**

**ACTIVE INGREDIENT:**

2,4-dichlorophenoxyacetic acid .....	0.69%
(+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid .....	0.15%
Dicamba: 3,6-dichloro-o-anisic acid .....	0.06%
<b>INERT INGREDIENTS: .....</b>	<b>99.10%</b>
<b>TOTAL .....</b>	<b>100.00%</b>

**THIS PRODUCT CONTAINS:**

- 13.8 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton
- 3.0 lbs. (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid equivalent per ton
- 1.2 lbs. 3,6-dichloro-o-anisic acid equivalent per ton

Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated:**

MAR 19 2010

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

2217-579

**STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

## **PRECAUTIONARY STATEMENTS**

### **Hazards to Humans and Domestic Animals**

**CAUTION:** Causes moderate eye irritation. Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin, eyes or clothing. Wear long-sleeved shirt, long pants, socks and shoes. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

<b>First Aid</b>	
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call toll-free, 1-877-800-5556 for emergency medical treatment information.</p>	

**Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate. This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

**[Environmental Hazards (alternate per PRN 2008-1)**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.]

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**GENERAL PRECAUTIONS AND RESTRICTIONS:**

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Do not use on dichondra, carpetgrass, or St. Augustinegrass. Do not use this product where desirable clovers are present. Do not exceed specified dosages for any area. Apply when the air is calm. Application of this product to newly seeded areas should be delayed until the third mowing.
- Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

**INSTRUCTIONS:**

Gordon's Trimec® Weed & Feed 30 contains Trimec® herbicide. Trimec® controls a wide range of lawn weeds, including dandelion, chickweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a (X-X-X analysis of fertilizer grade) fertilizer provides a supply of plant food. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days. Treated areas may be seeded 3 to 4 weeks after application.

**[Note to Supplemental Registrants:** Guarantees for plant nutrients other than nitrogen, phosphorous, and potassium may be permitted or required by regulation by the \_\_\_\_\_. The guarantees for such other

nutrients shall be expressed in the form of the element. The source (oxides, salts, chelates, etc.) of such other nutrients may be required to be stated and may be included on the label.]

**[For state registrations in Kentucky, add the following paragraph:**

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing (\*Number of pounds) of nitrogen per 1,000 square feet. Please consult your local extension office for questions regarding a regular fertilizer program suitable for your geographic region.]

**Labeling Requirements:**

- Information regarding the contents and levels of metals in this product is available on the Internet as <http://www.aapfco.org/metals.htm>.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when the dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

**SPREADERS AND CALIBRATION:**

1. Drop spreaders and rotary spreaders must be calibrated for accurate application of this product.
2. Close spreader opening (orifice).
3. Adjust spreader setting using the Spreader Suggestions chart

Application Rate = 3 lbs. of product per 1,000 square feet.

<b>SUGGESTED SPREADER SETTINGS:</b>			
<b>Centrifugal/Rotary Spreaders</b>	<b>Application Rate 3 lbs./1,000 sq.ft.</b>	<b>Drop Spreaders</b>	<b>Application Rate 3 lbs./1,000 sq.ft.</b>
Frank's Broadcast	2.25	Frank's Drop	8.25
Scotts Speedy Green	5	Greenview II	5.5
Radio Flyer #0047	8.25	True Temper CD20	4 to 4.5
True Temper Rotary CB2000	3.75	Scotts Accugreen AG1	5.5
Earthway Rotary #2050SU	14.5	Earthway Drop #7300SU	13.5

These spreader settings are approximate. Differences may occur due to the condition of spreader, speed at which applied and roughness of area treated. Check your spreader on 330 square feet and adjust to apply at the rate of 1 pound per 1,000 square feet.


**4. CALIBRATION INSTRUCTIONS:**

- For 3.0 pounds of product per 1,000 square feet of turfgrass: Check you spreader on 330 square feet and adjust to apply at the rate of 1 pound of product per 330 square feet.
  - For 3.2 pounds of product per 1,000 square feet of turfgrass: Check you spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
  - For 3.6 pounds of product per 1,000 square feet of turfgrass: Check you spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
  - For 4.0 pounds of product per 1,000 square feet of turfgrass: Check you spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.
5. Fill spreader. Fill spreader on driveway or sidewalk to eliminate the chance of spillage on the lawn.
  6. Apply the product to lawn. Refer to the patch charts below:


The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application (1.2 lbs 2,4-D ae, 0.27 lb MCPP-p ae and 0.11 lb dicamba ae per acre per application). The maximum number

of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. (2.4 lbs 2,4-D ae, 0.54 lb MCPP-p ae and 0.22 lb dicamba ae per acre).

Path A for centrifugal/rotary spreaders.



Path B for drop spreaders.



7. Close the spreader openings when turning to change directions or when stopping.
8. If using a centrifugal/rotary spreader, avoid spreading product into adjacent landscape, flowers or vegetable plantings.

NOTE: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 square feet.

This \_\_\_ pound bag should cover approximately \_\_\_ square feet (\_\_\_ ft. x \_\_\_ ft.) Refer to the table and picture to determine the size of your lawn.

**COVERAGE CHART**


\_\_\_\_\_ Length

X \_\_\_\_\_ Width

= \_\_\_\_\_ Total Sq. Ft.

- \_\_\_\_\_ Non-Lawn Areas

= \_\_\_\_\_ Total Sq. Ft. Coverage you need.



24-4-8 GUARANTEED ANALYSIS	
Total Nitrogen (N) .....	24%
Water Insoluble Nitrogen .....	5%
Available Phosphate (P2O5) .....	4%
Soluble Potash (K2O) .....	8%

**[Note to Supplemental Registrants:** Fertilizer sources may vary in micronutrient composition, and iron may discolor or stain concrete. For fertilizers containing iron: Sweep all fertilizer from concrete surfaces immediately after application. Do not wash this product from concrete surfaces because fertilizers which contain iron may cause discoloration upon oxidation.]

WEEDS			
Bedstraw	Ground ivy	Peppergrass	Speedwell
Black medic	Healall	Pigweed	Spurge
Buckhorn	Henbit	Plantain	Thistle
Burdock	Knotweed	Poison ivy	Wild carrot
Chickweed	Lambsquarters	Poison oak	Wild garlic
Chicory	Lespedeza	Purslane	Wild lettuce
Clover	Mallow	Ragweed	Wild onion
Dandelion	Morningglory	Sheep sorrel	Yarrow
Dock	Oxalis	Shepherdspurse	



### STORAGE AND DISPOSAL

**PESTICIDE STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

**PESTICIDE DISPOSAL: If Empty** — Do not reuse this container. Place in trash or offer for recycling if available. **If Partly Filled** — Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

### LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

### APPENDIX

#### 1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Weight: \_\_\_\_\_
- EPA Est. No. \_\_\_\_\_

#### 2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

- **Previously Approved Fertilizer Grades:** Fertilizer grades with nutrient contents expressed as the percentages of N,P2O5, and K2O include the following:

10-6-4	16-3-5	20-10-0-(1)	23-7-7	25-4-8	28-3-7	30-4-8
10-16-20	16-6-9	20-10-5	24-3-7	26-3-3	28-3-8	32-4-8
10-18-10	18-3-5	21-5-7	24-4-8	26-3-5	29-3-4	22-3-14
12-5-5	18-6-12	22-6-6	25-3-3	26-4-6	30-3-4	30-3-4
15-5-10	20-4-6	22-7-7	25-3-5	26-4-13	30-3-5	
16-2-3	20-4-8	22-10-10	25-3-8	28-3-3	30-3-10	

The fertilizer grade guaranteed analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations. The following guaranteed analyses presentations have previously been submitted as notifications:

<b>10-16-20 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	10%
6.0% Ammoniacal Nitrogen	
4.0% Urea Nitrogen	
Available Phosphate (P2O5) .....	16%
Soluble Potash (K2O) .....	20%
Magnesium (Mg) .....	1.0%
0.1% Water Soluble Magnesium (Mg)	
Sulfur (S) .....	2.0%
2.0% Combined Sulfur (S)	
Iron (Fe) .....	1.0%
Manganese (Mn) .....	0.5%
0.1% Water Soluble Manganese (Mn)	
Nutrient Sources: Urea, Ammonium Phosphate, Ammonium Sulfate, Muriate of Potash, Ferrous Sulfate, Manganese Oxide, Manganese Sulfate, Sulfate of Potash Magnesia and Magnesium oxide.	

<b>29-3-4 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	29%
1.1% Ammoniacal Nitrogen	
27.9% Urea Nitrogen	
Available Phosphate (P2O5) .....	3%
Soluble Potash (K2O) .....	4%
Derived from: Polymer-coated urea, urea, diammonium phosphate, and muriate of potash. *11.0% slowly available urea nitrogen from polymer coated urea.	
Chlorine (Cl), note more than .....	4.0%

<b>10-18-10 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	10%
10.0% Ammoniacal Nitrogen	
Available Phosphate (P2O5) .....	18%
Soluble Potash (K2O) .....	10%
Sulfur (S) .....	2.0%
5.0% Combined Sulfur (S)	
Nutrient Sources: Ammonium Sulfate, Ammonium Phosphate and Muriate of Potash.	

<b>26-4-6 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	26%
1.8% Ammoniacal Nitrogen	
24.4% Urea Nitrogen	
Available Phosphoric Acid (P2O5) .....	4%
Soluble Potash (K2O) .....	6%
Boron (B) .....	0.02%
Copper (Cu) .....	0.05%
Iron (Fe) .....	1.00%
Total Manganese (Mn) .....	0.06%
Molybenum (Mo) .....	0.0005%
Zinc (Zn) .....	0.05%
Primary Plant Food Sources: Sulfur coated urea, urea, diammonium phosphate, muriate of potash. Secondary Plant Food Sources: Fertilizer grade borate, sulfates and oxides of copper, sulfates and oxides of iron, oxides of manganese, fertilizer grade molybdate, and sulfates and oxides of zinc. *4.0% slowly available urea nitrogen from sulfur coated urea.	
Chlorine (Cl), not more than .....	4.8%

<b>28-3-7 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	28%
1.2% Ammoniacal Nitrogen	
26.8% Urea Nitrogen	
Available Phosphoric Acid(P2O5) .....	3%
Soluble Potash (K2O) .....	7%
Primary Plant Food Sources: Polymer-coated urea, urea, diammonium phosphate, muriate of potash. *7.0% slowly available urea nitrogen from polymer-coated urea.	
Chlorine (Cl), not more than .....	6.4%

- (Fifteen, Sixteen, Eighteen or Twenty) Pounds Covers Up To 5,000 Square Feet.
- Controls/Kills dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf weeds; some of which are listed on the back panel
- An 18-6-12 fertilizer provides slow-release feeding to promote winter hardiness and drought tolerance in lawns.
- Feeds your lawn for up to 12 weeks.
- Controls/Kills over 30 varieties of weeds while feeding your lawn.
- Greens lawns and develops deep roots while controlling weeds.
- Bring out the best with the turf magic touch!
- Controls/Kills dandelion, plantain, and other weeds.
- Feeds the lawn with two (2) types of nitrogen and is pelleted for easy application.
- Controls (Number) weeds! (Listing of weeds may include the following.)
- Kills dandelion, chickweed, clover and most common lawn weeds.
- Triple Action Feeding works at every level.
  - Grass- Greener, Thicker lawn
  - Roots – Stronger Root System
  - Soil – Enriches & Fortifies Soil
- Builds a better lawn – Guaranteed
- Gordon's Trimec Weed & Feed 30 kills unsightly weeds such as dandelion, chickweed and clover while it nourishes your lawn. Its Triple Action formula contains slow-release nitrogen which continuously feeds the grass, roots and soil for up to two full months [all winter long]. And it's guaranteed not to burn. The result is a greener, thicker lawn [this Fall and quicker green-up next Spring]. You're just one application away from a better lawn. That's the [company name] Better Lawn Guarantee.

- Triple Action Feeding works at every level for a Deep-Down Green!
- Why is Triple Action better? By feeding at all levels, your lawn receives the primary nutrients it needs to thrive. This ensures thicker, greener grass, stronger root system, and richer soil – the key to building a lush lawn that lasts.
- Grass: Builds thicker, greener grass and enhances overall appearance.
- Roots: Builds stronger, deeper root systems which transport nutrients from the soil to the grass to help lawn withstand heat-stress and drought.
- Soil: Enriches and fortifies your soil. Rich, fertile soil provides a foundation for establishing roots and replenishes vital primary nutrients.
- For questions or comments, call toll-free 1-877-bayerAL (1-877-229-3725)
- Made in the U.S.A.
- Kills weeds and builds a thick, green Fall lawn.
- Nourishes all Winter for a quick Spring green-up.



- Controls most annual broadleaf weeds
- Premium slow release fertilizer stimulates grass growth to quickly conceal where weeds have died.
  - NITROGEN for Dense Green Lawns
  - PHOSPHORUS Improves Root Growth
  - POTASSIUM Vital for Growth and Drought Resistance

◆ Controls (Number) weeds! (Listing of weeds may include the following.)

WEEDS			
Annual yellow sweetclover	Cockle	Mustard	Spiny cocklebur
Aster	Cocklebur	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Common mullein	Narrowleaf vetch	Spotted catsear
Bedstraw	Creeping Charlie	Nettle	Spotted spurge
Beggarticks	Creeping jenny	Orange hawkweed	Spurweed
Betony, Florida	Cudweed	Oriental cocklebur	Stinging nettle
Birdweed, field	Curly dock	Oxalis	Strawberry, India mock
Bird vetch	Daisy fleabane	Parsley-piert	Tall nettle
Bitter wintercress	Daisy, oxeye	Parsnip	Tall vervain
Bittercress, hairy	Dandelion	Pearlwort	Tansy ragwort
Bitterweed	Dichondra	Pennycress	Tansy mustard
Black-eyed Susan	Dogfennel	Pepperweed	Tanweed
Black medic	Dollarweed	Pigweed	Thistle
Black mustard	False dandelion	Pineywoods bedstraw	Trailing crownvetch
Blackseed plantain	Falseflax	Plains coreopsis	Tumble mustard
Blessed thistle	False sunflower	Plantain	Tumble pigweed
Bloodflower milkweed	Fiddleneck	Poison ivy	Velvetleaf
Blue lettuce	Florida pusley	Poison oak	Venice mallow
Blue vervain	Frenchweed	Pokeweed	Virginia buttonweed
Bracted plantain	Galinsoga	Poorjoe	Virginia creeper
Bristly oxtongue	Goldenrod	Prairie sunflower	Virginia pepperweed
Broadleaf dock	Ground ivy	Prickly lettuce	Wavyleaf bullthistle
Broadleaf plantain	Gumweed	Prickly sida	Western salsify
Broomweed	Hairy fleabane	Prostrate knotweed	White mustard
Buckhorn	Hawkweed	Prostrate pigweed	Wild aster
Buckhorn plantain	Healall	Prostrate spurge	Wild buckwheat
Bulbous buttercup		Prostrate vervain	

WEEDS			
Bull thistle	Heartleaf drymary	Puncturevine	Wild carrot
Bullnettle	Heath aster	Purslane, common	Wild four-o'clock
Burclover	Hedge bindweed	Ragweed	Wild garlic
Burdock	Hedge mustard	Red sorrel	Wild geranium
Burning nettle	Henbit	Redroot pigweed	Wild lettuce
Burweed	Hoary cress	Redstem filaree	Wild marigold
Buttercup	Hoary plantain	Rough cinquefoil	Wild mustard
Buttonweed	Hoary vervain	Rough fleabane	Wild onion
Canada thistle	Horsenettle	Russian pigweed	Wild parsnip
Carolina geranium	Jimsonweed	Russian thistle	Wild radish
Carpetweed	Knawel	Scarlet pimpernel	Wild rape
Catchweed bedstraw	Knotweed	Scotch thistle	Wild strawberry
Catnip	Kochia	Sheep sorrel	Wild sweet potato
Catsear	Lambsquarters	Shepherdspurse	Wild vetch
Chickweed, common	Lespedeza	Slender plantain	Woodsorrel
Chickweed, mouseear	Mallow	Smallflower galinsoga	Woolly croton
Chicory	Matchweed	Smooth dock	Woolly morningglory
Cinquefoil	Mexicanweed	Smooth pigweed	Woolly plantain
Clover, crimson	Milk vetch	Sorrel	Wormseed
Clover, hop	Morningglory	Sowthistle	Yarrow
Clover, red	Mouseear hawkweed	Spanishneedles	Yellow rocket
Clover, strawberry	Mugwort	Speedwell	Yellowflower pepperweed
Clover, sweet	Musk thistle	Spiny amaranth	and other broadleaf weeds
Clover, white			

**SUBLABEL 2: For use by supplemental distributor for Bayer Advanced Triple Action Lawn Fertilizer Plus Weed Control II**

# **GORDON'S TRIMEC® WEED & FEED 30 24-4-8**

## **EPA Reg. No. 2217-579**

**ACTIVE INGREDIENT:**

2,4-dichlorophenoxyacetic acid .....	0.69%
(+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid .....	0.15%
Dicamba: 3,6-dichloro-o-anisic acid .....	0.06%
<b>INERT INGREDIENTS: .....</b>	<b>99.10%</b>
<b>TOTAL .....</b>	<b>100.00%</b>

**THIS PRODUCT CONTAINS:**

- 13.8 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton
- 3.0 lbs. (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid equivalent per ton
- 1.2 lbs. 3,6-dichloro-o-anisic acid equivalent per ton

Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

- 2,4-D (CAS No. 94-75-7)
- Mecoprop-p (CAS No. 16484-77-8)
- Dicamba (CAS No. 1918-00-9)

**KEEP OUT OF REACH OF CHILDREN**

### **CAUTION**

**STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

## **PRECAUTIONARY STATEMENTS**

### **Hazards to Humans and Domestic Animals**

**CAUTION:** Causes moderate eye irritation. Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin, eyes or clothing. Wear long-sleeved shirt, long pants, socks and shoes. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

<b>First Aid</b>	
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call toll-free, 1-877-800-5556 for emergency medical treatment information.</p>	

**Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate. This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

**[Environmental Hazards (alternate per PRN 2008-1)**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.]

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**GENERAL PRECAUTIONS AND RESTRICTIONS:**

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Do not use on dichondra, carpetgrass, closely mowed bentgrass putting greens or St. Augustinegrass. Do not use this product where desirable clovers are present. Do not exceed specified dosages for any area. Apply when the air is calm. Application of this product to newly seeded areas should be delayed until the third mowing.
- Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

**INSTRUCTIONS:**

Gordon's Trimec® Weed & Feed 30 contains Trimec® herbicide. Trimec® controls a wide range of lawn weeds, including dandelion, chickweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a (X-X-X analysis of fertilizer grade) fertilizer provides a supply of plant food. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds or, (b) if adverse low

moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days. Treated areas may be seeded 3 to 4 weeks after application.

**[Note to Supplemental Registrants:** Guarantees for plant nutrients other than nitrogen, phosphorous, and potassium may be permitted or required by regulation by the \_\_\_\_\_. The guarantees for such other nutrients shall be expressed in the form of the element. The source (oxides, salts, chelates, etc.) of such other nutrients may be required to be stated and may be included on the label.]

**[For state registrations in Kentucky, add the following paragraph:**

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing (\*Number of pounds) of nitrogen per 1,000 square feet. Please consult your local extension office for questions regarding a regular fertilizer program suitable for your geographic region.]

**Labeling Requirements:**

- Information regarding the contents and levels of metals in this product is available on the Internet as <http://www.aapfco.org/metals.htm>. [<http://www.regulatory-info-ba.com>]

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when the dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application (1.2 lbs 2,4-D ae, 0.27 lb MCP-p ae and 0.11 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. (2.4 lbs 2,4-D ae, 0.54 lb MCP-p ae and 0.22 lb dicamba ae per acre).

WEEDS			
Annual yellow sweetclover	Cocklebur	Narrowleaf plantain	Spiny cocklebur
Aster	Common mullein	Narrowleaf vetch	Spiny sowthistle
Austrian fieldcress	Creeping Charlie	Nettle	Spotted catsear
Bedstraw	Creeping jenny	Orange hawkweed	Spotted spurge
Beggarticks	Cudweed	Oriental cocklebur	Spurweed
Betony, Florida	Curly dock	Oxalis	Stinging nettle
Bindweed, field	Daisy, English	Parsley-piert	Strawberry, India mock
Bird vetch	Daisy fleabane	Parsnip	Tall nettle
Bitter wintercress	Daisy, oxeye	Pearlwort	Tall vervain
Bittercress, hairy	Dandelion	Pennycress	Tansy ragwort
Bitterweed	Dichondra	Pennywort	Tansy mustard
Black-eyed Susan	Dogbane	Peppergrass (Peppergrass)	Tanweed
Black medic	Dogfennel	Pepperweed	Thistle
Black mustard	Dollarweed	Pigweed	Trailing crownvetch
Blackseed plantain	Elderberry	Pineywoods bedstraw	Tumble mustard
Blessed thistle	False dandelion	Plains coreopsis (tickseed)	Tumble pigweed
Bloodflower milkweed	False flax	Plantain	Velvetleaf
Blue lettuce	False sunflower	Poison ivy	Venice mallow
Blue vervain	Fiddleneck	Poison oak	Virginia buttonweed
Bracted plantain	Florida pusley	Pokeweed	Virginia creeper
Bristly oxtongue	Frenchweed	Poorjoe	Virginia pepperweed
Broadleaf dock	Galinsoga	Prairie sunflower	Water pennywort
Broadleaf plantain	Goldenrod	Prickly lettuce	Wavyleaf bullthistle
Broomweed	Ground ivy	Prickly sida	Western clematis



<b>WEEDS</b>			
Buckhorn	Gumweed	Prostrate knotweed	Western salsify
Buckhorn plantain	Hairy fleabane	Prostrate pigweed	White mustard
Bulbous buttercup	Hawkweed	Prostrate spurge	Wild aster
Bull thistle	Healall	Prostrate vervain	Wild buckwheat
Bullnettle	Heartleaf drymary	Puncturevine	Wild carrot
Burclover	Heathaster	Purslane, common	Wild four-o'clock
Burdock	Hedge bindweed	Ragweed	Wild garlic
Burning nettle	Hedge mustard	Red sorrel	Wild geranium
Burweed	Henbit	Redroot pigweed	Wild lettuce
Buttercup	Hoary cress	Redstem filaree	Wild marigold
Buttonweed	Hoary plantain	Rough cinquefoil	Wild mustard
Canada thistle	Hoary vervain	Rough fleabane	Wild onion
Carolina geranium	Horsenettle	Russian pigweed	Wild parsnip
Carpetweed	Jimsonweed	Russian thistle	Wild radish
Catchweed bedstraw	Knawel	Scarlet pimpernel	Wild rape
Catnip	Knotweed	Scotch thistle	Wild strawberry
Catsear	Kochia	Sheep sorrel	Wild sweet potato
Chickweed, common	Lambsquarters	Shepherdspurse	Wild vetch
Chickweed, mouseear	Lespedeza	Slender plantain	Woodsorrel
Chicory	Mallow	Smallflower galinsoga	Woolly croton
Cinquefoil	Matchweed	Smooth dock	Woolly morningglory
Clover, crimson	Mexicanweed	Smooth pigweed	Woolly plantain
Clover, hop	Milk vetch	Sorrel	Wormseed
Clover, red	Morningglory	Sowthistle	Yarrow
Clover, strawberry	Mouseear hawkweed	Spanishneedles	Yellow rocket
Clover, sweet	Mugwort	Speedwell	Yellowflower pepperweed
Clover, white	Musk thistle	Spiny amaranth	and other broadleaf weeds
Cockle	Mustard		

**SPREADERS AND CALIBRATION:**

1. Drop spreaders and rotary spreaders must be calibrated for uniform application of this product.
2. Close spreader opening (orifice).
3. Adjust spreader setting using the Spreader Suggestions chart

Application Rate = 3 lbs. of product per 1,000 square feet.

<b>SUGGESTED SPREADER SETTINGS:</b>			
<b>Centrifugal/Rotary Spreaders</b>	<b>Application Rate 3 lbs./1,000 sq.ft.</b>	<b>Drop Spreaders</b>	<b>Application Rate 3 lbs./1,000 sq.ft.</b>
Frank's Broadcast	2.25	Frank's Drop	8.25
Scotts Speedy Green	5	Greenview II	5.5
Radio Flyer #0047	8.25	True Temper CD20	4 to 4.5
True Temper Rotary CB2000	3.75	Scotts Accugreen AG1	5.5
Earthway Rotary #2050SU	14.5	Earthway Drop #7300SU	13.5

These spreader settings are approximate. Differences may occur due to the condition of spreader, speed at which applied and roughness of area treated. Check your spreader on 330 square feet and adjust to apply at the rate of 1 pound per 1,000 square feet.



**4. CALIBRATION INSTRUCTIONS:**

- For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 330 square feet and adjust to apply at the rate of 1 pound of product per 330 square feet.
- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

5. Fill spreader. Fill spreader on driveway or sidewalk to eliminate the chance of spillage on the lawn.
6. Apply the product to lawn. Refer to the patch charts below:

Path A for centrifugal/rotary spreaders.

Path B for drop spreaders.




7. Close the spreader opening when turning to change directions or when stopping.
8. If using a centrifugal/rotary spreader, avoid spreading product into adjacent landscape, flowers or vegetable plantings.

NOTE: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 square feet.

This \_\_\_ pound bag should cover approximately \_\_\_ square feet (\_\_\_ ft. x \_\_\_ ft.) Refer to the table and picture to determine the size of your lawn.

**COVERAGE CHART**

\_\_\_\_\_ Length  
 x \_\_\_\_\_ Width  
 = \_\_\_\_\_ Total Sq. Ft.  
 - \_\_\_\_\_ Non-Lawn Areas  
 = \_\_\_\_\_ Total Sq. Ft. Coverage you need.



24-4-8 GUARANTEED ANALYSIS	
Total Nitrogen (N) .....	24%
Water Insoluble Nitrogen .....	5%
Available Phosphate (P2O5) .....	4%
Soluble Potash (K2O) .....	8%

[Note to Supplemental Registrants: Fertilizer sources may vary in micronutrient composition, and iron may discolor or stain concrete. For fertilizers containing iron: Sweep all fertilizer from concrete surfaces immediately after application. Do not wash this product from concrete surfaces because fertilizers which contain iron may cause discoloration upon oxidation.]

WEEDS			
Bedstraw	Ground ivy	Peppergrass	Speedwell
Black medic	Healall	Pigweed	Spurge
Buckhorn	Henbit	Plantain	Thistle
Burdock	Knotweed	Poison ivy	Wild carrot
Chickweed	Lambsquarters	Poison oak	Wild garlic
Chicory	Lespedeza	Purslane	Wild lettuce
Clover	Mallow	Ragweed	Wild onion
Dandelion	Morningglory	Sheep sorrel	Yarrow
Dock	Oxalis	Shepherdspurse	

**STORAGE AND DISPOSAL**  
**PESTICIDE STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.  
**PESTICIDE DISPOSAL: If Empty** — Do not reuse this container. Place in trash or offer for recycling if available. **If Partly Filled** — Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

**LIMITED WARRANTY AND DISCLAIMER**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once to the seller.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

**APPENDIX**

**1. Statements which may appear on different label components depending on packaging configuration.**

- See next panel for additional Precautionary Statements and First Aid
- Net Weight: \_\_\_\_\_
- EPA Est. No. \_\_\_\_\_

**2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:**

- **Previously Approved Fertilizer Grades:** Fertilizer grades with nutrient contents expressed as the percentages of N,P2O5, and K2O include the following:

10-6-4	16-3-5	20-10-0-(1)	23-7-7	25-4-8	28-3-7	30-4-8
10-16-20	16-6-9	20-10-5	24-3-7	26-3-3	28-3-8	32-4-8
10-18-10	18-3-5	21-5-7	24-4-8	26-3-5	29-3-4	
12-5-5	18-6-12	22-6-6	25-3-3	26-4-6	30-3-4	
15-5-10	20-4-6	22-7-7	25-3-5	26-4-13	30-3-5	
16-2-3	20-4-8	22-10-10	25-3-8	28-3-3	30-3-10	

The fertilizer grade guaranteed analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations. The following guaranteed analyses presentations have previously been submitted as notifications:

<b>10-16-20 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	10%
6.0% Ammoniacal Nitrogen	
4.0% Urea Nitrogen	
Available Phosphate (P2O5) .....	16%
Soluble Potash (K2O) .....	20%
Magnesium (Mg) .....	1.0%
0.1% Water Soluble Magnesium (Mg)	
Sulfur (S) .....	2.0%
2.0% Combined Sulfur (S)	
Iron (Fe) .....	1.0%
Manganese (Mn) .....	0.5%
0.1% Water Soluble Manganese (Mn)	
Nutrient Sources: Urea, Ammonium Phosphate, Ammonium Sulfate, Muriate of Potash, Ferrous Sulfate, Manganese Oxide, Manganese Sulfate, Sulfate of Potash Magnesia and Magnesium oxide.	

<b>29-3-4 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	29%
1.1% Ammoniacal Nitrogen	
27.9% Urea Nitrogen	
Available Phosphate (P2O5) .....	3%
Soluble Potash (K2O) .....	4%
Derived from: Polymer-coated urea, urea, diammonium phosphate, and muriate of potash. *11.0% slowly available urea nitrogen from polymer coated urea.	
Chlorine (Cl), not more than .....	4.0%

<b>10-18-10 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	10%
10.0% Ammoniacal Nitrogen	
Available Phosphate (P3O5) .....	18%
Soluble Potash (K2O) .....	10%
Sulfur (S) .....	2.0%
5.0% Combined Sulfur (S)	
Nutrient Sources: Ammonium Sulfate, Ammonium Phosphate and Muriate of Potash.	

<b>26-4-6 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	26%
1.8% Ammoniacal Nitrogen	
24.4% Urea Nitrogen	
Available Phosphoric Acid (P2O5) .....	4%
Soluble Potash (K2O) .....	6%
Boron (B) ..... 0.02%	
Copper (Cu) ..... 0.05%	
Iron (Fe) ..... 1.00%	
Total Manganese (Mn) ..... 0.06%	
Molybdenum (Mo) ..... 0.0005%	
Zinc (Zn) ..... 0.05%	
Primary Plant Food Sources: Sulfur coated urea, urea, diammonium phosphate, muriate of potash. Secondary Plant Food Sources: Fertilizer grade borate, sulfates and oxides of copper, sulfates and oxides of iron, oxides of manganese, fertilizer grade molybdate, and sulfates and oxides of zinc. *4.0% slowly available urea nitrogen from sulfur coated urea.	
Chlorine (Cl), not more than .....	4.8%

<b>28-3-7 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	28%
1.2% Ammoniacal Nitrogen	
26.8% Urea Nitrogen	
Available Phosphoric Acid (P2O5) .....	3%
Soluble Potash (K2O) .....	7%
Primary Plant Food Sources: Polymer-coated urea, urea, diammonium phosphate, muriate of potash. *7.0% slowly available urea nitrogen from polymer-coated urea.	
Chlorine (Cl), not more than .....	6.4%

- (Fifteen, Sixteen, Eighteen or Twenty) Pounds Covers Up To 5,000 Square Feet.
- Controls/Kills dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf weeds; some of which are listed on the back panel
- An 18-6-12 fertilizer provides slow-release feeding to promote winter hardiness and drought tolerance in lawns.
- Feeds your lawn for up to 12 weeks.
- Controls/Kills over 30 varieties of weeds while feeding your lawn.
- Greens lawns and develops deep roots while controlling weeds.
- Bring out the best with the turf magic touch!
- Controls/Kills dandelion, plantain, and other weeds.
- Feeds the lawn with two (2) types of nitrogen and is pelleted for easy application.
- Controls (Number) weeds! (Listing of weeds may include the following.)
- Kills dandelion, chickweed, clover and most common lawn weeds.
- Triple Action Feeding works at every level.
  - Grass- Greener, Thicker lawn
  - Roots – Stronger Root System
  - Soil – Enriches & Fortifies Soil
- Builds a better lawn – Guaranteed
- Gordon's Trimec Weed & Feed 30 kills unsightly weeds such as dandelion, chickweed and clover while it nourishes your lawn. Its Triple Action formula contains slow-release nitrogen which continuously feeds the grass, roots and soil for up to two full months [all winter long]. And it's guaranteed not to burn. The result is a greener, thicker lawn [this Fall and quicker green-up next Spring]. You're just one application away from a better lawn. That's the [company name] Better Lawn Guarantee.

- Triple Action Feeding works at every level for a Deep-Down Green!
- Why is Triple Action better? By feeding at all levels, your lawn receives the primary nutrients it needs to thrive. This ensures thicker, greener grass, stronger root system, and richer soil – the key to building a lush lawn that lasts.
- Grass: Builds thicker, greener grass and enhances overall appearance.
- Roots: Builds stronger, deeper root systems which transport nutrients from the soil to the grass to help lawn withstand heat-stress and drought.
- Soil: Enriches and fortifies your soil. Rich, fertile soil provides a foundation for establishing roots and replenishes vital primary nutrients.
- For questions or comments, call toll-free 1-877-bayerAL (1-877-229-3725)
- Made in the U.S.A.
- Kills weeds and builds a thick, green Fall lawn.
- Nourishes all Winter for a quick Spring green-up.



23/  
31

**SUBLABEL 3: For use by supplemental distributor for  
Bayer Advanced Triple Action Winterizer Fertilizer Plus Weed Control II**

# **GORDON'S TRIMEC® WEED & FEED 30 24-4-8**

## **EPA Reg. No. 2217-579**

**ACTIVE INGREDIENT:**

2,4-dichlorophenoxyacetic acid .....	0.69%
(+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid .....	0.15%
Dicamba: 3,6-dichloro-o-anisic acid .....	0.06%
<b>INERT INGREDIENTS: .....</b>	<b>99.10%</b>
<b>TOTAL .....</b>	<b>100.00%</b>

**THIS PRODUCT CONTAINS:**

- 13.8 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton
- 3.0 lbs. (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid equivalent per ton
- 1.2 lbs. 3,6-dichloro-o-anisic acid equivalent per ton

Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

- 2,4-D (CAS No. 94-75-7)
- Mecoprop-p (CAS No. 16484-77-8)
- Dicamba (CAS No. 1918-00-9)

### **KEEP OUT OF REACH OF CHILDREN**

### **CAUTION**

**STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

## **PRECAUTIONARY STATEMENTS**

### **Hazards to Humans and Domestic Animals**

**CAUTION:** Causes moderate eye irritation. Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin, eyes or clothing. Wear long-sleeved shirt, long pants, socks and shoes. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

<b>First Aid</b>	
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call toll-free, 1-877-800-5556 for emergency medical treatment information.</p>	

**Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate. This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

**[Environmental Hazards (alternate per PRN 2008-1)**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.]

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**GENERAL PRECAUTIONS AND RESTRICTIONS:**

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Do not use on dichondra, carpetgrass, bentgrass or St. Augustinegrass. Do not use this product where desirable clovers are present. Do not exceed specified dosages for any area. Apply when the air is calm. Application of this product to newly seeded areas should be delayed until the third mowing.
- Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

**INSTRUCTIONS:**

Gordon's Trimec® Weed & Feed 30 contains Trimec® herbicide. Trimec® controls a wide range of lawn weeds, including dandelion, chickweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a (X-X-X analysis of fertilizer grade) fertilizer provides a supply of plant food. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds or, (b) if adverse low



moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in 30 days. Treated areas may be seeded 3 to 4 weeks after application.

**[Note to Supplemental Registrants:** Guarantees for plant nutrients other than nitrogen, phosphorous, and potassium may be permitted or required by regulation by the \_\_\_\_\_. The guarantees for such other nutrients shall be expressed in the form of the element. The source (oxides, salts, chelates, etc.) of such other nutrients may be required to be stated and may be included on the label.]

**[For state registrations in Kentucky, add the following paragraph:**

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing (\*Number of pounds) of nitrogen per 1,000 square feet. Please consult your local extension office for questions regarding a regular fertilizer program suitable for your geographic region.]

**Labeling Requirements:**

- Information regarding the contents and levels of metals in this product is available on the Internet as <http://www.aapfco.org/metals.htm>. [<http://www.regulatory-info-ba.com>]

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when the dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application (1.2 lbs 2,4-D ae, 0.27 lb MCP-p ae and 0.11 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. (2.4 lbs 2,4-D ae, 0.54 lb MCP-p ae and 0.22 lb dicamba ae per acre).

WEEDS			
Annual yellow sweetclover	Cocklebur	Narrowleaf plantain	Spiny cocklebur
Aster	Common mullein	Narrowleaf vetch	Spiny sowthistle
Austrian fieldcress	Creeping Charlie	Nettle	Spotted catsear
Bedstraw	Creeping jenny	Orange hawkweed	Spotted spurge
Beggarticks	Cudweed	Ornamental cocklebur	Sturweed
Betony, Florida	Curly dock	Oxalis	Stinging nettle
Bindweed, field	Daisy, English	Parsley-piert	Strawberry, India mock
Bird vetch	Daisy fleabane	Parsnip	Tall nettle
Bitter wintercress	Daisy, oxeye	Pearlwort	Tall vervain
Bittercress, hairy	Dandelion	Pennycress	Tansy ragwort
Bitterweed	Dichondra	Pennywort	Tansy mustard
Black-eyed Susan	Dogbane	Peppergrass (Peppergrass)	Tanweed
Black medic	Dogfennel	Pepperweed	Thistle
Black mustard	Dollarweed	Pigweed	Trailing crownvetch
Blackseed plantain	Elderberry	Pineywoods bedstraw	Tumble mustard
Blessed thistle	False dandelion	Plains coreopsis (tickseed)	Tumble pigweed
Bloodflower milkweed	False flax	Plantain	Velvetleaf
Blue lettuce	False sunflower	Poison ivy	Venice mallow
Blue vervain	Fiddleneck	Poison oak	Virginia buttonweed
Bracted plantain	Florida pusley	Pokeweed	Virginia creeper
Bristly oxtongue	Frenchweed	Poorjoe	Virginia pepperweed
Broadleaf dock	Galinsoga	Prairie sunflower	Water pennywort
Broadleaf plantain	Goldenrod	Prickly lettuce	Wavyleaf bullthistle
Broomweed	Ground ivy	Prickly sida	Western clematis

<b>WEEDS</b>			
Buckhorn	Gumweed	Prostrate knotweed	Western salsify
Buckhorn plantain	Hairy fleabane	Prostrate pigweed	White mustard
Bulbous buttercup	Hawkweed	Prostrate spurge	Wild aster
Bull thistle	Healall	Prostrate vervain	Wild buckwheat
Bullnettle	Heartleaf drymary	Puncturevine	Wild carrot
Burclover	Heathaster	Purslane, common	Wild four-o'clock
Burdock	Hedge bindweed	Ragweed	Wild garlic
Burning nettle	Hedge mustard	Red sorrel	Wild geranium
Burweed	Henbit	Redroot pigweed	Wild lettuce
Buttercup	Hoary cress	Redstem filaree	Wild marigold
Buttonweed	Hoary plantain	Rough cinquefoil	Wild mustard
Canada thistle	Hoary vervain	Rough fleabane	Wild onion
Carolina geranium	Horsenettle	Russian pigweed	Wild parsnip
Carpetweed	Jimsonweed	Russian thistle	Wild radish
Catchweed bedstraw	Knawel	Scarlet pimpernel	Wild rape
Catnip	Knotweed	Scotch thistle	Wild strawberry
Catsear	Kochia	Sheep sorrel	Wild sweet potato
Chickweed, common	Lambsquarters	Shepherdspurse	Wild vetch
Chickweed, mouseear	Lespedeza	Slender plantain	Woodsorrel
Chicory	Mallow	Smallflower galinsoga	Woolly croton
Cinquefoil	Matchweed	Smooth dock	Woolly morningglory
Clover, crimson	Mexicanweed	Smooth pigweed	Woolly plantain
Clover, hop	Milk vetch	Sorrel	Wormseed
Clover, red	Morningglory	Sowthistle	Yarrow
Clover, strawberry	Mouseear hawkweed	Spanishneedles	Yellow rocket
Clover, sweet	Mugwort	Speedwell	Yellowflower pepperweed and other broadleaf weeds
Clover, white	Musk thistle	Spiny amaranth	
Cockle	Mustard		

**SPREADERS AND CALIBRATION:**

1. Drop spreaders and rotary spreaders must be calibrated for uniform application of this product.
2. Close spreader opening (orifice).
3. Adjust spreader setting using the Spreader Suggestions chart

Application Rate = 3 lbs. of product per 1,000 square feet.

<b>SUGGESTED SPREADER SETTINGS:</b>			
<b>Centrifugal/Rotary Spreaders</b>	<b>Application Rate 3 lbs./1,000 sq.ft.</b>	<b>Drop Spreaders</b>	<b>Application Rate 3 lbs./1,000 sq.ft.</b>
Frank's Broadcast	2.25	Frank's Drop	8.25
Scotts Speedy Green	5	Greenview II	5.5
Radio Flyer #0047	8.25	True Temper CD20	4 to 4.5
True Temper Rotary CB2000	3.75	Scotts Accugreen AG1	5.5
Earthway Rotary #2050SU	14.5	Earthway Drop #7300SU	13.5

These spreader settings are approximate. Differences may occur due to the condition of spreader, speed at which applied and roughness of area treated. Check your spreader on 330 square feet and adjust to apply at the rate of 1 pound per 1,000 square feet.



**4. CALIBRATION INSTRUCTIONS:**

- For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 330 square feet and adjust to apply at the rate of 1 pound of product per 330 square feet.
- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

5. Fill spreader. Fill spreader on driveway or sidewalk to eliminate the chance of spillage on the lawn.
6. Apply the product to lawn. Refer to the patch charts below:

Path A for centrifugal/rotary spreaders.

Path B for drop spreaders.



7. Close the spreader openings when turning to change directions or when stopping.
8. If using a centrifugal/rotary spreader, avoid spreading product into adjacent landscape, flowers or vegetable plantings.

NOTE: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 square feet.

This \_\_\_ pound bag should cover approximately \_\_\_ square feet (\_\_\_ ft. x \_\_\_ ft.) Refer to the table and picture to determine the size of your lawn.

**COVERAGE CHART**


\_\_\_\_\_ Length

x \_\_\_\_\_ Width

= \_\_\_\_\_ Total Sq. Ft.

- \_\_\_\_\_ Non-Lawn Areas

= \_\_\_\_\_ Total Sq. Ft. Coverage you need.



24-4-8 GUARANTEED ANALYSIS	
Total Nitrogen (N) .....	24%
Water Insoluble Nitrogen .....	5%
Available Phosphate (P2O5) .....	4%
Soluble Potash (K2O) .....	8%

**[Note to Supplemental Registrants:** Fertilizer sources may vary in micronutrient composition, and iron may discolor or stain concrete. For fertilizers containing iron: Sweep all fertilizer from concrete surfaces immediately after application. Do not wash this product from concrete surfaces because fertilizers which contain iron may cause discoloration upon oxidation.]

WEEDS			
Bedstraw	Ground ivy	Peppergrass	Speedwell
Black medic	Healall	Pigweed	Spurge
Buckhorn	Henbit	Plantain	Thistle
Burdock	Knotweed	Poison ivy	Wild carrot
Chickweed	Lambsquarters	Poison oak	Wild garlic
Chicory	Lespedeza	Purslane	Wild lettuce
Clover	Mallow	Ragweed	Wild onion
Dandelion	Morningglory	Sheep sorrel	Yarrow
Dock	Oxalis	Shepherdspurse	

### STORAGE AND DISPOSAL

**PESTICIDE STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.

**PESTICIDE DISPOSAL: If Empty** — Do not reuse this container. Place in trash or offer for recycling if available. **If Partly Filled** — Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

### LIMITED WARRANTY AND DISCLAIMER

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once to the seller.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

### APPENDIX

#### 1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Weight: \_\_\_\_\_
- EPA Est. No. \_\_\_\_\_

#### 2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

- Previously Approved Fertilizer Grades: Fertilizer grades with nutrient contents expressed as the percentages of N,P2O5, and K5O include the following:

10-6-4	16-3-5	20-10-0-(1)	23-7-7	25-4-8	28-3-7	30-4-8
10-16-20	16-6-9	20-10-5	24-3-7	26-3-3	28-3-8	32-4-8
10-18-10	18-3-5	21-5-7	24-4-8	26-3-5	29-3-4	
12-5-5	18-6-12	22-6-6	25-3-3	26-4-6	30-3-4	
15-5-10	20-4-6	22-7-7	25-3-5	26-4-13	30-3-5	
16-2-3	20-4-8	22-10-10	25-3-8	28-3-3	30-3-10	

The fertilizer grade guaranteed analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations. The following guaranteed analyses presentations have previously been submitted as notifications:

<b>10-16-20 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	10%
6.0% Ammoniacal Nitrogen	
4.0% Urea Nitrogen	
Available Phosphate (P2O5) .....	16%
Soluble Potash (K2O) .....	20%
Magnesium (Mg) .....	1.0%
0.1% Water Soluble Magnesium (Mg)	
Sulfur (S) .....	2.0%
2.0% Combined Sulfur (S)	
Iron (Fe) .....	1.0%
Manganese (Mn) .....	0.5%
0.1% Water Soluble Manganese (Mn)	
Nutrient Sources: Urea, Ammonium Phosphate, Ammonium Sulfate, Muriate of Potash, Ferrous Sulfate, Manganese Oxide, Manganese Sulfate, Sulfate of Potash Magnesia and Magnesium oxide.	

<b>29-3-4 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	29%
1.1% Ammoniacal Nitrogen	
27.9% Urea Nitrogen	
Available Phosphate (P2O5) .....	3%
Soluble Potash (K2O) .....	4%
Derived from: Polymer-coated urea, urea, diammonium phosphate, and muriate of potash. *11.0% slowly available urea nitrogen from polymer coated urea.	
Chlorine (Cl), not more than .....	4.0%

<b>10-18-10 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	10%
10.0% Ammoniacal Nitrogen	
Available Phosphate (P2O5) .....	18%
Soluble Potash (K2O) .....	10%
Sulfur (S) .....	2.0%
5.0% Combined Sulfur (S)	
Nutrient Sources: Ammonium Sulfate, Ammonium Phosphate and Muriate of Potash.	

<b>26-4-6 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	26%
1.8% Ammoniacal Nitrogen	
24.4% Urea Nitrogen	
Available Phosphoric Acid (P2O5) .....	4%
Soluble Potash (K2O) .....	6%
Boron (B) ..... 0.02%	
Copper (Cu) ..... 0.05%	
Iron (Fe) ..... 1.00%	
Total Manganese (Mn) ..... 0.06%	
Molybdenum (Mo) ..... 0.0005%	
Zinc (Zn) ..... 0.05%	
Primary Plant Food Sources: Sulfur coated urea, urea, diammonium phosphate, muriate of potash. Secondary Plant Food Sources: Fertilizer grade borate, sulfates and oxides of copper, sulfates and oxides of iron, oxides of manganese, fertilizer grade molybdate, and sulfates and oxides of zinc. *4.0% slowly available urea nitrogen from sulfur coated urea.	
Chlorine (Cl), not more than .....	4.8%

<b>28-3-7 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	28%
1.2% Ammoniacal Nitrogen	
26.8% Urea Nitrogen	
Available Phosphoric Acid(P2O5) .....	3%
Soluble Potash (K2O) .....	7%
Primary Plant Food Sources: Polymer-coated urea, urea, diammonium phosphate, muriate of potash *7.0% slowly available urea nitrogen from polymer-coated urea.	
Chlorine (Cl), not more than .....	6.4%

- (Fifteen, Sixteen, Eighteen or Twenty) Pounds Covers Up To 5,000 Square Feet.
- Controls/Kills dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf weeds; some of which are listed on the back panel
- An 18-6-12 fertilizer provides slow-release feeding to promote winter hardiness and drought tolerance in lawns.
- Feeds your lawn for up to 12 weeks.
- Controls/Kills over 30 varieties of weeds while feeding your lawn.
- Greens lawns and develops deep roots while controlling weeds.
- Bring out the best with the turf magic touch!
- Controls/Kills dandelion, plantain, and other weeds.
- Feeds the lawn with two (2) types of nitrogen and is pelleted for easy application.
- Controls (Number) weeds! (Listing of weeds may include the following.)
- Kills dandelion, chickweed, clover and most common lawn weeds.
- Triple Action Feeding works at every level.
  - Grass- Greener, Thicker lawn
  - Roots – Stronger Root System
  - Soil – Enriches & Fortifies Soil
- Builds a better lawn – Guaranteed
- Gordon's Trimec Weed & Feed 30 kills unsightly weeds such as dandelion, chickweed and clover while it nourishes your lawn. Its Triple Action formula contains slow-release nitrogen which continuously feeds the grass, roots and soil for up to two full months [all winter long]. And it's guaranteed not to burn. The result is a greener, thicker lawn [this Fall and quicker green-up next Spring]. You're just one application away from a better lawn. That's the [company name] Better Lawn Guarantee.

- Triple Action Feeding works at every level for a Deep-Down Green!
- Why is Triple Action better? By feeding at all levels, your lawn receives the primary nutrients it needs to thrive. This ensures thicker, greener grass, stronger root system, and richer soil – the key to building a lush lawn that lasts.
- Grass: Builds thicker, greener grass and enhances overall appearance.
- Roots: Builds stronger, deeper root systems which transport nutrients from the soil to the grass to help lawn withstand heat-stress and drought.
- Soil: Enriches and fortifies your soil. Rich, fertile soil provides a foundation for establishing roots and replenishes vital primary nutrients.
- For questions or comments, call toll-free 1-877-bayerAL (1-877-229-3725)
- Made in the U.S.A.
- Kills weeds and builds a thick, green Fall lawn.
- Nourishes all Winter for a quick Spring green-up.



## **DOCUMENT CONTROL INFORMATION**

1. **Unique Label Identifier:** 002217-00579.20081105.proposed.doc
2. **Reason for Issue:** Various changes pertaining to re-registration. Re-format label.