



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

January 23, 2017

Mr. Eric Smith  
Federal Registration Manager  
PBI/Gordon Corporation  
P.O. Box 014090  
Kansas City, MO 64101

Subject: Notification per PRN 98-10 – Added weeds, alternative warranty statement, marketing claims  
Product Name: Gordon's Trimec Weed & Feed 30 24-4-8  
EPA Registration Number: 2217-579  
Application Date: November 2, 2016  
Decision Number: 523315

Dear Mr. Smith:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If you have any questions, you may contact Kathryn Montague at 703-305-1243 or via email at [montague.kathryn@epa.gov](mailto:montague.kathryn@epa.gov).

Sincerely,

A handwritten signature in black ink that reads "Kathryn V. Montague".

Kathryn Montague, Product Manager 23  
Herbicide Branch  
Registration Division (7505P)  
Office of Pesticide Programs

# GORDON'S TRIMEC® WEED & FEED 30 24-4-8

NOTIFICATION

2217-579

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

01/23/2017

## EPA Reg. No. 2217-579

Three sublabels represent the entire master label of GORDON'S TRIMEC® WEED & FEED 30 24-4-8. Please refer to the appropriate section of the labeling as shown as follows:

<b>Section</b>
<b>Sublabel 1: Gordon's Trimec® Weed &amp; Feed 30 24-4-8</b>
<b>Sublabel 2: For use by supplemental distributor for Bayer Advanced(TM) Triple Action Lawn Fertilizer Plus Weed Control II</b>
<b>Sublabel 3: For use by supplemental distributor for Bayer Advanced(TM) Triple Action Winterizer Fertilizer Plus Weed Control II</b>
<b>Appendix</b>
<b>Document Control Information</b>

# GORDON'S TRIMEC® WEED & FEED 30 24-4-8

## EPA Reg. No. 2217-579

### ACTIVE INGREDIENTS:

2,4-dichlorophenoxyacetic acid (CAS Registry No. 94-75-7) .....	0.69%
(+)-(R)-2-(4-chloro-2 methylphenoxy) propionic acid (CAS Registry No. 16484-77-8) .....	0.15%
Dicamba: 3,6-dichloro-2-methoxybenzoic acid (CAS Registry No. 1918-00-9) .....	0.06%
<b>INERT OTHER</b> INGREDIENTS: .....	<u>99.10%</u>
TOTAL .....	100.00%

### THIS PRODUCT CONTAINS:

- 13.8 lb 2,4-dichlorophenoxyacetic acid equivalent per ton (CAS No. 94-75-7)
- 3.0 lb (+)-(R)-2-(4-chloro-2 methylphenoxy) propionic acid equivalent per ton (CAS No. 16484-77-8)
- 1.2 lb 3,6-dichloro-2-methoxybenzoic acid equivalent per ton (CAS No. 1918-00-9)

Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

**STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

See attached booklet for complete [Precautionary Statements and] Directions for Use, including Storage and Disposal, and back panel for [complete Precautionary Statements, including] First Aid. OR See back panel for complete Precautionary Statements, including First Aid, and Directions for Use, including Storage and Disposal.

**Guaranteed Analysis:** See appendix for example.

**Net Contents:** \_\_\_\_\_  
**EPA Est. No.** \_\_\_\_\_

### Company Name and Address:

PBI/Gordon Corporation  
1217 West 12<sup>th</sup> Street  
Kansas City, Missouri 64101



# **PRECAUTIONARY STATEMENTS**

## **Hazards to Humans and Domestic Animals**

**CAUTION:** Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wear long-sleeved shirt, long pants, shoes, socks, and chemical resistant gloves made of any waterproof material. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

<b>First Aid</b>	
<b>If on skin or on clothing:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to mouth if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call toll-free, 1-877-800-5556 for emergency medical treatment information.	

## **Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

## **[Environmental Hazards (alternate per PRN 2008-1)**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.]

---

**SUBLABEL 1: For use for Gordon's Trimec® Weed & Feed 30 24-4-8, and supplemental registrations.**

---

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### **PRECAUTIONS AND USE RESTRICTIONS:**

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Do not use on dichondra, carpetgrass, or St. Augustinegrass. Do not use this product where desirable clovers are present. Do not exceed specified dosages for any area. Apply when the air is calm. Application of this product to newly seeded areas should be delayed until the third mowing.
- Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, because injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

### **INSTRUCTIONS:**

Gordon's Trimec® Weed & Feed 30 contains Trimec® herbicide. Trimec® controls a wide range of lawn weeds, including dandelion, chickweed, plantain, henbit, spurge and many other listed weeds commonly found in home lawns. In addition, a (X-X-X [analysis of fertilizer grade]) fertilizer provides a supply of plant food. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made 30 days after the initial treatment. Treated areas may be seeded 3 to 4 weeks after application.

The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application [(1.2 lb 2,4-D ae, 0.27 lb MCP-P ae and 0.11 lb dicamba ae per acre per application)]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. [(2.4 lb 2,4-D ae, 0.54 lb MCP-P ae and 0.22 lb dicamba ae per acre)].

**[Note to Reviewer: The paragraph above was moved - not added or deleted.]**

**[Note to Supplemental Registrants: Guarantees for plant nutrients other than nitrogen, phosphorous, and potassium may be permitted or required by regulation by the \_\_\_\_\_. The guarantees for such other nutrients shall be expressed in the form of the element. The source (oxides, salts, chelates, etc.) of such other nutrients may be required to be stated and may be included on the label.]**

**[Note to Reviewer: The section above was incorporated into the Fertilizer Guaranteed Analysis in the appendix]**

**[For state registrations in Kentucky, add the following paragraph:]**

**A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing (\*Number of pounds) of nitrogen per 1,000 square feet. Please consult your local extension office for questions regarding a regular fertilizer program suitable for your geographic region.]**

### **Labeling Requirements:**

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.htm>.

**[Note to Reviewer: The section above was moved to the appendix]**

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when the dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

<b>Broadleaf Weeds Controlled [248 Weeds, 233 Weeds CA]</b>			
Amaranth, purple	Creeping Beggarweed*	Morningglory, mexican	Speedwell, bi-lobed
Amaranth, slender	Creeping Jenny	Morningglory, red	Speedwell, common
Amaranth, spiny	(creeping charlie)	Morningglory, tall	Speedwell, corn
Aster, heath	Cupid shaving Brush*	Morningglory, wolly	Speedwell, ivyleaf
Aster, white heath	Cudweed	Mugwort	Speedwell, Persian
Aster, purple	Daisy, English	Mustard, ball	Speedwell, purslane
Aster, white prairie	Daisy, oxeye	Mustard, black	Speedwell, slender
Aster, wild	Dandelion, common	Mustard, blue	Speedwell, snow
Artichoke*	Day flower*	Mustard, hedge	Speedwell, thymeleaf
Austrian fieldcress	Deadnettle	Mustard, Indian,	Spurge, spotted
Bedstraw, Catchweed	Dichondra	Mustard, tansy	Spurge, prostrate
Bedstraw, smooth	Dock, broadleaf	Mustard, tumble	Star of Bethlehem*
Bedstraw, Northern	Dock, curly	Mustard, white	Strawberry, India mock
Bedstraw, pineywoods	Dock, smooth	Mustard, wormseed	Tansy ragwort
Beggarticks, hairy	Dogbane	Nettle, stinging (burning)	Tanweed
Betony, Florida	Dogfennel	Nettle, tall	Thistle, blessed
Bindweed, Field	Dollarweed	Nettle, wood	Thistle, bull
Bindweed, Hedge	(Pennywort)	Parsley-piert	Thistle, Canada
Bitter Wintercress	Dove weed	Parsnip	Thistle, flodman (prairie)
Bittercress, hairy	Elderberry	Pearlwort	Thistle, musk
Bitterweed	Falseflax	Pennycress (frenchweed)	Thistle, platte
Black-eyed Susan	False sunflower	Pepperweed, perennial	Thistle, plumeless
Black Medic	Fiddleneck	Pepperweed, Virginia	Thistle, Russian
Blood flower milkweed	Filaree, redstem	(Peppergrass)	Thistle, scotch
Blue lettuce	Filaree, whitestem	Pigweed, prostrate	Thistle, wavyleaf
Brassbuttons	Fleabane, daisy	Pigweed, redroot	Thistle, yellow (horrible)
Bristly oxtongue	(annual)	Pigweed, Russian	Trailing crownvetch
Broomweed	Fleabane, hairy	Pigweed, smooth	Velvetleaf
Bullnettle	Fleabane, rough	Pigweed, tumble	Vervain, blue
Burclover	Galinsoga, hairy	Pineapple weed	Vervain, hoary
Burdock	Galinsoga, smallflower	Plains coreopsis (tickseed)	Vervain, prostrate
Burweed, lawn (spurweed)	Goldenrod	Plantain, blackseed	Vervain, tall
Buttercup, bulbous	Ground ivy	Plantain, bracted	Vetch, bird
Buttercup, creeping	Groundsel*	Plantain, broadleaf	Vetch, hairy
Buttonweed, common	Gumweed	Plantain, buckhorn	Vetch, milk
(poorjoe)	Hawkweed, meadow	Plantain, hoary	Vetch, narrowleaf
Buttonweed, Virginia	Hawkweed, mouseear	Plantain, narrowleaf	Vetch, wild
Carolina geranium (wild)	Hawkweed, orange	Plantain, slender	Virginia creeper
Carpetweed	Hawkweed, yellow	Plantain, woolly	Western clematis
Catnip	Healall	Poison ivy	Western salsify
Catsear, spotted (false	Heartleaf drymary	Poison oak	Wild buckwheat
dandelion)	Hemp	Pokeweed	Wild carrot
Chamber bitter*	Henbit	Prairie Sunflower	Wild four-o'clock
Chickweed, common	Hoary cress	Prickly lettuce (compass plant)	Wild garlic
Chickweed, mouseear	Horsenettle	Prickly sida	Wild lettuce
Chickweed, sticky	Horseweed*	Puncturevine	Wild marigold

Chicory	Jimsonweed	Purslane, common	Wild mustard (yellow rocket)
Cinquefoil	Johnny-Jumpup violet*	Pusley, Brazil	Wild onion
Cinquefoil, rough	Knawel	Pusley, Florida	Wild parsnip
Clover, alcyon	Knotweed, prostrate	Pusley, large flower	Wild radish
Clover, crimson	Kochia	Ragweed, bur	Wild rape
Clover, hop	Lambsquarters	Ragweed, common	Wild strawberry
Clover, rabbit foot	Lespedeza	Ragweed, lanceleaf	Wild sweet potato
Clover, red	Mallow, alkali	Ragweed, western	Wild violet
Clover, strawberry	Mallow, bristly	Scarlet pimpernel	Woodsorrel, creeping
Clover, yellow sweet	Mallow, common	Shepard's purse	Woodsorrel, violet
Clover, white sweet	Mallow, dwarf	Smartweed, ladysthumb*	Woodsorrel, yellow (Oxalis)
Clover, white	Mallow, venice	Smartweed, pale*	Woolly croton
Cockle	Marcela*	Smartweed, Pennsylvania*	Wormseed
Cocklebur	Matchweed	Smooth chaff-flower	Yarrow, common
Cocklebur, oriental	Mexicanweed	Sorrel, red (sheep)	Yarrow, western
Cocklebur, spiny	Moneywort*	Sowthistle, annual	Yarrow, yellow
Common mullein	Morningglory, bigroot	Sowthistle, spiny	
Corn Spurry*	Morningglory, ivyleaf	Spanish needles	

\*Not for use in California.

**[Note to Supplemental Registrants:** The following language on equipment settings/spreader settings is optional – it may be presented on the container at your discretion.]


#### SPREADERS AND CALIBRATION:

1. Drop spreaders and rotary spreaders must be calibrated for accurate application of this product.
2. Close spreader opening (orifice).
3. Adjust spreader setting using the Spreader Suggestions chart
4. CALIBRATION INSTRUCTIONS:
  - For 3.0 pounds of product per 1,000 square feet of turfgrass: Check you spreader on 330 square feet and adjust to apply at the rate of 1 pound of product per 330 square feet.
  - For 3.2 pounds of product per 1,000 square feet of turfgrass: Check you spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
  - For 3.6 pounds of product per 1,000 square feet of turfgrass: Check you spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
  - For 4.0 pounds of product per 1,000 square feet of turfgrass: Check you spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.
5. Fill spreader. Fill spreader on driveway or sidewalk to eliminate the chance of spillage on the lawn.
6. Apply the product to lawn. [Refer to the patch charts below:]


The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application [(1.2 lb 2,4-D ae, 0.27 lb MCPP p ae and 0.11 lb dicamba ae per acre per application)]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. [(2.4 lb 2,4 D ae, 0.54 lb MCPP p ae and 0.22 lb dicamba ae per acre)].

**[Note to Reviewer:** The paragraph above was moved into the Instructions section of the label – not added or deleted.]

Path A for centrifugal/rotary spreaders.



Path B for drop spreaders.




7. Close the spreader openings when turning to change directions or when stopping.
8. If using a centrifugal/rotary spreader, avoid spreading product into adjacent landscape, flowers or vegetable plantings.

NOTE: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 square feet.

NOTE: If your spreader model does not appear above, please call toll free 800-xxx-xxxx, [8 A.M. - 5 P.M. EDT, Monday - Friday] for additional model settings.

This \_\_\_ pound bag should covers approximately \_\_\_\_\_ square feet (\_\_\_ ft. x \_\_\_ ft.) Refer to the table and picture to determine the size of your lawn.

COVERAGE CHART	
_____	Length
X _____	Width
= _____	Total Sq. Ft.
- _____	Non-Lawn Areas
= _____	Total Sq. Ft. Coverage you need.



Application Rate = 3 lb of product per 1,000 square feet.

SUGGESTED SPREADER SETTINGS:			
Centrifugal/Rotary Spreaders	Application Rate 3 lb/1,000 sq.ft.	Drop Spreaders	Application Rate 3 lb/1,000 sq.ft.
Frank's Broadcast	xx	Frank's Drop	xx
Scotts Speedy Green	xx	Greenview II	xx
Radio Flyer #0047	xx	True Temper CD20	xx
True Temper Rotary CB2000	xx	Scotts Accugreen AG1	xx
Earthway Rotary #2050SU	xx	Earthway Drop #7300SU	xx

These spreader settings are approximate. Differences may occur due to the condition of spreader, speed at which applied and roughness of area treated. Check your spreader on 330 square feet and adjust to apply at the rate of 1 pound per 1,000 330 square feet.

[Note to Reviewer: The chart above was moved forward from the appendix – not added or deleted.]

24-4-8 GUARANTEED ANALYSIS	
Total Nitrogen (N) .....	24%
Water Insoluble Nitrogen .....	5%
Available Phosphate (P2O5) .....	4%
Soluble Potash (K2O) .....	8%

[Note to Reviewer: The chart above was incorporated into the Guaranteed Analysis in the appendix.]

[Note to Supplemental Registrants: Fertilizer sources may vary in micronutrient composition, and iron may discolor or stain concrete. For fertilizers containing iron: Sweep all fertilizer from concrete surfaces immediately after application. Do not wash this product from concrete surfaces because fertilizers which contain iron may cause discoloration upon oxidation.]

[Note to Reviewer: The paragraph above was incorporated into the Appendix, not deleted.]

[End Optional Text]

## **STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or [www.website.com](http://www.website.com) for disposal instructions. Never place unused product down any indoor or outdoor drain.

## **LIMITED WARRANTY AND DISCLAIMER**

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

### **[ALTERNATE LIMITED WARRANTY**

The Distributor warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. To the extent consistent with applicable law, neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Distributor, and Buyer assumes the risk of any such use.]

[Alternate for -2935]

[Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using the product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

**ALL STATEMENTS MADE HEREIN ARE SUBJECT TO APPLICABLE LAW, AND TO THE EXTENT THERE IS ANY INCONSISTENCY OR CONTENTION, APPLICABLE LAW SHALL GOVERN.**

The Directions for Use of the product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of many different factors including, without limitation, manner of use or application, weather, combination with other products, or crop conditions. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Manufacturer and Seller harmless from any claims relating to such factors.

**Seller warrants that this product conforms to the chemical description on the label. EXCEPT FOR THIS WARRANTY, THE PRODUCT IS FURNISHED "AS-IS," AND NEITHER SELLER NOR MANUFACTURER MAKES ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT; SELLER AND MANUFACTURER SPECIFICALLY DISCLAIM ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE BEYOND WHAT IS STATED ON THE LABEL. Buyer and User accept all risks arising from any use of this product, including without limitation, uses contrary to label**

**instructions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or Manufacturer.**

Neither Manufacturer nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE BUYER OR USER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT, OR, AT THE ELECTION OF MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

These Conditions of Sale and Limitation of Warranty and Liability shall be interpreted, unless otherwise required by the law of the state of purchase, in accordance with the laws of the State of California, excluding its conflicts of laws rules, and may not be amended by any oral or written agreement.]

[Alternate for-2935]

[Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using the product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of the product ~~should~~ must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of many different factors including, without limitation, manner of use or application, weather, combination with other products, or crop conditions. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Manufacturer and Seller harmless from any claims relating to such factors.

**Seller warrants that this product conforms to the chemical description on the label, EXCEPT FOR THIS WARRANTY, THE PRODUCT IS FURNISHED “AS-IS,” AND NEITHER SELLER NOR MANUFACTURER MAKES ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT; SELLER AND MANUFACTURER SPECIFICALLY DISCLAIM ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Buyer and User accept all risks arising from any use of this product, including without limitation users contrary to label instructions, under abnormal conditions not reasonably foreseeable to (or beyond the control of) Seller or Manufacturer.**

To the extent permitted by law, neither Manufacturer nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE BUYER OR USER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT, OR, AT THE ELECTION OF MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

These Conditions of Sale and Limitation of Warranty and Liability shall be interpreted in accordance with the laws of the State of California, excluding its conflicts of laws rules, and may not be amended by any oral or written agreement.]

[Alternate for -41948]

[Conditions of Sale – Limited Warranty and Limitations of Liabilities and Remedies

Notice: Read the warranty and limitations of liability and remedies BEFORE using this product. If the terms and conditions of sale of this product to you are not acceptable, return the product unopened, and the full purchase price will be refunded.

Pride Ag Resources, (the “Company”) warrants only that this product conforms to the chemical description on the label and, if properly applied in accordance with label instructions, is reasonably suited for the purpose of supplying plant nutrients to turf grasses for lawns and golf courses.

The effectiveness of this product is dependent upon a number of factors, including without limitation, failure to read and follow label instructions, failure to use correct applications practices or appropriate

application equipment, soil conditioners, and weather conditions. No agent or employee of the Company or the seller is authorized to amend the terms of the conditions of sale of the limited warranty set forth on this label or to make and presentation or recommendation different from or inconsistent with the label of this product.

Failure to follow label instructions may cause injury to plant, animals or the environment. People with highly sensitive skin may develop skin irritations after contact with some fertilizers. If you develop a skin irritation in connection with the use of this product, you should either wear protective clothing or discontinue use of the product.

The Company makes no other warranties or representation of any kind, express or implied concerning the product, including without limitation warranties of merchantability and fitness for a particular purpose, nor are any such warranties intended or implied.

The exclusive remedy against the Company for any cause of action relating to the handling or use of this product shall be limited to, at the Company's election, one of the following:

1. Refund of the purchase price paid by the buyer or user for product bought, or
2. Replacement of the product used.

To the fullest extent under applicable law, the company shall not be liable for special, indirect, incidental, consequential, or punitive damages or expenses of any nature. The Company and seller offer the Product for sale, and the buyer and the user purchase this Product and accept it, subject to the foregoing conditions of sale and limitation of warranty, liabilities and remedies. Further, to the fullest extent under applicable law, the buyer and the user assume all risk of use, storage, and handling of the Product.]

## APPENDIX

### 1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Weight: \_\_\_\_\_
- EPA Est. No. \_\_\_\_\_

### 21. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

- **Previously Approved Fertilizer Grades:** Fertilizer grades with nutrient contents expressed as the percentages of N, P<sub>2</sub>O<sub>5</sub>, and K<sub>2</sub>O include the following:

10-6-4	16-3-5	20-10-0 (1)	23-7-7	25-4-8	28-3-7	30-4-8
10-16-20	16-6-9	20-10-5	24-3-7	26-3-3	28-3-8	32-4-8
10-18-10	18-3-5	21-5-7	24-4-8	26-3-5	29-3-4	22-3-14
12-5-5	18-6-12	22-6-6	25-3-3	26-4-6	30-3-4	30-3-4
15-5-10	20-4-6	22-7-7	25-3-5	26-4-13	30-3-5	
16-2-3	20-4-8	22-10-10	25-3-8	28-3-3	30-3-10	

The fertilizer grade guaranteed analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations. The following guaranteed analyses presentations have previously been submitted as notifications:

10-16-20 GUARANTEED ANALYSIS	
Total Nitrogen (N) .....	10%
6.0% Ammoniacal Nitrogen	
4.0% Urea Nitrogen	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) .....	16%
Soluble Potash (K <sub>2</sub> O) .....	20%
Magnesium (Mg) .....	1.0%
0.1% Water Soluble Magnesium (Mg)	
Sulfur (S) .....	2.0%

2.0% Combined Sulfur (S)	
Iron (Fe) .....	4.0%
Manganese (Mn) .....	0.5%
0.1% Water Soluble Manganese (Mn)	
Nutrient Sources: Urea, Ammonium Phosphate, Ammonium Sulfate, Muriate of Potash, Ferrous Sulfate, Manganese Oxide, Manganese Sulfate, Sulfate of Potash Magnesia and Magnesium oxide.	

<b>29-3-4 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	29%
1.1% Ammoniacal Nitrogen	
27.9% Urea Nitrogen	
Available Phosphate (P2O5) .....	3%
Soluble Potash (K2O) .....	4%
Derived from: Polymer-coated urea, urea, diammonium phosphate, and muriate of potash. *11.0% slowly available urea nitrogen from polymer coated urea.	
Chlorine (Cl), not more than .....	4.0%

<b>10-18-10 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	10%
10.0% Ammoniacal Nitrogen	
Available Phosphate (P2O5) .....	18%
Soluble Potash (K2O) .....	10%
Sulfur (S) .....	2.0%
5.0% Combined Sulfur (S)	
Nutrient Sources: Ammonium Sulfate, Ammonium Phosphate and Muriate of Potash.	

<b>26-4-6 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	26%
1.8% Ammoniacal Nitrogen	
24.4% Urea Nitrogen	
Available Phosphoric Acid (P2O5) .....	4%
Soluble Potash (K2O) .....	6%
Boron (B) ..... 0.02%	
Copper (Cu) ..... 0.05%	
Iron (Fe) ..... 1.00%	
Total Manganese (Mn) ..... 0.06%	
Molybdenum (Mo) ..... 0.0005%	
Zinc (Zn) ..... 0.05%	
Primary Plant Food Sources: Sulfur coated urea, urea, diammonium phosphate, muriate of potash. Secondary Plant Food Sources: Fertilizer grade borate, sulfates and oxides of copper, sulfates and oxides of iron, oxides of manganese, fertilizer grade molybdate, and sulfates and oxides of zinc. *4.0% slowly available urea nitrogen from sulfur coated urea.	
Chlorine (Cl), not more than .....	4.8%

<b>28-3-7 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	28%
1.2% Ammoniacal Nitrogen	
26.8% Urea Nitrogen	
Available Phosphoric Acid(P2O5) .....	3%
Soluble Potash (K2O) .....	7%
Primary Plant Food Sources: Polymer-coated urea, urea, diammonium phosphate, muriate of potash. *7.0% slowly available urea nitrogen from polymer-coated urea.	
Chlorine (Cl), not more than .....	6.4%

**Herbicide Benefit Claims:**

- With Trimec® Herbicide
- Controls/Kills dandelion, chickweed, knotweed, plantain, henbit, spurge and many other **[[listed]]** broadleaf weeds; ~~some of which are listed on the back panel~~
- Controls/Kills over 30 varieties of **[[listed]]** weeds while feeding your lawn.
- Controls/Kills dandelion, plantain, and other **[[listed]]** weeds.
- Controls **(Number)** weeds! (Listing of weeds may include the following.)
- Kills dandelion, chickweed, clover and most common lawn weeds **[as listed]**.
- Controls **most** annual broadleaf weeds **[as listed]**
- Kills **[Controls] XXX [Plus] [Listed] [Broadleaf] Weeds [As Listed]**



- Controls **(Number)** weeds! (Listing of weeds may include the following.)
- Application Rate = Apply 4 lbs. of GORDON'S TRIMEC® WEED & FEED 30 24-4-8 per 1,000 sq.ft. or 175 lbs. per acre (1 lb. of nitrogen per 1,000 sq.ft.)

<b>WEEDS</b>			
Annual yellow sweetclover	Cockle	Mustard	Spiny cocklebur
Aster	Cocklebur	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Common mullein	Narrowleaf vetch	Spotted catsear

**WEEDS**

Bedstraw	Creeping Charlie	Nettle	Spotted spurge
Beggarticks	Creeping jenny	Orange hawkweed	Spurweed
Betony, Florida	Cudweed	Oriental cocklebur	Stinging nettle
Bindweed, field	Curly dock	Oxalis	Strawberry, India mock
Bird vetch	Daisy, English	Parsley-piert	Tall nettle
Bitter wintercress	Daisy fleabane	Parsnip	Tall vervain
Bittercress, hairy	Daisy, oxeye	Pearlwort	Tansy ragwort
Bitterweed	Dandelion	Pennycress	Tansy mustard
Black-eyed Susan	Dichondra	Pepperweed	Tanweed
Black medic	Dogfennel	Pigweed	Thistle
Black mustard	Dollarweed	Pineywoods bedstraw	Trailing crownvetch
Blackseed plantain	False dandelion	Plains coreopsis	Tumble mustard
Blessed thistle	Falseflax	Plantain	Tumble pigweed
Bloodflower milkweed	False sunflower	Poison ivy	Velvetleaf
Blue lettuce	Fiddleneck	Poison oak	Venice mallow
Blue vervain	Florida pusley	Pokeweed	Virginia buttonweed
Bracted plantain	Frenchweed	Poorjoe	Virginia creeper
Bristly ox tongue	Galinsoga	Prairie sunflower	Virginia pepperweed
Broadleaf dock	Goldenrod	Prickly lettuce	Wavyleaf bullthistle
Broadleaf plantain	Ground ivy	Prickly sida	Western clematis
Broomweed	Gumweed	Prostrate knotweed	Western salsify
Buckhorn	Hairy fleabane	Prostrate pigweed	White mustard
Buckhorn plantain	Hawkweed	Prostrate spurge	Wild aster
Bulbous buttercup	Healall	Prostrate vervain	Wild buckwheat
Bull thistle	Heartleaf drymary	Puncturevine	Wild carrot
Bullnettle	Heath aster	Purslane, common	Wild four o'clock
Burclover	Hedge bindweed	Ragweed	Wild garlic
Burdock	Hedge mustard	Red sorrel	Wild geranium
Burning nettle	Henbit	Redroot pigweed	Wild lettuce
Burweed	Hoary cross	Redstem filaree	Wild marigold
Buttercup	Hoary plantain	Rough cinquefoil	Wild mustard
Buttonweed	Hoary vervain	Rough fleabane	Wild onion
Canada thistle	Horsenettle	Russian pigweed	Wild parsnip
Carolina geranium	Jimsonweed	Russian thistle	Wild radish
Carpetweed	Knawel	Scarlet pimpernel	Wild rape
Catchweed bedstraw	Knotweed	Scotch thistle	Wild strawberry
Catnip	Kochia	Sheep sorrel	Wild sweet potato
Catsear	Lambsquarters	Shepherd's purse	Wild vetch
Chickweed, common	Lespedeza	Slender plantain	Woodsorrel
Chickweed, mouseear	Mallow	Smallflower galinsoga	Woolly croton
Chicory	Matchweed	Smooth dock	Woolly morningglory
Cinquefoil	Mexicanweed	Smooth pigweed	Woolly plantain
Clover, crimson	Milk vetch	Serrel	Wormseed
Clover, hop	Morningglory	Sowthistle	Yarrow
Clover, red	Mouseear hawkweed	Spanish needles	Yellow rocket
Clover, strawberry	Mugwort	Speedwell	Yellowflower pepperweed
Clover, sweet	Musk thistle	Spiny amaranth	and other broadleaf weeds
Clover, white			

**Fertilizer Benefit Claims:**

- An 18-6-12 fertilizer provides slow-release feeding to promote winter hardiness and drought tolerance in lawns.
- Feeds your lawn for up to 12 weeks.
- Feeds the lawn with two (2) types of nitrogen and is pelleted for easy application.
- Triple Action Feeding works at every level.
  - Grass- Greener, Thicker lawn
  - Roots – Stronger Root System
  - Soil – Enriches & Fortifies Soil
- Triple Action Feeding works at every level for a Deep-Down Green!

- Why is Triple Action better? By feeding at all levels, your lawn receives the primary nutrients it needs to thrive. This ensures thicker, greener grass, stronger root system, and richer soil – the key to building a lush lawn that lasts.
- Grass: Builds thicker, greener grass and enhances overall appearance.
- Roots: Builds stronger, deeper root systems which transport nutrients from the soil to the grass to help lawn withstand heat-stress and drought.
- Soil: Enriches and fortifies your soil. Rich, fertile soil provides a foundation for establishing roots and replenishes vital primary nutrients.
- Nourishes all Winter for a quick Spring green-up.
- Premium slow release fertilizer stimulates grass growth to quickly conceal where weeds have died.
  - NITROGEN for Dense Green Lawns
  - PHOSPHORUS Improves Root Growth
  - POTASSIUM Vital for Growth and Drought Resistance
- Feeds your lawn for [up to] XX weeks.
- Zero Phosphate Formula
- The Easy Way To A Greener, Thicker Lawn
- Grow plants faster
- Promotes root growth for a thicker, greener lawn
- Enhances nutrient absorption
- Larger and stronger root system results in a more robust plant [lawn] and reduces the negative effects of stress such as drought on the turf [lawn]
- Formulated to release nutrients [fertilizer] to the plants for easy absorption [uptake]
- Increase nutrient uptake in grass up to 20%
- More nutrient absorption leads to stronger and deeper root systems that can take in water deeper from the soil resulting in a more robust lawn that is more drought resistant and requires less additional watering
- Allows for easier nutrient uptake into the plant [lawn] thus promoting stronger and deeper rooting
- Enhances nutrient absorption for stronger, deeper roots
- NutriLife [GreenSmart]-nourished lawns are more robust because their root systems are larger, deeper and better developed. Vigorous roots mean the lawns are hardier and better able to tolerate the stress that comes with heat and drought.
- Lawn and plants nourished with [company name's] Green Smart [GreenSmart] are able to tolerate the stress from heat and drought. Green Smart [GreenSmart] produces large, deep, and well developed root systems that can retain water. This enables lawns and plants to withstand stressful conditions and reduces the stress on you in caring for them. Green Smart [GreenSmart] produces thick, green lawns
- Enriches Soil to Promote a Vibrant, Hardy Lawn/Effective, Easy Way to Enhance the Soil
- Enriches Soil to Promote a Vibrant, Hardy Lawn
- Effective, Easy Way to Enhance the Soil
- Recharges Your Soil With Nutrients!
- [Company name's] Green Smart [GreenSmart] recharges your soil by enriching it with beneficial micro-organisms and nourishing them, making it easy for lawns and plants to absorb nutrient and use them efficiently. Green Smart [GreenSmart] enhances poor soil, creating an environment in which lawns and plants can thrive. Green Smart [GreenSmart] Formula produces thick, green lawns
- Beneath your grass lies a dynamic community of microorganisms vital to the lawn lifecycle and sustainability. Poor soil may have fewer of the nutrients microorganisms need to promote a vigorous lawn. GreenSmart works by enriching your soil with key microbes. GreenSmart revitalizes the soil by nourishing beneficial microorganisms, which aid in the development of a larger, deeper and better root system. The enhanced habitat is proven to produce a green, thick, vigorous lawn, and reduces the need for watering.
- GreenSmart creates the condition in the soil for the development of a large, extensive root system, by making it easier for the lawn to absorb nutrients and use them more efficiently. A vigorous root system enhances overall plant vigor, resulting in thick, green lawns.
- [Fertilizer Grade] Broadleaf Treatment with [ Other Fertilizer Nutrients]

- [Augusta Green™] “America’s Premium Turf Fertilizer”
- Phosphorus Free! [For use only on products containing no phosphorus]

**General Marketing Claims:**

- Builds a better lawn – Guaranteed
- (Fifteen, Sixteen, Eighteen or Twenty) Pounds Covers Up To 5,000 Square Feet.
- Greens lawns and develops deep roots while controlling weeds.
- Bring out the best with the turf magic touch!
- Gordon’s Trimec Weed & Feed 30 kills unsightly weeds such as dandelion, chickweed and clover while it nourishes your lawn. Its Triple Action formula contains slow-release nitrogen which continuously feeds the grass, roots and soil for up to two full months [all winter long]. And it’s guaranteed not to burn. The result is a greener, thicker lawn [this Fall and quicker green-up next Spring]. You’re just one application away from a better lawn. That’s the [company name] Better Lawn Guarantee.
- For questions or comments, call toll-free 1-877-bayerAL (1-877-229-3725)
- Made in the U.S.A.
- **Manufactured in the U.S.A.**
- Kills weeds and builds a thick, green Fall lawn.



- For Use In Spring Or When Weeds Are Present
- Covers [Up To] XXXX sq.ft.
- Care And Handling
- Quality Results
- Use Instructions
- Contains GreenSmart [NutraLife]
- Reduces Stress on Lawns, Protects from Drought and Heat
- For chemical spills, leaks, fire or exposure, call CHEMTREC: 800-424-9300
- Controls listed broadleaf weeds while feeding your lawn

Application Rate = 3 lb of product per 1,000 square feet.

<b>SUGGESTED SPREADER SETTINGS:</b>			
<b>Centrifugal/Rotary Spreaders</b>	<b>Application Rate 3 lb/1,000 sq.ft.</b>	<b>Drop Spreaders</b>	<b>Application Rate 3 lb/1,000 sq.ft.</b>
Frank’s Broadcast	xx	Frank’s Drop	xx
Scotts Speedy Green	xx	Greenview II	xx
Radio Flyer #0047	xx	True Temper CD20	xx
True Temper Rotary CB2000	xx	Scotts Accugreen AG4	xx
Earthway Rotary #2050SU	xx	Earthway Drop #7300SU	xx

These spreader settings are approximate. Differences may occur due to the condition of spreader, speed at which applied and roughness of area treated. Check your spreader on 330 square feet and adjust to apply at the rate of 4 pound per 1,000-330 square feet.

**2. State mandated fertilizer labeling requirements:**

[Note to Supplemental Registrants: Please select the statements relevant to the areas of the country in which your specific product is registered and sold; if any of these statements are

relevant to your labeling, please place in the Directions for Use under the heading “State Restrictions”]

<b>GUARANTEED ANALYSIS XX-X-X</b>	
Total Nitrogen (N) .....	XX%
x.x% ammoniacal nitrogen	
x.x% urea nitrogen*	
Available Phosphate (P2O5) .....	X%
Soluble Potash (K2O) .....	X%
Sulfur (S) .....	X%
x.x% free sulfur (S)	
Iron (Fe).....	X%
Derived from: polymer-coated sulfur-coated urea, urea, ammonium phosphate, potassium chloride.	
*A portion of the urea has been coated to provide x.x% coated slow-release nitrogen.	
This product and/or its manufacture is covered by one or more of the following U.S. patents and international counterparts thereof: US Patent Numbers: xxxxxx	

[Note: Fertilizer grades with nutrient contents expressed as the percentages of N, P2O5 and K2O may include the following: 23-3-7 or x-x-x. The fertilizer grade analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.]

- California, Oregon, Washington: “Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>”
- Florida: [Note to reviewer: maximum number of applications in this section is dually based upon maximum ai load as well as fertilizer analysis effect on maximum nitrogen rates in different regions in Florida]:
  - In North Florida: Do not exceed X applications per year on bermudagrass. Do not exceed X applications per year on centipedegrass. Do not exceed X applications per year on St. Augustine grass. Do not exceed X applications per year on zoysiagrass.
  - In Central Florida: Do not exceed X applications per year on bermudagrass. Do not exceed X applications per year on centipedegrass. Do not exceed X applications per year on St. Augustine grass. Do not exceed X applications per year on zoysiagrass.
  - In South Florida: Do not exceed X applications per year on bermudagrass. Do not exceed X applications per year on centipedegrass. Do not exceed X applications per year on St. Augustine grass. Do not exceed X applications per year on zoysiagrass.
- Florida: “Apply only to actively growing turf.”
- Florida: “Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices. Check with your county or city government to determine if there are local regulations for fertilizer use.
- Idaho: “Apply only as directed”
- Maryland: “This fertilizer contains phosphorus and may not be used on turf in the state of Maryland except when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory identified under § 8-803.7 of the Agriculture Article, Annotated Code of Maryland, performed no more than 3 years before the application; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State; or 3) Reestablishing or repairing a turf area.”
- New Jersey: “In the state of New Jersey, this product may be used only when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary pursuant to a soil test conducted by a laboratory identified pursuant to subsection b. of section 6 of P.L.2010, c.112 (C.58:10A-66) and performed no more than three years before the application, and pursuant to the associated annual fertilizer recommendation issued by the New Jersey Agricultural Experiment Station at Rutgers, the State University; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is in accordance with the standards and requirements

- established under the "Soil Erosion and Sediment Control Act," P.L.1975, c.251 (C.4:24-39 et seq.) and the rules and regulations adopted pursuant thereto; 3) Reestablishing or repairing a turf area; or 4) Delivering liquid or granular fertilizer under the soil surface, directly to the feeder roots."
- New Hampshire: In the state of New Hampshire, this product may only be used on 1) Establishing new lawns, for repairing a lawn, for seeding, or for use when a soil test indicates a phosphorus deficiency.
  - Virginia: "In the state of Virginia, this product may be used only on 1) Newly established turf areas from sod or seed during the first growing season; 2) Turf areas being repaired or renovated, and; 3) Turf areas where soil tests performed within the past three years indicate nutrient deficiency."
  - [Virginia, Florida, Maryland, others] DO NOT APPLY NEAR WATER, STORM DRAINS, OR DRAINAGE DITCHES. DO NOT APPLY IF HEAVY RAIN IS EXPECTED. APPLY THIS PRODUCT ONLY TO YOUR LAWN/GARDEN. [AND SWEEP ANY PRODUCT THAT LANDS ON THE DRIVEWAY, SIDEWALK, OR STREET, BACK INTO YOUR LAWN/GARDEN.] [Note to Supplemental Registrant: The bracketed part of this statement is not required for liquid products.]
  - [For products containing Iron:] Keep this product off sidewalks, patios, driveways, etc. This fertilizer contains iron which will stain concrete and painted surfaces. If it comes in contact with concrete surface, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.
  - [Fertilizer sources may vary in micronutrient composition, and iron may discolor or stain concrete. For fertilizers containing iron:] Sweep all fertilizer from concrete surfaces immediately after application. Do not wash this product from concrete surfaces because fertilizers which contain iron may cause discoloration upon oxidation.]
  - [Note to Supplemental Distributor: In Kentucky if the product supplies less than 1lb of Nitrogen per application, include the following:] A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing XX pounds of nitrogen per 1,000 square feet [and herbicide to control listed weeds]. Please consult your local extension office for questions regarding a regular fertilizer program [suitable for your geographic region.]

**SUBLABEL 2: For use by supplemental distributor for Bayer Advanced Triple Action Lawn Fertilizer Plus Weed Control II**

**GORDON'S TRIMEC® WEED & FEED 30  
24-4-8**

**EPA Reg. No. 2217-579**

**ACTIVE INGREDIENT:**

2,4-dichlorophenoxyacetic acid (CAS Registry No. 94-75-7) .....	0.69%
(+)-(R)-2-(4-chloro-2-methylphenoxy) propionic acid (CAS Registry No. 16484-77-8) .....	0.15%
Dicamba: 3,6-dichloro-2-methoxybenzoic acid (CAS Registry No. 1918-00-9) .....	0.06%
<b>INERT INGREDIENTS:</b> .....	<b>99.10%</b>
<b>TOTAL</b> .....	<b>100.00%</b>

**THIS PRODUCT CONTAINS:**

- 13.8 lb 2,4-dichlorophenoxyacetic acid equivalent per ton
- 3.0 lb (+)-(R)-2-(4-chloro-2-methylphenoxy) propionic acid equivalent per ton
- 1.2 lb 3,6-dichloro-2-methoxybenzoic acid equivalent per ton

Isomer Specific by AOAC Methods:

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals**

**CAUTION:** Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wear long-sleeved shirt, long pants, shoes, socks, and chemical resistant gloves made of any waterproof material. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

<b>First Aid</b>	
<b>If on skin or on clothing:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If inhaled:</b>	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call toll free, 1-877-800-5556 for emergency medical treatment information.</p>	

### **Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

### **[Environmental Hazards (alternate per PRN 2008-1)]**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run-off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### **PRECAUTIONS AND RESTRICTIONS:**

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Do not use on dichondra, carpetgrass, closely mowed bentgrass putting greens or St. Augustinegrass. Do not use this product where desirable clovers are present. Do not exceed specified dosages for any area. Apply when the air is calm. Application of this product to newly seeded areas should be delayed until the third mowing.
- Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, because injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not

over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

**INSTRUCTIONS:**

Gordon's Trimec® Weed & Feed 30 contains Trimec® herbicide. Trimec® controls a wide range of lawn weeds, including dandelion, chickweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a (X-X-X analysis of fertilizer grade) fertilizer provides a supply of plant food. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days. Treated areas may be seeded 3 to 4 weeks after application.

**[Note to Supplemental Registrants:** Guarantees for plant nutrients other than nitrogen, phosphorous, and potassium may be permitted or required by regulation by the \_\_\_\_\_. The guarantees for such other nutrients shall be expressed in the form of the element. The source (oxides, salts, chelates, etc.) of such other nutrients may be required to be stated and may be included on the label.]

**[For state registrations in Kentucky, add the following paragraph:**

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing (\*Number of pounds) of nitrogen per 1,000 square feet. Please consult your local extension office for questions regarding a regular fertilizer program suitable for your geographic region.]

**Labeling Requirements:**

- Information regarding the contents and levels of metals in this product is available on the Internet as <http://www.aapfco.org/metals.htm>. [<http://www.regulatory-info-ba.com>]

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when the dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application (1.2 lb 2,4-D ae, 0.27 lb MCP-P p ae and 0.11 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. (2.4 lb 2,4-D ae, 0.54 lb MCP-P p ae and 0.22 lb dicamba ae per acre).

<b>WEEDS</b>			
Annual yellow sweetclover	Cocklebur	Narrowleaf plantain	Spiny cocklebur
Aster	Common mullein	Narrowleaf vetch	Spiny sowthistle
Austrian fieldcress	Creeping Charlie	Nettle	Spotted catsear
Bedstraw	Creeping jenny	Orange hawkweed	Spotted spurge
Beggarticks	Cudweed	Oriental cocklebur	Spurweed
Betony, Florida	Curly dock	Oxalis	Stinging nettle
Bindweed, field	Daisy, English	Parsley-piert	Strawberry, India mock
Bird vetch	Daisy fleabane	Parsnip	Tall nettle
Bitter wintercress	Daisy, oxeye	Pearlwort	Tall vervain
Bittercress, hairy	Dandelion	Pennycress	Tansy ragwort
Bitterweed	Dichondra	Pennywort	Tansy mustard
Black-eyed Susan	Dogbane	Peppergrass (Peppergrass)	Tanweed
Black medic	Dogfennel	Pepperweed	Thistle
Black mustard	Dollarweed	Pigweed	Trailing crownvetch
Blackseed plantain	Elderberry	Pineywoods bedstraw	Tumble mustard

**WEEDS**



Blessed thistle	False dandelion	Plains coreopsis (tickseed)	Tumble pigweed
Bloodflower milkweed	False flax	Plantain	Velvetleaf
Blue lettuce	False sunflower	Poison ivy	Venice mallow
Blue vervain	Fiddleneck	Poison oak	Virginia buttonwood
Bracted plantain	Florida pusley	Pokeweed	Virginia creeper
Bristly oxtongue	Frenchweed	Poorjoe	Virginia pepperweed
Broadleaf dock	Galinsoga	Prairie sunflower	Water pennywort
Broadleaf plantain	Goldenrod	Prickly lettuce	Wavyleaf bullthistle
Broomweed	Ground ivy	Prickly sida	Western clematis
Buckhorn	Gumweed	Prostrate knotweed	Western salsify
Buckhorn plantain	Hairy fleabane	Prostrate pigweed	White mustard
Bulbous buttercup	Hawkweed	Prostrate spurge	Wild aster
Bull thistle	Healall	Prostrate vervain	Wild buckwheat
Bullnettle	Heartleaf drymary	Puncturevine	Wild carrot
Burclover	Heathaster	Purslane, common	Wild four o'clock
Burdock	Hedge bindweed	Ragweed	Wild garlic
Burning nettle	Hedge mustard	Red sorrel	Wild geranium
Burweed	Henbit	Redroot pigweed	Wild lettuce
Buttercup	Hoary cress	Redstem filaree	Wild marigold
Buttonwood	Hoary plantain	Rough cinquefoil	Wild mustard
Canada thistle	Hoary vervain	Rough fleabane	Wild onion
Carolina geranium	Horsenettle	Russian pigweed	Wild parsnip
Carpetweed	Jimsonweed	Russian thistle	Wild radish
Catchweed bedstraw	Knawel	Scarlet pimpernel	Wild rape
Catnip	Knotweed	Scotch thistle	Wild strawberry
Catsear	Kochia	Sheep sorrel	Wild sweet potato
Chickweed, common	Lambsquarters	Shepherdspurse	Wild vetch
Chickweed, mouseear	Lespedeza	Slender plantain	Woodsorrel
Chicory	Mallow	Smallflower galinsoga	Woolly croton
Cinquefoil	Matchweed	Smooth dock	Woolly morningglory
Clover, crimson	Mexicanweed	Smooth pigweed	Woolly plantain
Clover, hop	Milk vetch	Sorrel	Wormseed
Clover, red	Morningglory	Sowthistle	Yarrow
Clover, strawberry	Mouseear hawkweed	Spanishneedles	Yellow rocket
Clover, sweet	Mugwort	Speedwell	Yellowflower pepperweed
Clover, white	Musk thistle	Spiny amaranth	and other broadleaf weeds
Cockle	Mustard		

**SPREADERS AND CALIBRATION:**

1. Drop spreaders and rotary spreaders must be calibrated for uniform application of this product.
2. Close spreader opening (orifice).
3. Adjust spreader setting using the Spreader Suggestions chart
4. CALIBRATION INSTRUCTIONS:
  - For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 330 square feet and adjust to apply at the rate of 1 pound of product per 330 square feet.
  - For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
  - For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
  - For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.
5. Fill spreader. Fill spreader on driveway or sidewalk to eliminate the chance of spillage on the lawn.
6. Apply the product to lawn. Refer to the patch charts below:

Path A for centrifugal/rotary spreaders.

Path B for drop spreaders.



7. Close the spreader opening when turning to change directions or when stopping.


8. If using a centrifugal/rotary spreader, avoid spreading product into adjacent landscape, flowers or vegetable plantings.

NOTE: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 square feet.

NOTE: If your spreader model does not appear above, please call toll free 800-xxx-xxxx, [8 A.M. - 5 P.M. EDT, Monday - Friday] for additional model settings.

This \_\_\_\_\_ pound bag should cover approximately \_\_\_\_\_ square feet (\_\_\_\_ ft. x \_\_\_\_ ft.). Refer to the table and picture to determine the size of your lawn.

COVERAGE CHART	
_____	Length
X _____	Width
= _____	Total Sq. Ft.
- _____	Non-Lawn Areas
= _____	Total Sq. Ft. Coverage you need.



#### 24-4-8 GUARANTEED ANALYSIS

Total Nitrogen (N) .....	24%
Water Insoluble Nitrogen .....	5%
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) .....	4%
Soluble Potash (K <sub>2</sub> O) .....	8%

**[Note to Supplemental Registrants:** Fertilizer sources may vary in micronutrient composition, and iron may discolor or stain concrete. For fertilizers containing iron: Sweep all fertilizer from concrete surfaces immediately after application. Do not wash this product from concrete surfaces because fertilizers which contain iron may cause discoloration upon oxidation.]

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or [www.website.com](http://www.website.com) for disposal instructions. Never place unused product down any indoor or outdoor drain.

## LIMITED WARRANTY AND DISCLAIMER

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once to the seller.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

## APPENDIX

### 1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Weight: \_\_\_\_\_
- EPA Est. No. \_\_\_\_\_

### 2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

- **Previously Approved Fertilizer Grades:** Fertilizer grades with nutrient contents expressed as the percentages of N,P2O5, and K2O include the following:

10-6-4	16-3-5	20-10-0 (1)	23-7-7	25-4-8	28-3-7	30-4-8
10-16-20	16-6-9	20-10-5	24-3-7	26-3-3	28-3-8	32-4-8
10-18-10	18-3-5	21-5-7	24-4-8	26-3-5	29-3-4	
12-5-5	18-6-12	22-6-6	25-3-3	26-4-6	30-3-4	
15-5-10	20-4-6	22-7-7	25-3-5	26-4-13	30-3-5	
16-2-3	20-4-8	22-10-10	25-3-8	28-3-3	30-3-10	

The fertilizer grade guaranteed analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations. The following guaranteed analyses presentations have previously been submitted as notifications:

<b>10-16-20 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	10%
— 6.0% Ammoniacal Nitrogen	
— 4.0% Urea Nitrogen	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) .....	16%
Soluble Potash (K <sub>2</sub> O) .....	20%
Magnesium (Mg) .....	1.0%
— 0.1% Water Soluble Magnesium (Mg)	
Sulfur (S) .....	2.0%
— 2.0% Combined Sulfur (S)	
Iron (Fe) .....	1.0%
Manganese (Mn) .....	0.5%
— 0.1% Water Soluble Manganese (Mn)	
Nutrient Sources: Urea, Ammonium Phosphate, Ammonium Sulfate, Muriate of Potash, Ferrous Sulfate, Manganese Oxide, Manganese Sulfate, Sulfate of Potash Magnesia and Magnesium oxide.	

<b>29-3-4 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	20%
— 1.1% Ammoniacal Nitrogen	
— 27.9% Urea Nitrogen	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) .....	3%
Soluble Potash (K <sub>2</sub> O) .....	4%
Derived from: Polymer coated urea, urea, diammonium phosphate, and muriate of potash. *11.0% slowly available urea nitrogen from polymer coated urea.	
Chlorine (Cl), not more than .....	4.0%

<b>10-18-10 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	10%
— 10.0% Ammoniacal Nitrogen	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) .....	18%
Soluble Potash (K <sub>2</sub> O) .....	10%
Sulfur (S) .....	2.0%
— 5.0% Combined Sulfur (S)	
Nutrient Sources: Ammonium Sulfate, Ammonium Phosphate and Muriate of Potash.	

<b>26-4-6 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	26%
1.8% Ammoniacal Nitrogen	
24.4% Urea Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) .....	4%
Soluble Potash (K <sub>2</sub> O) .....	6%
Boron (B) ..... 0.02%	
Copper (Cu) ..... 0.05%	
Iron (Fe) ..... 1.00%	
Total Manganese (Mn) ..... 0.06%	
Molybdenum (Mo) ..... 0.0005%	
Zinc (Zn) ..... 0.05%	
Primary Plant Food Sources: Sulfur coated urea, urea, diammonium phosphate, muriate of potash.	
Secondary Plant Food Sources: Fertilizer grade borate, sulfates and oxides of copper, sulfates and oxides of iron, oxides of manganese, fertilizer grade molybdate, and sulfates and oxides of zinc.	
*4.0% slowly available urea nitrogen from sulfur coated urea.	
Chlorine (Cl), not more than .....	4.8%

<b>28-3-7 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	28%
1.2% Ammoniacal Nitrogen	
26.8% Urea Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) .....	3%
Soluble Potash (K <sub>2</sub> O) .....	7%
Primary Plant Food Sources: Polymer-coated urea, urea, diammonium phosphate, muriate of potash.	
*7.0% slowly available urea nitrogen from polymer-coated urea.	
Chlorine (Cl), not more than .....	6.4%

**SUBLABEL 3: For use by supplemental distributor for  
Bayer Advanced Triple Action Winterizer Fertilizer Plus Weed Control II**

# **GORDON'S TRIMEC® WEED & FEED 30 24-4-8**

## **EPA Reg. No. 2217-579**

### **ACTIVE INGREDIENT:**

2,4-dichlorophenoxyacetic acid (CAS Registry No. 94-75-7) .....	0.69%
(+)-(R)-2-(4-chloro-2-methylphenoxy) propionic acid (CAS Registry No. 16484-77-8) .....	0.15%
Dicamba: 3,6-dichloro-2-methoxybenzoic acid (CAS Registry No. 1918-00-9) .....	0.06%
<b>INERT INGREDIENTS:</b> .....	<b>99.10%</b>
<b>TOTAL</b> .....	<b>100.00%</b>

### **THIS PRODUCT CONTAINS:**

13.8 lb 2,4-dichlorophenoxyacetic acid equivalent per ton  
3.0 lb (+)-(R)-2-(4-chloro-2-methylphenoxy) propionic acid equivalent per ton  
1.2 lb 3,6-dichloro-2-methoxybenzoic acid equivalent per ton  
Isomer Specific by AOAC Methods:  
TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

**KEEP OUT OF REACH OF CHILDREN**

## **CAUTION**

**STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

## **PRECAUTIONARY STATEMENTS**

### **Hazards to Humans and Domestic Animals**

**CAUTION:** Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wear long-sleeved shirt, long pants, shoes, socks, and chemical resistant gloves made of any waterproof material. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

<b>First Aid</b>	
<b>If on skin or on clothing:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If inhaled:</b>	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call toll free, 1-877-800-5556 for emergency medical treatment information.</p>	

### **Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

### **[Environmental Hazards (alternate per PRN 2008-1)]**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run-off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### **PRECAUTIONS AND RESTRICTIONS:**

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Do not use on dichondra, carpetgrass, bentgrass or St. Augustinegrass. Do not use this product where desirable clovers are present. Do not exceed specified dosages for any area. Apply when the air is calm. Application of this product to newly seeded areas should be delayed until the third mowing.
- Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, because injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not

over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

**INSTRUCTIONS:**

Gordon's Trimec® Weed & Feed 30 contains Trimec® herbicide. Trimec® controls a wide range of lawn weeds, including dandelion, chickweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a (X-X-X analysis of fertilizer grade) fertilizer provides a supply of plant food. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in 30 days. Treated areas may be seeded 3 to 4 weeks after application.

**[Note to Supplemental Registrants:** Guarantees for plant nutrients other than nitrogen, phosphorous, and potassium may be permitted or required by regulation by the \_\_\_\_\_. The guarantees for such other nutrients shall be expressed in the form of the element. The source (oxides, salts, chelates, etc.) of such other nutrients may be required to be stated and may be included on the label.]

**[For state registrations in Kentucky, add the following paragraph:**

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing (\*Number of pounds) of nitrogen per 1,000 square feet. Please consult your local extension office for questions regarding a regular fertilizer program suitable for your geographic region.]

**Labeling Requirements:**

- Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.htm>. [<http://www.regulatory-info-ba.com>]

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when the dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application (1.2 lb 2,4-D ae, 0.27 lb MCPP-p ae and 0.11 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. (2.4 lb 2,4-D ae, 0.54 lb MCPP-p ae and 0.22 lb dicamba ae per acre).

<b>WEEDS</b>			
Annual yellow sweetclover	Cocklebur	Narrowleaf plantain	Spiny cocklebur
Aster	Common mullein	Narrowleaf vetch	Spiny sowthistle
Austrian fieldcress	Creeping Charlie	Nettle	Spotted catsear
Bedstraw	Creeping jenny	Orange hawkweed	Spotted spurge
Beggarticks	Cudweed	Oriental cocklebur	Spurweed
Betony, Florida	Curly dock	Oxalis	Stinging nettle
Bindweed, field	Daisy, English	Parsley-piert	Strawberry, India mock
Bird vetch	Daisy fleabane	Parsnip	Tall nettle
Bitter wintercress	Daisy, oxeye	Pearlwort	Tall vervain
Bittercress, hairy	Dandelion	Pennycress	Tansy ragwort
Bitterweed	Dichondra	Pennywort	Tansy mustard
Black-eyed Susan	Dogbane	Peppergrass (Peppergrass)	Tanweed
Black medic	Dogfennel	Pepperweed	Thistle
Black mustard	Dollarweed	Pigweed	Trailing crownvetch
Blackseed plantain	Elderberry	Pineywoods bedstraw	Tumble mustard

## WEEDS

Blessed thistle	False dandelion	Plains coreopsis (tickseed)	Tumble pigweed
Bloodflower milkweed	False flax	Plantain	Velvetleaf
Blue lettuce	False sunflower	Poison ivy	Venice mallow
Blue vervain	Fiddleneck	Poison oak	Virginia buttonwood
Bracted plantain	Florida pusley	Pokeweed	Virginia creeper
Bristly oxtongue	Frenchweed	Poorjoe	Virginia pepperweed
Broadleaf dock	Galinsoga	Prairie sunflower	Water pennywort
Broadleaf plantain	Goldenrod	Prickly lettuce	Wavyleaf bullthistle
Broomweed	Ground ivy	Prickly sida	Western clematis
Buckhorn	Gumweed	Prostrate knotweed	Western salsify
Buckhorn plantain	Hairy fleabane	Prostrate pigweed	White mustard
Bulbous buttercup	Hawkweed	Prostrate spurge	Wild aster
Bull thistle	Healall	Prostrate vervain	Wild buckwheat
Bullnettle	Heartleaf drymary	Puncturevine	Wild carrot
Burclover	Heathaster	Purslane, common	Wild four o'clock
Burdock	Hedge bindweed	Ragweed	Wild garlic
Burning nettle	Hedge mustard	Red sorrel	Wild geranium
Burweed	Henbit	Redroot pigweed	Wild lettuce
Buttercup	Hoary cross	Redstem filaree	Wild marigold
Buttonwood	Hoary plantain	Rough cinquefoil	Wild mustard
Canada thistle	Hoary vervain	Rough fleabane	Wild onion
Carolina geranium	Horsenettle	Russian pigweed	Wild parsnip
Carpetweed	Jimsonweed	Russian thistle	Wild radish
Catchweed bedstraw	Knawel	Scarlet pimpernel	Wild rape
Catnip	Knotweed	Scotch thistle	Wild strawberry
Catsear	Kochia	Sheep sorrel	Wild sweet potato
Chickweed, common	Lambsquarters	Shepherdspurse	Wild vetch
Chickweed, mouseear	Lespedeza	Slender plantain	Woodsorrel
Chicory	Mallow	Smallflower galinsoga	Woolly croton
Cinquefoil	Matchweed	Smooth dock	Woolly morningglory
Clover, crimson	Mexicanweed	Smooth pigweed	Woolly plantain
Clover, hop	Milk vetch	Sorrel	Wormseed
Clover, red	Morningglory	Sowthistle	Yarrow
Clover, strawberry	Mouseear hawkweed	Spanishneedles	Yellow rocket
Clover, sweet	Mugwort	Speedwell	Yellowflower pepperweed
Clover, white	Musk thistle	Spiny amaranth	and other broadleaf weeds
Cockle	Mustard		

## SPREADERS AND CALIBRATION:

1. Drop spreaders and rotary spreaders must be calibrated for uniform application of this product.
2. Close spreader opening (orifice).
3. Adjust spreader setting using the Spreader Suggestions chart.
4. CALIBRATION INSTRUCTIONS:
  - For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 330 square feet and adjust to apply at the rate of 1 pound of product per 330 square feet.
  - For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
  - For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
  - For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.
5. Fill spreader. Fill spreader on driveway or sidewalk to eliminate the chance of spillage on the lawn.
6. Apply the product to lawn. Refer to the patch charts below:

Path A for centrifugal/rotary spreaders.

Path B for drop spreaders.



7. Close the spreader openings when turning to change directions or when stopping.


8. If using a centrifugal/rotary spreader, avoid spreading product into adjacent landscape, flowers or vegetable plantings.

NOTE: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 square feet.

NOTE: If your spreader model does not appear above, please call toll free 800-xxx-xxxx, [8 A.M. - 5 P.M. EDT, Monday - Friday] for additional model settings.

This \_\_\_ pound bag should cover approximately \_\_\_ square feet (\_\_\_ ft. x \_\_\_ ft.) Refer to the table and picture to determine the size of your lawn.

COVERAGE CHART	
_____	Length
X _____	Width
= _____	Total Sq. Ft.
- _____	Non-Lawn Areas
= _____	Total Sq. Ft. Coverage you need.



24-4-8 GUARANTEED ANALYSIS	
Total Nitrogen (N) .....	24%
Water Insoluble Nitrogen .....	5%
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) .....	4%
Soluble Potash (K <sub>2</sub> O) .....	8%

**[Note to Supplemental Registrants:** Fertilizer sources may vary in micronutrient composition, and iron may discolor or stain concrete. For fertilizers containing iron: Sweep all fertilizer from concrete surfaces immediately after application. Do not wash this product from concrete surfaces because fertilizers which contain iron may cause discoloration upon oxidation.]

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or [www.website.com](http://www.website.com) for disposal instructions. Never place unused product down any indoor or outdoor drain.

## LIMITED WARRANTY AND DISCLAIMER

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once to the seller.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

## APPENDIX

### 1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Weight: \_\_\_\_\_
- EPA Est. No. \_\_\_\_\_

### 2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

- Previously Approved Fertilizer Grades: Fertilizer grades with nutrient contents expressed as the percentages of N,P2O5, and K5O include the following:

10-6-4	16-3-5	20-10-0-(1)	23-7-7	25-4-8	28-3-7	30-4-8
10-16-20	16-6-9	20-10-5	24-3-7	26-3-3	28-3-8	32-4-8
10-18-10	18-3-5	21-5-7	24-4-8	26-3-5	29-3-4	
12-5-5	18-6-12	22-6-6	25-3-3	26-4-6	30-3-4	
15-5-10	20-4-6	22-7-7	25-3-5	26-4-13	30-3-5	
16-2-3	20-4-8	22-10-10	25-3-8	28-3-3	30-3-10	

The fertilizer grade guaranteed analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations. The following guaranteed analyses presentations have previously been submitted as notifications:

<b>10-16-20 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	10%
— 6.0% Ammoniacal Nitrogen	
— 4.0% Urea Nitrogen	
Available Phosphate (P2O5) .....	16%
Soluble Potash (K2O) .....	20%
Magnesium (Mg) .....	1.0%
— 0.1% Water Soluble Magnesium (Mg)	
Sulfur (S) .....	2.0%
— 2.0% Combined Sulfur (S)	
Iron (Fe) .....	1.0%
Manganese (Mn) .....	0.5%
— 0.1% Water Soluble Manganese (Mn)	
Nutrient Sources: Urea, Ammonium Phosphate, Ammonium Sulfate, Muriate of Potash, Ferrous Sulfate, Manganese Oxide, Manganese Sulfate, Sulfate of Potash Magnesia and Magnesium oxide.	

<b>29-3-4 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	20%
— 1.1% Ammoniacal Nitrogen	
— 27.9% Urea Nitrogen	
Available Phosphate (P2O5) .....	3%
Soluble Potash (K2O) .....	4%
Derived from: Polymer-coated urea, urea, diammonium phosphate, and muriate of potash. *11.0% slowly available urea nitrogen from polymer coated urea.	
Chlorine (Cl), not more than .....	4.0%

<b>10-18-10 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	10%
— 10.0% Ammoniacal Nitrogen	
Available Phosphate (P2O5) .....	18%
Soluble Potash (K2O) .....	10%
Sulfur (S) .....	2.0%
— 5.0% Combined Sulfur (S)	
Nutrient Sources: Ammonium Sulfate, Ammonium Phosphate and Muriate of Potash.	

<b>26-4-6 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	26%
— 1.8% Ammoniacal Nitrogen	
— 24.4% Urea Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) .....	4%
Soluble Potash (K <sub>2</sub> O) .....	6%
Boron (B) .....	0.02%
Copper (Cu) .....	0.05%
Iron (Fe) .....	1.00%
Total Manganese (Mn) .....	0.06%
Molybdenum (Mo) .....	0.0005%
Zinc (Zn) .....	0.05%
Primary Plant Food Sources: Sulfur coated urea, urea, diammonium phosphate, muriate of potash. Secondary Plant Food Sources: Fertilizer grade borate, sulfates and oxides of copper, sulfates and oxides of iron, oxides of manganese, fertilizer grade molybdate, and sulfates and oxides of zinc. *4.0% slowly available urea nitrogen from sulfur coated urea.	
Chlorine (Cl), not more than .....	4.8%

<b>28-3-7 GUARANTEED ANALYSIS</b>	
Total Nitrogen (N) .....	28%
— 1.2% Ammoniacal Nitrogen	
— 26.8% Urea Nitrogen	
Available Phosphoric Acid(P <sub>2</sub> O <sub>5</sub> ) .....	3%
Soluble Potash (K <sub>2</sub> O) .....	7%
Primary Plant Food Sources: Polymer-coated urea, urea, diammonium phosphate, muriate of potash *7.0% slowly available urea nitrogen from polymer-coated urea.	
Chlorine (Cl), not more than .....	6.4%

- (Fifteen, Sixteen, Eighteen or Twenty) Pounds Covers Up To 5,000 Square Feet.
- Controls/Kills dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf weeds; some of which are listed on the back panel
- An 18-6-12 fertilizer provides slow-release feeding to promote winter hardiness and drought tolerance in lawns.
- Feeds your lawn for up to 12 weeks.
- Controls/Kills over 30 varieties of weeds while feeding your lawn.
- Greens lawns and develops deep roots while controlling weeds.
- Bring out the best with the turf magic touch!
- Controls/Kills dandelion, plantain, and other weeds.
- Feeds the lawn with two (2) types of nitrogen and is pelleted for easy application.
- Controls (Number) weeds! (Listing of weeds may include the following.)
- Kills dandelion, chickweed, clover and most common lawn weeds.
- Triple Action Feeding works at every level.
  - Grass— Greener, Thicker lawn
  - Roots— Stronger Root System
  - Soil— Enriches & Fortifies Soil
- Builds a better lawn — Guaranteed
- Gordon's Trimec Weed & Feed 30 kills unsightly weeds such as dandelion, chickweed and clover while it nourishes your lawn. Its Triple Action formula contains slow-release nitrogen which continuously feeds the grass, roots and soil for up to two full months [all winter long]. And it's guaranteed not to burn. The result is a greener, thicker lawn [this Fall and quicker green up next Spring]. You're just one application away from a better lawn. That's the [company name] Better Lawn Guarantee.

- Triple Action Feeding works at every level for a Deep-Down Green!
- Why is Triple Action better? By feeding at all levels, your lawn receives the primary nutrients it needs to thrive. This ensures thicker, greener grass, stronger root system, and richer soil—the key to building a lush lawn that lasts.
- Grass: Builds thicker, greener grass and enhances overall appearance.
- Roots: Builds stronger, deeper root systems which transport nutrients from the soil to the grass to help lawn withstand heat stress and drought.
- Soil: Enriches and fortifies your soil. Rich, fertile soil provides a foundation for establishing roots and replenishes vital primary nutrients.
- For questions or comments, call toll-free 1-877-bayerAL (1-877-229-3725)
- Made in the U.S.A.
- Kills weeds and builds a thick, green Fall lawn.
- Nourishes all Winter for a quick Spring green-up.



Application Rate = 3 lb of product per 1,000 square feet.

<b>SUGGESTED SPREADER SETTINGS:</b>			
<b>Centrifugal/Rotary Spreaders</b>	<b>Application Rate</b> 3 lb/1,000 sq.ft.	<b>Drop Spreaders</b>	<b>Application Rate</b> 3 lb/1,000 sq.ft.
Frank's Broadcast	xx	Frank's Drop	xx
Scotts Speedy Green	xx	Greenview II	xx
Radio Flyer #0047	xx	True Temper CD20	xx
True Temper Rotary CB2000	xx	Scotts Accugreen AG4	xx
Earthway Rotary #2050SU	xx	Earthway Drop #7300SU	xx

These spreader settings are approximate. Differences may occur due to the condition of spreader, speed at which applied and roughness of area treated. Check your spreader on 330 square feet and adjust to apply at the rate of 1 pound per 1,000-330 square feet.

## **DOCUMENT CONTROL INFORMATION**

1. **Unique Label Identifier:** 002217-00579.20170111.notif-proposed-highlighted
2. **Reason for Issue:** updated weeds list; additional warranty statements for distributors; typos; state fertilizer language; formatting + EPA Comments #1

<b>Internal Usage Notes</b>
Associated Container Label: Supplemental Distributors