

6/16/97

PM 23

2217-580

pg 124

GORDON'S TRIMEC® WEED & FEED 20 16-4-8

Controls Dandelion, Chickweed, Knotweed, Plantains, Henbit, Spurge
and many other broadleaf weeds including those listed on the back panel.

ACTIVE INGREDIENTS:

*2,4-dichlorophenoxyacetic acid.....	0.456%
**2-(2-methyl-4-chlorophenoxy)propionic acid.....	0.204%
***Dicamba: 3,6-dichloro-o-anisic acid.....	0.043%
INERT INGREDIENTS.....	99.297%

TOTAL 100.000%

THIS PRODUCT CONTAINS:

- *9.117 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.
- **4.081 lbs. 2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton.
- ***0.861 lb. 3,6-dichloro-o-anisic acid equivalent per ton.

Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

KEEP OUT OF REACH OF CHILDREN

CAUTION

Statement of Practical Treatment

In case of contact, wash skin with plenty of soap and water.
For eyes, flush with water for 15 minutes and get medical attention.
See back panel for additional Precautionary Statements.

97 MAY 29 A9:27

REC'D EPA/OPP/DPD1

NET WEIGHT (20, 25, 30, 40, or 50) POUNDS

576/ APXXXXXX
EPA REG. NO. 2217-580
EPA EST. NO. 2217-KS-1
MANUFACTURED BY:



ACCEPTED

JUN 16 1997

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 2217-580



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

ENVIRONMENTAL HAZARDS:

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, *to areas where surface water is present or to intertidal areas below the mean high tide mark.* Do not contaminate target area. Do not contaminate domestic or irrigation waters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.

PESTICIDE DISPOSAL: Securely wrap bag in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

USE PRECAUTIONS:

Do not use this product where desirable clovers are present or allow the material to wind drift onto flowers, vegetables or ornamental shrubbery. Do not apply in areas underlaid by roots of desirable trees or shrubs. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning.

INSTRUCTIONS:

Gordon's Trimec® Weed & Feed 20 controls a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantains, henbit, spurge and many others commonly found in home lawns. This product may be applied to established lawns of bermudagrass, bentgrass, bahiagrass, centipedegrass, Kentucky bluegrass, fescues, perennial ryegrass, zoysiagrass, and St. Augustinegrass. Do not apply to dichondra or carpetgrass. Do not apply this product to 'Floritam' St. Augustinegrass in Florida. In addition, a 16-4-8 fertilizer provides a balanced supply of plant nutrients or plant food, needed for a normal season's growth.

Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days.

The maximum application rate to turf is 1.2 pounds of 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

Spreader Settings	
Scott.....	6
Lawn Beauty.....	5
Jackson.....	7½
Sears.....	4

The above spreader settings are approximate. (20 pound bag: Calibrate your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet) (25 pound bag: Calibrate your spreader on 200 square feet and adjust to apply at the rate of 1 pound of product per 200 square feet) (30 pound bag: Calibrate your spreader on 166 square feet and adjust to apply at the rate of 1 pound of product per 166 square feet)

GUARANTEED ANALYSIS 16-4-8	
Total Nitrogen (N).....	16%
Water Insoluble Nitrogen	5%
Available Phosphate (P ₂ O ₅).....	4%
Soluble Potash (K ₂ O).....	8%

- WEEDS -		
Bedstraw	Knotweed	Ragweed
Black Medic	Lambsquarters	Scarlet Pimpernel
Buckhorn	Lespedeza	Sheep Sorrel
Burdock	Mallow	Shepherdspurse
Chickweed	Morningglory	Speedwell
Chicory	Oxalis (yellow woodsorrel)	Spurge
Clover	Peppergrass	Thistle
Dandelion	Pigweed	Wild Carrot
Dock	Plantains	Wild Garlic
Ground Ivy	Poison Ivy	Wild Lettuce
Healall	Poison Oak	Wild Onion
Henbit	Purslane	Yarrow

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

APPENDIX

I. Previously Approved Fertilizer Grades:

Fertilizer grades with nutrient contents expressed as the percentages of N,P₂O₅, and K₂O include the following:

12-3-5	16-4-8	23-3-5
12-4-8	20-2-4	25-3-3
15-5-7	21-2-4	28-3-3
15-7-7	22-2-4	

The fertilizer grade guaranteed analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations. The following guaranteed analyses presentations have previously been submitted as notifications:

ii. Container Packaging and Size

Note to Supplemental Registrants: Select the appropriate bag size and present only the information provided for that bag size on your product labeling.

Bag Size, Pounds	Desired Application Rates		Treated Area for This Bag Size, Sq. Ft.
	Pounds/Area	Pounds/1000 Sq. Ft.	
20	1/250	4.0	5,000
	1/200	5.0	4,000
	1/166	6.0	3,333
25	1/250	4.0	6,250
	1/200	5.0	5,000
	1/166	6.0	4,167
30	1/250	4.0	7,500
	1/200	5.0	6,000
	1/166	6.0	5,000
40	1/250	4.0	10,000
	1/200	5.0	8,000
	1/166	6.0	6,667
50	1/250	4.0	12,500
	1/200	5.0	10,000
	1/166	6.0	8,333

iii. Advertising claims that may be presented on the retail containers:

- ◆ _____ pounds of product cover up to _____ square feet of turf.
- ◆ Controls over 30 weeds and builds a vigorous green lawn in one application.
- ◆ Provides lawns with a deep green color.
- ◆ Kills dandelion, spurge and many other broadleaf weeds.
- ◆ Controls dandelion, spurge, and many other broadleaf weeds.
- ◆ Kills the weeds and feeds the lawn in one easy application.
- ◆ Controls the weeds and feeds the lawn in one easy application.
- ◆ Also for other grass lawns including fescue and bermudagrass.
- ◆ Contains Trimec® Herbicide.
- ◆ Referrals to sequential applications of other pesticides:
 - ◇ Control existing broadleaf weeds with (supplemental registrant's brand name) containing Trimec® Herbicide during the growing season.
 - ◇ Stop broadleaf weeds seeds before they sprout with Bandini Pre Sprout 1, September through March/April.
 - ◇ Stop annual weed grass seeds before they sprout with Bandini Pre Sprout 2, August through March/April.