



**UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460**

**OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION**

June 2, 2022

Eric D. Smith
Director, Regulatory Affairs
PBI/Gordon Corporation
P.O. Box 860350
Shawnee, KS 66286

Subject: Notification per PRN 98-10 – Add optional spreader pattern statement and image and update the company address
Product Name: GORDON'S TRIMEC WEED & FEED 20
EPA Registration Number: 2217-580
Application Date: October 25, 2021
Decision Number: 583523

Dear Eric Smith:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped "NOTIFICATION" and placed in our records.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If you have any questions, please contact Endia Blunt at 202-566-2505 or at blunt.endia@epa.gov.

Sincerely,

A handwritten signature in cursive script that reads "Mindy Ondish".

Mindy Ondish
Product Manager 23
Herbicide Branch
Registration Division (7505T)
Office of Pesticide Programs

GORDON'S TRIMEC® WEED & FEED 20

EPA Reg. No. 2217-580

ACTIVE INGREDIENTS:

2,4-dichlorophenoxyacetic acid	0.46%
(+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid	0.10%
Dicamba: 3,6-dichloro-o-anisic acid	0.04%
OTHER INGREDIENTS:	99.40%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- 9.2 lb 2,4-dichlorophenoxyacetic acid equivalent per ton.
- 2.0 lb (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid equivalent per ton.
- 0.8 lb 3,6-dichloro-o-anisic acid equivalent per ton.

Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

NOTIFICATION

2217-580

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

06/02/2022

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

See attached booklet for complete [Precautionary Statements and] Directions for Use, including Storage and Disposal, and back panel for [complete Precautionary Statements, including] First Aid. OR See back panel for complete Precautionary Statements, including First Aid, and Directions for Use, including Storage and Disposal.

Net Contents: _____

EPA Est. No. _____

Company Name and Address:

PBI/Gordon Corporation

1217 West 12th Street P.O. Box 860350

Kansas City, Missouri 64104 Shawnee, KS 66286



PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves made of any waterproof material. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid	
If on skin or on clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 - 20 minutes.• Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferable mouth-to-mouth if possible.
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call toll-free, 1-877-800-5556 for emergency medical treatment information.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate. This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

[Environmental Hazards (alternate per PRN 2008-1)]

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow,

may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.]

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Do not use this product where desirable clovers are present or allow the material to drift onto flowers, vegetables or ornamental shrubbery. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning.
- Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.
- Do not apply to dichondra or carpetgrass.
- Do not apply this product to 'Floritam' St. Augustinegrass in Florida

INSTRUCTIONS:

Gordon's Trimec® Weed & Feed 20 controls a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others as listed, which are commonly found in home lawns. This product may be applied to established lawns of bermudagrass, bentgrass, bahiagrass, centipedegrass, Kentucky bluegrass, fescues, perennial ryegrass, zoysiagrass, and St. Augustinegrass. In addition, a X-X-X fertilizer provides a supply of plant nutrients or plant food.

Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in 30 days. [(Note: Specific to slow release fertilizers with high nitrogen content, such as 37-0-2) A retreatment may be made in 6 months.]

The maximum broadcast application rate is 6.0 pounds of product per 1,000 sq.ft. per application [1.2 lb 2,4-D ae, 0.27 lb MCP-P ae and 0.11 lb dicamba ae per acre per application]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. [(Note: Specific to slow release fertilizers with high nitrogen content, such as 37-0-2) The maximum number of broadcast applications is limited to 2 per year with a minimum of 6 months between applications.] The maximum seasonal rate is 12.0 pounds of product per 1,000 sq.ft. [2.4 lb 2,4-D ae, 0.54 lb MCP-P ae and 0.22 lb dicamba ae per acre].

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

WEEDS			
Annual yellow sweetclover	Clover, white	Mustard	Spiny cocklebur
Aster	Cockle	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Cocklebur	Narrowleaf vetch	Spotted catsear
Bedstraw	Common mullein	Nettle	Spotted spurge
Beggarticks	Creeping jenny	Orange hawkweed	Spurweed
Betony, Florida	Cudweed	Oriental cocklebur	Stinging nettle
Bindweed, field	Curly dock	Oxalis	Strawberry, India mock
Bird vetch	Daisy, English	Parsley-piert	Tall nettle
Bitter wintercress	Daisy fleabane	Parsnip	Tall vervain
Bittercress, hairy	Daisy, oxeye	Pearlwort	Tansy ragwort
Bitterweed	Dandelion	Pennycress	Tansy mustard
Black-eyed Susan	Dichondra	Pepperweed	Tanweed
Black medic	Dogfennel	Pigweed	Thistle
Black mustard	Dollarweed	Pineywoods bedstraw	Trailing crownvetch
Blackseed plantain	False dandelion	Plains coreopsis	Tumble mustard
Blessed thistle	Falseflax	Plantain	Tumble pigweed
Bloodflower milkweed	False sunflower	Poison ivy	Velvetleaf
Blue lettuce	Fiddleneck	Poison oak	Venice mallow
Blue vervain	Florida pusley	Pokeweed	Virginia buttonweed
Bracted plantain	Frenchweed	Poorjoe	Virginia creeper
Bristly oxtongue	Galinsoga	Prairie sunflower	Virginia pepperweed
Broadleaf dock	Goldenrod	Prickly lettuce	Wavyleaf bullthistle
Broadleaf plantain	Ground ivy	Prickly sida	Western clematis
Broomweed	Gumweed	Prostrate knotweed	Western salsify
Buckhorn	Hairy fleabane	Prostrate pigweed	White mustard
Buckhorn plantain	Hawkweed	Prostrate spurge	Wild aster
Bulbous buttercup	Healall	Prostrate vervain	Wild buckwheat
Bull thistle	Heartleaf drymary	Puncturevine	Wild carrot
Bullnettle	Heath aster	Purslane, common	Wild four-o'clock
Burclover	Hedge bindweed	Ragweed	Wild garlic
Burdock	Hedge mustard	Red sorrel	Wild geranium
Burning nettle	Henbit	Redroot pigweed	Wild lettuce
Burweed	Hoary cress	Redstem filaree	Wild marigold
Buttercup	Hoary plantain	Rough cinquefoil	Wild mustard
Buttonweed	Hoary vervain	Rough fleabane	Wild onion
Canada thistle	Horsenettle	Russian pigweed	Wild parsnip
Carolina geranium	Jimsonweed	Russian thistle	Wild radish
Carpetweed	Knawel	Scarlet pimpernel	Wild rape
Catchweed bedstraw	Knotweed	Scotch thistle	Wild strawberry
Catnip	Kochia	Sheep sorrel	Wild sweet potato
Catsear	Lambsquarters	Shepherdspurse	Wild vetch
Chickweed, common	Lespedeza	Slender plantain	Woodsorrel
Chickweed, mouseear	Mallow	Smallflower galinsoga	Woolly croton
Chicory	Matchweed	Smooth dock	Woolly morningglory
Cinquefoil	Mexicanweed	Smooth pigweed	Woolly plantain
Clover, crimson	Milk vetch	Sorrel	Wormseed
Clover, hop	Morningglory	Sowthistle	Yarrow
Clover, red	Mouseear hawkweed	Spanishneedles	Yellow rocket
Clover, strawberry	Mugwort	Speedwell	Yellowflower pepperweed
Clover, sweet	Musk thistle	Spiny amaranth	

[Begin Optional Language

(Note to Supplemental Registrants: Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, equipment settings of drop spreaders, rotary spreaders, and other granular applicators may be presented on the container labels.)

SPREADERS AND CALIBRATION:

1. Drop spreaders and rotary spreaders must be calibrated for accurate applications of this product.
2. Close spreader opening (orifice).

3. Adjust spreader settings using the spreader settings in the following chart.

Spreader Settings	
Scott	6
Lawn Beauty	5
Jackson	7.5
Sears	4

The above spreader settings are approximate.

20 pound bag: Calibrate your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet. Application rate: 4.0 pounds of product/ 1,000 sq.ft.

25 pound bag: Calibrate your spreader on 200 square feet and adjust to apply at the rate of 1 pound of product per 200 square feet. Application rate: 5.0 pounds of product/ 1,000 sq.ft.

30 pound bag: Calibrate your spreader on 166 square feet and adjust to apply at the rate of 1 pound of product per 166 square feet. Application rate: 6.0 pounds of product/ 1,000 sq.ft.

Bag Size, Pounds	Desired Application Rates		Treated Area for This Bag Size, sq.ft.
	Pounds/Area (sq.ft.)	Pounds/1000 sq.ft.	
20	1/250	4.0	5,000
	1/200	5.0	4,000
	1/166	6.0	3,333
25	1/250	4.0	6,250
	1/200	5.0	5,000
	1/166	6.0	4,167
30	1/250	4.0	7,500
	1/200	5.0	6,000
	1/166	6.0	5,000
40	1/250	4.0	10,000
	1/200	5.0	8,000
	1/166	6.0	6,667
50	1/250	4.0	12,500
	1/200	5.0	10,000
	1/166	6.0	8,333

End Optional Language]

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

APPENDIX

1. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

Herbicide Benefit Claims:

- Controls (Kills) over (Number) weeds and builds a vigorous green lawn in one application.
- Kills (Controls) dandelion, spurge and many other broadleaf weeds as listed.
- Controls Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and many other broadleaf weeds including those listed on the back panel.
- Kills (Controls) the weeds and feeds the lawn in one easy application.
- Banish weeds and enjoy a greener lawn that looks as good as it feels.
- Fertilizes & controls broadleaf weeds in one easy step.
- Controls (Kills) most major broadleaf weeds.
- Kills (Controls) many broadleaf weeds
- Controls (Kills) (Number) weeds! (Listing of weeds may include the following.)
- (Listing of weeds may include the following.)
- Kills dandelions & feeds grass
- Weed & Feed
- [For use with sublabel 2 only] Season long fertilizer with dandelion control
- Dandelion control
- Broadleaf control

Fertilizer Benefit Claims:

- Provides lawns with a deep green color.
- Also for other grass lawns including fescue and bermudagrass.
- Longer lasting fertilizer formula.
- Plus a quick greening lawn fertilizer.
- Promotes thick, green lawns.
- Also for other grass lawns
- Provides deep green color [and thick green lawns]
- A (X-X-X) fertilizer provides a balanced supply of plant food needed for a normal season's growth.
- [Alternate Spreader Listing and Graphics]: Bandini - XX; Vigoro 4000 Rotary - XX; Scotts Drop - XX; Vigoro 4000 Drop - XX
- Builds healthy green grass
- (Note: Specific to slow release fertilizers with high nitrogen content, such as 37-0-2) When applied as directed, this product will apply less than 2.5 lb of nitrogen per 1,000 sq.ft. and will release less than 0.7 lb of nitrogen per month.
- (Note: Specific to slow release fertilizers with high nitrogen content, such as 37-0-2) Feeds [your lawn] for up to 6 months
- (Note: Specific to slow release fertilizers with high nitrogen content, such as 37-0-2) One application [feeds] [fertilizes] for 6 months
- (Note: Specific to slow release fertilizers with high nitrogen content, such as 37-0-2) Feeds all season
- (Note: Specific to slow release fertilizers with high nitrogen content, such as 37-0-2) Fertilizer lasts [up to 5 months] [all season]
- Non-burning fertilizer
- 80% [coated] slow release nitrogen
- (Note: Specific to slow release fertilizers with high nitrogen content, such as 37-0-2) Uses less nitrogen than standard 4 application program and keep lawn green for 6 months
- Less surge growth throughout season
- More green – less clippings
- Reduced surge growth of clippings
- Same money – Save time
- Mow less frequently

- (Note: Specific to slow release fertilizers with high nitrogen content, such as 37-0-2) Polyon coating technology releases fertilizer slowly when the lawn needs it
- (Note: Specific to slow release fertilizers with high nitrogen content, such as 37-0-2) Fertilizer is done in one application
- (Note: Specific to slow release fertilizers with high nitrogen content, such as 37-0-2) Why spread fertilizer 4 times a season when you can do it in one application


General Marketing Claims:

- **SATISFACTION GUARANTEED OR YOUR MONEY BACK.** If you apply (Brand Name) as directed and are not completely satisfied, send proof of purchase from this package and a cash register receipt to the address on the front of the bag. We will promptly refund your money.
- **Guarantee of Satisfaction:** If this (Company Name) product is not performing to your satisfaction, please send proof of purchase and a brief description of your problem to (Company Name and address), for a prompt response. Also enclose a daytime telephone number.
 - _____ pounds of product cover up to _____ square feet of turf.
 - Contains Trimec® Herbicide.
 - Covers [up to] _____ sq.ft
 - One pass
 - Single [app] [pass]

Referrals to other products used sequentially:

- Control existing broadleaf weeds with (Brand Name) containing Trimec® Herbicide during the growing season.
- Control existing broadleaf weeds with Bandini Super Green Weed & Feed, Bandini St. Augustine Weed & Feed, & Bandini Super Plush Weed & Feed during the growing season.
- Stop broadleaf weeds seeds before they sprout with (Brand Name), September through March/April.
- Stop broadleaf weeds seeds before they sprout with Bandini Pre-Sprout 1, applications made September through April.
- Stop annual weed grass seeds before they sprout with (Brand Name), August through March/April.
- Stop annual weed grass seeds before they sprout with Bandini Pre-Sprout 2 applications made August through April.

Other optional spreader language/statements and related images:



- **Spreader Pattern: Broadcast Spreader - First spread the borders of your lawn. Walk at an even speed making sure that particles [granules] are overlapping the previous pass. Shut off spreader when stopped or when you reach a border strip.**

2. State mandated fertilizer labeling requirements:

[Note to Supplemental Registrants: Please select the statements relevant to the areas of the country in which your specific product is registered and sold; if any of these statements are relevant to your labeling, please place in the Directions for Use under the heading “3.1 State Restrictions”]

GUARANTEED ANALYSIS XX-X-X	
Total Nitrogen (N)	XX%
x.x% ammoniacal nitrogen	
x.x% urea nitrogen*	

Available Phosphate (P ₂ O ₅)	X%
Soluble Potash (K ₂ O)	X%
Sulfur (S)	X%
x.x% free sulfur (S)	
Derived from: polymer-coated sulfur-coated urea, urea, ammonium phosphate, potassium chloride. *A portion of the urea has been coated to provide x.x% coated slow-release nitrogen. This product and/or its manufacture is covered by one or more of the following U.S. patents and international counterparts thereof: US Patent Numbers: xxxxxx	

[Note: Fertilizer grades with nutrient contents expressed as the percentages of N, P₂O₅ and K₂O may include the following: 23-3-7 or x-x-x. The fertilizer grade analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.]

- California, Oregon, Washington: “[When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials.]Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>”
- Florida: [Note to reviewer: maximum number of applications in this section is dually based upon maximum ai load as well as fertilizer analysis effect on maximum nitrogen rates in different regions in Florida]:
 - In North Florida: Do not exceed X applications per year on bermudagrass. Do not exceed X applications per year on centipedegrass. Do not exceed X applications per year on St. Augustine grass. Do not exceed X applications per year on zoysiagrass.
 - In Central Florida: Do not exceed X applications per year on bermudagrass. Do not exceed X applications per year on centipedegrass. Do not exceed X applications per year on St. Augustine grass. Do not exceed X applications per year on zoysiagrass.
 - In South Florida: Do not exceed X applications per year on bermudagrass. Do not exceed X applications per year on centipedegrass. Do not exceed X applications per year on St. Augustine grass. Do not exceed X applications per year on zoysiagrass.
- Florida: “Apply only to actively growing turf.”
- Florida: “Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices. Check with your county or city government to determine if there are local regulations for fertilizer use.
- Maryland: “This fertilizer contains phosphorus and may not be used on turf in the state of Maryland except when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory identified under § 8-803.7 of the Agriculture Article, Annotated Code of Maryland, performed no more than 3 years before the application; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State; or 3) Reestablishing or repairing a turf area.”
- New Jersey: “In the state of New Jersey, this product may be used only when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary pursuant to a soil test conducted by a laboratory identified pursuant to subsection b. of section 6 of P.L.2010, c.112 (C.58:10A-66) and performed no more than three years before the application, and pursuant to the associated annual fertilizer recommendation issued by the New Jersey Agricultural Experiment Station at Rutgers, the State University; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is in accordance with the standards and requirements established under the “Soil Erosion and Sediment Control Act,” P.L.1975, c.251 (C.4:24-39 et seq.) and the rules and regulations adopted pursuant thereto; 3) Reestablishing or repairing a turf area; or 4) Delivering liquid or granular fertilizer under the soil surface, directly to the feeder roots.”
- New Hampshire: In the state of New Hampshire, this product may only be used on 1) Establishing new lawns, for repairing a lawn, for seeding, or for use when a soil test indicates a phosphorus deficiency.

- Virginia: "In the state of Virginia, this product may be used only on 1) Newly established turf areas from sod or seed during the first growing season; 2) Turf areas being repaired or renovated, and; 3) Turf areas where soil tests performed within the past three years indicate nutrient deficiency."
- [Virginia, Florida, Maryland, others] DO NOT APPLY NEAR WATER, STORM DRAINS, OR DRAINAGE DITCHES. DO NOT APPLY IF HEAVY RAIN IS EXPECTED. APPLY THIS PRODUCT ONLY TO YOUR LAWN/GARDEN. [AND SWEEP ANY PRODUCT THAT LANDS ON THE DRIVEWAY, SIDEWALK, OR STREET, BACK INTO YOUR LAWN/GARDEN.] [Note to Supplemental Registrant: The bracketed part of this statement is not required for liquid products.]
- [For products containing Iron:] Keep this product off sidewalks, patios, driveways, etc. This fertilizer contains iron which will stain concrete and painted surfaces. If it comes in contact with concrete surface, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.
- [Note to Supplemental Distributor: In Kentucky if the product supplies less than 1lb of Nitrogen per application, include the following:] A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing XX pounds of nitrogen per 1,000 square feet and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program."

DOCUMENT CONTROL INFORMATION

1. Unique Label Identifier: 002217-00580.20211025.notif-proposed-highlighted.doc

2. Reason for Issue: Spreader pattern image and optional spreader language

Internal Usage Notes
Sublabel 1. For use with fertilizers which can be applied on a 30-day interval Associated Container Label: Supplemental Distributors
Sublabel 2. For use with fertilizers which require 6 months between applications Associated Container Label: Supplemental Distributors