

Gordon's

*TRIMEC

Weed & Feed 24

18-4-8

Contains *TRIMEC Herbicide

Twenty-five Pounds Covers up to 5,000 Sq. Ft.

KILLS: Dandelion, Chickweed, Knotweed, Plantains, Henbit, Spurge and many other broadleaf weeds; some of which are listed on the back panel

ACTIVE INGREDIENTS

- **2,4-Dichlorophenoxyacetic acid
- **2,4-Dimethyl-4-chlorophenoxyacetic acid
- **2,4-Dimethyl-6-chlorophenoxyacetic acid

INERT INGREDIENTS

THIS PRODUCT CONTAINS

- **10.940 lbs. 2,4-Dichlorophenoxyacetic acid
- **4.897 lbs. 2,4-Dimethyl-4-chlorophenoxyacetic acid
- **1.033 lbs. 2,4-Dimethyl-6-chlorophenoxyacetic acid

*Isomer Specific by AOCAC Method 111-11

KEEP OUT OF REACH OF CHILDREN CAUTION


STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of water for 15 minutes and get medical attention.

See back panel for additional information.

*TRIMEC® is a registered trademark of PBI GORDON CORPORATION, U.S. Patent No. 3,284,186.

NET WEIGHT 25 POUNDS



Step 1. Read the entire label first.
Observe all cautions and follow directions carefully.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

Caution: May be harmful if swallowed. Causes eye irritation. Avoid contact with skin and clothing. Wash thoroughly after handling.

Environmental Hazards

DIRECTIONS FOR USE

Apply to lawns and areas where weeds are present or growing. Do not apply to areas where desirable plants are growing.

STORAGE & DISPOSAL

USE PRECAUTIONS

1. Do not apply to areas where desirable plants are present or growing. Do not apply to areas where desirable plants are growing. Do not apply to areas where desirable plants are growing.

INSTRUCTIONS

1. Apply to lawns and areas where weeds are present or growing. Do not apply to areas where desirable plants are growing.
2. Water lawns thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to add moisture and prevent drift onto non-target plants.
4. Do not water lawns for 1 to 2 days after application. At this time a light watering should be made.

Be patient. TRIMEC works slowly and gently and is friendly to your lawn.

SPREADER SETTINGS

Scott	6
Lawn Beauty	5
Jackson	7 1/2
Scot	4

The above spreader settings are approximate. Check your spreader on 250 sq. ft. and adjust to give the rate of 1 lb. per 250 sq. ft.

FERTILIZER GUARANTEE

Total Nitrogen	5%
Water-soluble Nitrogen	4%
Available Phosphoric Acid	8%
Soluble Potash	18%

WEEDS

BEDSTRAW	DANDELION	LESPEDEZA	POISON IVY	SPURGE
BLACK MEDIC	DOCK	MALLOW	POISON OAK	THISTLE
BUCKMORN	GROUND IVY	MORNING GLORY	PURSLANE	WILD GARLIC
BURDOCK	HEAL-ALL	OXALIS	RAGWEED	WILD LETTUCE
CHICORY	HENBIT	PEPPERGRASS	SHEEP SORREL	WILD ONION
CHICKWEED	KNOTWEED	PIGWEEED	SHEPHERD'S PURSE	YARROW
CLOVER	LAMBSQUARTER	PLANTAINS	SPEEDWELL	

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

EPA Reg. No. 227-501

EPA Est. No. 1-KS-1

574-881

