2217-585 7/1/2003.

Page 137

GORDON'S BROADLEAF AND POISON IVY SPOT-WEEDER

READY TO USE

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	0.311%
Mecoprop, dimethylamine salt	0.075%
Dicamba, dimethylamine salt	0.034%
INERT INGREDIENTS:	99.580%
TOTAL	100.000%

THIS PRODUCT CONTAINS:

0.258% 2,4-dichlorophenoxyacetic acid equivalent 0.062% (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent 0.028% 3,6-dichloro-o-anisic acid equivalent Isomer Specific by AOAC Methods.

CONTAINS SINGLE ISOMER FORM OF MECOPROP-D

CAS Registry Numbers:

2,4-D, dimethylamine salt CAS# 2008-39-1 Mecoprop, dimethylamine salt CAS# 66423-09-4 Dicamba, dimethylamine salt CAS#2300-66-5

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for Precautionary Statements and First Aid.


NET CONTENTS: 24 Fluid Ounces, 32 Fluid Ounces (1 Quart), 64 Fluid Ounces (2 Quarts), 128 Fluid Ounces (1 Gallon)

864 AP----EPA REG. NO. 2217-585
EPA EST. NO. 2217-KS-1
MANUFACTURED BY

Bill Conclor

An Employee-Owned Company
1217 West 1218 Street

Telephone: 1-800-821-7925





READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Avoid contact with skin, eyes or clothing. Wear goggles when applying this product. When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves.

After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

If in eyes:	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing:	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.
If swallowed	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
	ct container or label with you when calling a poison control center or doctor or going for may also contact 1-877-800-5556 for emergency medical treatment advice.

ENVIRONMENTAL HAZARDS:

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater.

SUBLABEL I. DIRECTIONS FOR USE PREVIOUSLY APPROVED ON **OCTOBER 22, 1998.**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

STORAGE AND DISPOSAL

STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse container. Rinse thoroughly before discarding in trash.

USE PRECAUTIONS:

Avoid spraying susceptible plants (flowers, shrubs, etc.). Clovers, bentgrass and St. Augustinegrass may be injured if contacted. Do not apply within dripline of desirable plants. Avoid accidental drift or application to nearby garden plants, flowers or shrubs. Do not use on dichondra lawns.

NO MIXING OR POURING - AIM AT WEED AND SPRAY

Use to kill poison ivy and any other troublesome broadleaf weeds. For best results use Gordon's Broadleaf and Poison Ivy Spot Weeder when weeds are actively growing. Rainfall or watering soon after application may reduce effectiveness.

Gordon's Broadleaf and Poison Ivy Spot Weeder is designed to permit easy treatment of small weed patches without mess or trouble and without bending or stooping. Wet poison ivy or other weeds thoroughly using short bursts of spray. Overspraying only wastes the spray. Effects begin to show in a few days and weeds slowly die and disappear.

ALSO KILLS:

Bedstraw	Filaree	Pennywort
Black medic (Yellow clover)	Fleabane	Pepperweed
Buckhorn	Florida pusley	Pigweed
Canada thistle	Ground ivy	Plantain
Chickweed	(Creeping Charlie)	Poison ivy
Clover	Healali	Prickly lettuce
Cudweed	Henbit	Purslane
Curled dock	Knotweed	Ragweed
Dandelion	Lambsquarters	Sheep sorrel
Dollarweed	Matchweed	Shepherdspurse
English daisy	Mustard	Yarrow

SUBLABEL II. PROPOSED DIRECTIONS FOR USE.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried.

STORAGE AND DISPOSAL

STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children and domestic animals.

CONTAINER DISPOSAL:

If Empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partially filled: Call your local solid waste agency or toll free number (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRODUCT DESCRIPTION:

Gordon's Broadleaf and Poison Ivy Spot Weeder contains TRIMEC[®] turf herbicide, a combination of three proven weed killers - 2,4-D, mecoprop, and dicamba. Together they control a wide range of broadleaf weeds commonly found in home (residential) lawns. This ready-to-use herbicide is intended for spot treatments only. Weeds that have not emerged at the time of application will not be controlled.

WHERE TO USE:

Northern Grasses - Kentucky bluegrass, perennial ryegrass and fescues.

Southern Grasses - St. Augustinegrass, centipedegrass, bentgrass, bahiagrass, bermudagrass and zoysiagrass.

477

PROHIBITIONS:

- Do not apply to garden plants, flowers or shrubs.
- Do not use this product on carpetgrass, dichondra, nor on lawns with desirable clovers.

HOW TO USE:

Use to spot treat weeds in the lawn. Gordon's Broadleaf and Poison Ivy Spot Weeder is designed for easy treatment of small weed patches without mess or trouble and without bending or stooping.

NO MIXING OR POURING - AIM AT WEED AND SPRAY

For best results, use Gordon's Broadleaf and Poison Ivy Spot Weeder when weeds are actively growing. Rainfall or watering soon after application may reduce effectiveness.

FOR CONTAINERS WITH TRIGGER SPRAYERS:

Gordon's Broadleaf and Poison Ivy Spot Weeder is ready-to-use and is applied directly from the container fitted with the trigger sprayer. Direct the spray to the target weeds to provide uniform coverage of the upper leaf surfaces. [OR: Wet the broadleaf weeds thoroughly using short bursts of spray.] Usually, two (2) pumps of this trigger sprayer will provide adequate coverage of individual weeds. Excessive spraying can injure some lawns. [OR: Overspraying may injure lawns including bentgrass, centipedegrass, and St. Augustinegrass.]

FOR REFILL CONTAINERS:

Gordon's Broadleaf and Poison Ivy Spot Weeder is ready-to-use and is applied directly from the container. Remove the cap from the container and attach the trigger sprayer. Direct the spray to the target weeds to provide uniform coverage of the upper leaf surfaces. [OR: Wet the broadleaf weeds thoroughly using short bursts of spray.] Excessive spraying can injure some lawns. [OR: Overspraying may injure lawns including bentgrass, centipedegrass, and St. Augustinegrass.]

CARE AND INSTRUCTIONS FOR TRIGGER SPRAYERS

OPERATION:

- 1. Remove sprayer from bag.
- 2. Unscrew cap from bottle and remove the foil seal.
- 3. Making sure that tube is properly inserted through hole in cap, insert tube into bottle and screw on cap.
- 4. Open spray nozzle approximately one full turn.
- 5. TO PRIME With sprayer directed at the target area, operate trigger until spray begins (approximately 10 to 15 times).
- 6. Adjust the sprayer nozzle for desired spray pattern.

STORAGE OF TRIGGER SPRAYER:

- 1. Close spray nozzle.
- 2. Hook the trigger on or under the bottle handle so that trigger is above the level of contents.
- To prevent clogging of trigger, remove sprayer from bottle and pump clean water through the hose and trigger.

HOW TO PRIME AND CARE FOR YOUR SPRAYER:

- 1. After shaking product well, remove cap from container and replace with sprayer hose and cap assembly. Tighten securely.
- 2. Turn sprayer nozzle one half turn counter-clockwise. Hold sprayer below the level of container and prime by pumping trigger 10 15 times until product is dispensed. Adjust nozzle to coarse stream.
- 3. To store, turn nozzle to "OFF" position and store with sprayer above level of liquid in container to prevent leakage

Use Gordon's Broadleaf and Poison Ivy Spot Weeder to kill poison ivy and any other troublesome broadleaf weeds.

- WEEDS -					
Annual yellow sweetclover	Cockle	Musk thistle	Spiny cocklebur		
Aster	Cocklebur	Mustard	Spiny sowthistle		
Austrian fieldcress	Common mullein	Narrowleaf plantain	Spotted catsear		
Bedstraw	Creeping jenny	Narrowleaf vetch	Spotted spurge		
Beggarticks	Cudweed	Nettle	Spurweed		
Betony, Florida	Curly dock	Orange hawkweed	Stinging nettle		
Bindweed, field	Daisy, English	Oriental cocklebur	Strawberry, India mock		
Bird vetch	Daisy fleabane	Oxalis	Tall nettle		
Bitter wintercress	Daisy, oxeye	Parsley-piert	Tall vervain		
Bittercress, hairy	Dandelion	Parsnip	Tansy ragwort		
Bitterweed	Dichondra	Pearlwort	Tansy mustard		
Black-eyed Susan	Dogbane	Pennycress	Tanweed		
Black medic	Dogfennel	Pennywort	Thistle		
Black mustard	Dollarweed	Pepperweed	Trailing crownvetch		
Blackseed plantain	Elderberry	Pigweed	Tumble mustard		
Blessed thistle	False dandelion	Pineywoods bedstraw	Tumble pigweed		
Bloodflower milkweed	Falseflax	Plains coreopsis (tickseed)	Velvetleaf		
Blue lettuce	False sunflower	Plantain	Venice mallow		
Blue vervain	Fiddleneck	Poison ivy	Virginia buttonweed		
Bracted plantain	Florida pusley	Poison oak	Virginia creeper		
Brassbuttons	Frenchweed	Pokeweed	Virginia pepperweed		
Bristly oxtongue	Galinsoga	Poorjoe	Water pennywort		
Broadleaf dock	Goathead	Prairie sunflower	Wavyleaf bullthistle		
Broadleaf plantain	Goldenrod	Prickly lettuce	Western clematis		
Broomweed	Ground ivy	Prickly sida	Western salsify		
Buckhorn	Gumweed	Prostrate knotweed	White mustard		
Buckhorn plantain	Hairy fleabane	Prostrate pigweed	Wild aster		
Bulbous buttercup	Hawkweed	Prostrate spurge	Wild buckwheat		
Bull thistle	Healall	Prostrate vervain	Wild carrot		
Bullnettle	Heartleaf drymary	Puncturevine	Wild four-o'clock		
Burclover	Heath aster	Purslane, common	Wild garlic		
Burdock	Hedge bindweed	Ragweed	Wild geranium		
Burning nettle	Hedge mustard	Red sorrel	Wild lettuce		
Burweed	Hemp	Redroot pigweed	Wild marigold		
Buttercup	Henbit	Redstem filaree	Wild mustard		
Buttonweed	Hoary cress	Rough cinquefoil	Wild onion		
Canada thistle	Hoary plantain	Rough fleabane	Wild parsnip		
Carolina geranium	Hoary vervain	Russian pigweed	Wild radish		
Carpetweed	Horsenettle	Russian thistle	Wild rape		
Catchweed bedstraw	Jimsonweed	Scarlet pimpernel	Wild strawberry		
Catnip	Knawel	Scotch thistle	Wild sweet potato		
Catsear	Knotweed	Sheep sorrel	Wild vetch		
Chickweed, common	Kochia	Shepherdspurse	Woodsorrel		
Chickweed, mouseear	Lambsquarters	Slender plantain	Woolly croton		
Chicory	Lespedeza	Smallflower galinsoga	Woolly morningglory		
Cinquefoil	Mallow	Smooth dock	Woolly plantain		
Clover, crimson	Matchweed	Smooth pigweed	Wormseed		
Clover, hop	Mexicanweed	Sorrel	Yarrow		
Clover, red	Milk vetch	Sowthistle	Yellow rocket		
Clover, strawberry	Morningglory	Spanishneedles	Yellowflower pepperweed		
Clover, sweet	Mouseear hawkweed	Speedwell	and other		
Clover, white	Mugwort	Spiny amaranth	broadleaf weeds		

6 2 7

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

フォフ

APPENDIX

- I. Advertising claims that may be used by supplemental registrants and be presented on the container label.
 - No mixing No pouring
 - · Ready-To-Use Aim at weed and spray.
 - ♦ Contains TRIMEC®
 - ◆ TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.
 - Kills (Controls) (Number) weed types, roots and all
 - Water-based or Water-based formulation
 - Kills the weeds, not the lawn
 - Kills the root.
 - Kills the entire weed or plant roots and all.
 - Kills (Controls) a wide range of broadleaf weeds roots and all.
 - ◆ Controls (Kills) (desired weed species from attached listing may be inserted here,) and many other broadleaf lawn weeds.
 - ◆ Kills (Controls) (desired weed species from attached listing may be inserted here,) and over (Number) broadleaf weeds
 - ♦ Kills (Number) weeds
 - Kills weed roots