

DIRECTIONS FOR LIVESTOCK PEST CONTROL

Mix 2 quarts of Dairy and Livestock Spray per 6 gallons of water. Spray to thoroughly cover all parts of the animal, including the legs, using 1-2 pints of spray per large animal and proportionately less for smaller animals.

For pen or corral operations involving spray application to livestock with high pressure equipment, do not exceed a 0.5% concentration. Mix 2 quarts of Dairy and Livestock Spray per 12 gallons of water (0.5% concentration). Spray to thoroughly cover the animal's body and legs (avoid excessive spraying of the head) using 1-2 quarts of spray per large animal and proportionately less for smaller animals. Repeat above application as necessary to maintain control, but not more often than once every 7 days.

Improved face fly control may be achieved by adding sugar to the spray solution at the rate of 1 pound per 6-10 gallons of spray. Repeat above application as necessary to maintain control, but not more often than once every 7 days.

LOW VOLUME APPLICATION FOR CONTROL OF FACE FLIES

Mix 4 tablespoons (2 fluid oz.) of Dairy and Livestock Spray per 1 1/2 pints of water (1% concentration). Using a calibrated sprayer, apply 5 teaspoons of spray to the face and a total of 5-10 teaspoons to the back and sides of the animal 3 times per week. Improved control may be achieved by the addition of 1 level teaspoon of sugar per pint of spray solution.

FOR CONTROL OF LICE AND WINTER TICKS ON DAIRY CATTLE, BEEF CATTLE, SHEEP, GOATS AND HOGS


Mix 2 1/2 pints of Dairy and Livestock Spray per 16 gallons of water (0.25% concentration). Spray animals thoroughly using up to 1 gallon of spray. Apply a second application 14 days later. Repeat applications as necessary to maintain control, but not more often than once every 7 days.

BACKRUBBER APPLICATIONS

For use in backrubbers or face rubbers for the control of horn flies and face flies, mix 1 gallon of Dairy and Livestock Spray per 16 gallons of back rubber base oil (1% concentration). This may be diesel fuel or fuel oil in accordance with local recommendations.



DAIRY and LIVESTOCK SPRAY



FOR CONTROL OF HOUSE FLIES, FACE FLIES AND STABLE FLIES IN AND AROUND LIVESTOCK BUILDINGS

Mix 2 quarts of Dairy and Livestock Spray per 6 gallons of water (1% concentration). Spray premise such as inside and outside barn walls (including dairy barns) eaves, fences and other resting or breeding places at least twice monthly using approximately 1 gallon of finished spray per 1000 square feet (40' x 25'). Improved house fly and face fly control may be achieved by adding sugar to the spray solution at the rate of 1 pound per 6-10 gallons of spray.

PRECAUTIONS IN USING

KEEP OUT OF THE REACH OF CHILDREN AND DOMESTIC ANIMALS

WARNING! HAZARDOUS IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN. Wash thoroughly with soap and water after handling and before eating and smoking. Wear clean clothing. In case of spill age on person or clothing, immediately remove clothing and flush skin or eyes with plenty of water; for eyes, get medical attention. Keep away from heat and open flame. Atropine is antidotal. During commercial or exposure in spray-mixing and loading operations, wear clean rubber gloves and a mask or respirator of a type found suitable by the U.S. Bureau of Mines for protection against Dairy and Livestock Spray. Use this material only for recommended purposes and at recommended dosages.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME Do not contaminate feed or foodstuffs. Do not contaminate streams, lakes or ponds with this material.


This product is toxic to fish and wildlife. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not use in poultry houses.

Do not treat animals within 1 day of slaughter.

DECONTAMINATE AND DESTROY USED CONTAINERS. Do not reuse for any purpose. Decontaminate containers by rinsing thoroughly with water and alkaline detergent. Destroy containers by breaking and burying fragments in isolated area. Dispose of rinsings in a way as not to constitute a hazard, or contaminate water supplies.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.



DIRECTIONS FOR LIVESTOCK PEST CONTROL

Mix 2 quarts of Dairy and Livestock Spray per 6 gallons of water. Spray liberally over all parts of the animal, including the legs, using a fine mist of spray per large animal and proportionately less for small animals.

For pest or control operations involving spray application to livestock with low pressure equipment, use maximum 0.5% concentration. Mix 2 quarts of Dairy and Livestock Spray per 12 gallons of water. 0.5% concentration. Spray to thoroughly cover the animal's body and head and extend spraying at the head using 1-2 quarts of spray per animal depending on the size of the animal. Repeat above concentration every 7 days to maintain control, but not more often than once a week.

For control of fly pests, spray liberally over the animal's body and head and extend spraying at the head using 1-2 quarts of spray per animal depending on the size of the animal. Repeat above concentration every 7 days to maintain control, but not more often than once a week.

LOW VOLUME APPLICATION FOR CONTROL OF FACE FLIES

Mix 4 quarts of Dairy and Livestock Spray per 6 gallons of water. Spray liberally over all parts of the animal, including the legs, using a fine mist of spray per large animal and proportionately less for small animals. Repeat above concentration every 7 days to maintain control, but not more often than once a week.

FOR CONTROL OF LICE AND WINTER TICKS ON

DAIRY CATTLE, BEEF CATTLE, SHEEP, GOATS AND HOGS

Mix 2 quarts of Dairy and Livestock Spray per 6 gallons of water. Spray liberally over all parts of the animal, including the legs, using a fine mist of spray per large animal and proportionately less for small animals. Repeat above concentration every 7 days to maintain control, but not more often than once a week.

BACK RUBBER APPLICATIONS

Mix 2 quarts of Dairy and Livestock Spray per 6 gallons of water. Spray liberally over all parts of the animal, including the legs, using a fine mist of spray per large animal and proportionately less for small animals. Repeat above concentration every 7 days to maintain control, but not more often than once a week.

DAIRY and LIVESTOCK SPRAY

For Control of Horn Flies, Stable Flies, Face Flies, House Flies, Lice and Lone Star Ticks on Dairy Cattle, Beef Cattle, Sheep, Goats and Hogs

Contains CIODRIN® and VAPONA® Insecticides

ACTIVE INGREDIENTS	BY WEIGHT
Dimethyl phosphate of alpha-methylbenzyl 3-hydroxy-cis-crotonate	10.00%
2,2-dichlorovinyl dimethyl phosphate	2.3%*
Related Compounds	0.2%*
Heavy Aromatic Naphtha	77.0%
INERT INGREDIENTS	10.5%
TOTAL	100.0%

WARNING! Keep out of Reach of Children
See Side Panels for Additional Warnings

*Equivalent 2.5% VAPONA® Insecticide
NET CONTENTS 1 GALLON
E.P.A. Reg. No. 2217-592

963-572

FOR CONTROL OF HOUSE FLIES, FACE FLIES AND STABLE FLIES IN AND AROUND LIVESTOCK BUILDINGS

Mix 2 quarts of Dairy and Livestock Spray per 6 gallons of water. Spray liberally over all parts of the animal, including the legs, using a fine mist of spray per large animal and proportionately less for small animals. Repeat above concentration every 7 days to maintain control, but not more often than once a week.

PRECAUTIONS IN USING

KEEP OUT OF THE REACH OF CHILDREN AND DOMESTIC ANIMALS

WARNING! HAZARDOUS IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN. Avoid contact with eyes, nose, mouth, and skin. If contact occurs, flush with water. If swallowed, do not induce vomiting. If inhaled, move to fresh air. If on skin, wash with soap and water. Do not use in confined spaces. Do not use near food or feed. Do not use on animals that are pregnant or nursing. Do not use on animals that are sick or stressed. Do not use on animals that are under the influence of other drugs. Do not use on animals that are under the influence of alcohol. Do not use on animals that are under the influence of any other substance. Do not use on animals that are under the influence of any other substance.

Do not use in confined spaces. Do not use near food or feed. Do not use on animals that are pregnant or nursing. Do not use on animals that are sick or stressed. Do not use on animals that are under the influence of other drugs. Do not use on animals that are under the influence of alcohol. Do not use on animals that are under the influence of any other substance.

DECONTAMINATE AND DESTROY USED CONTAINERS
Do not reuse containers. Do not use containers for food or feed. Do not use containers for any other purpose. Do not use containers for any other purpose.