

Use herbicides safely
Do not overdose
Read DIRECTIONS FOR USE
carefully

NET WT 35 Lbs.
TREATS
Up To 5000 Sq Ft.



ST. AUGUSTINE GRASS LAWN WEEDER & FEEDER

For Use On St. Augustine Grass
And Other Grasses For
Control Of Growing Broadleaf Weeds

CAUTION: Keep Out Of Reach Of Children.
See Back Panel for Additional Cautions

WEEDS AS IT FEEDS

ACTIVE INGREDIENTS	PERCENTAGE
**2,4-dichlorophenoxyacetic acid	0.800%
**2-(2-methoxy-4-chlorophenoxy)propionic acid	1.950%
***Dicamba (3,6-dichlorobenzoic acid)	0.0295%
INERT INGREDIENTS	99.5933%
TOTAL	100.0000%

Used at 35 lbs. per acre on St. Augustine grass
**2,4-D ACID 0.25 lbs. per acre
**MCPA ACID 0.90 lbs. per acre
***Dicamba ACID 0.19 lbs. per acre

U.S. Patent No. 3,244,186

MANUFACTURED BY
GORDON CORPORATION
KANSAS CITY, KANSAS

NET WEIGHT 35 POUNDS COVERS UP TO 5000 SQUARE FEET

GORDON'S

ST. AUGUSTINE GRASS LAWN WEEDER & FEEDER

For Control of Growing Broadleaf Weeds In
St. Augustine and Other Grasses

14-49
GUARANTEED ANALYSIS

TOTAL NITROGEN (not less than)	16.00%
NITRATE NITROGEN (not less than)	None
AMMONIACAL NITROGEN (not less than)	0.66%
WATER SOLUBLE ORGANIC NITROGEN (not less than)	5.67%
WATER INSOLUBLE NITROGEN (not less than)	4.67%
AVAILABLE PHOSPHORIC ACID (not less than)	4.00%
INSOLUBLE PHOSPHORIC ACID (not less than)	0.50%
WATER SOLUBLE POTASH (not less than)	8.00%
TOTAL AVAILABLE PRIMARY PLANT FOOD (not less than)	28.00%
CHLORINE (not more than)	1.00%

DIRECTIONS FOR USE

GORDON'S ST. AUGUSTINE GRASS LAWN WEEDER AND FEEDER destroys growing broadleaf weeds in St. Augustine, Bermuda, Centipede, Zoysia, and other turf grasses.

CONTROLS THESE WEEDS: Chickweed, Chickweed, Clover, Cockspur, Dandelion, Dock, Plantain, Ground Squirrel, Mustard, Oxalis, Peppergrass, Pitted Purslane, Poison Ivy, Purslane, Ragweed, Spurge, Wild Lettuce, and many other broadleaf weeds.

TIME TO APPLY: Apply to St. Augustine grass in lawn except used for golf greens. Apply in St. Augustine grass in lawn through September when it is growing rapidly. Because temporary foliage discoloration may occur, it is best to apply when weeds are young and growing rapidly. If weeds are growing slowly, control will be slow and more than one application may be needed. Moisture from dew or watering will help control. Do not water immediately after application.

HOW MUCH TO APPLY: Apply 35 lbs. per acre (1000 sq. ft.) for St. Augustine grass and other turf grasses. Do not apply to temporary foliage discoloration may occur. Do not apply to Bermuda, Zoysia, Escalade, and other turf grasses.

GUARANTEED SPREADER SETTINGS

Make sure you apply the correct amount of spreader and adjust the spreader by trial on a known area.

SPREADER SETTINGS	Bentgrass & Bahia Grass	Other Grasses
Rate	3 1/2	4 1/2
Spreader	2 1/2	8 1/2
Spreader	1 1/2	5
Spreader	1 1/2	6 1/2
Spreader	1 1/2	6 1/2

CAUTION: Keep children and pets away from treated and untreated areas. Avoid application in contact with eyes, nose, mouth, or clothing. Do not use on empty bag. Destroy when empty. Do not apply to areas underlain by roots of desirable trees and shrubs.

Apply to areas where plants are newly seeded.

Do not use on lawns treated with insecticides.

516-670

→ (SI) ↑

10

Use herbicides safely
 Do not overdose
 Read DIRECTIONS FOR USE
 carefully

Net Wt. 35 Lbs.

AREATS

Up To 5000 Sq. Ft.



ACCEPTED
July 17-1971
 UNDER THE FEDERAL INSECTICIDE AND RODENTICIDE ACT AND ECONOMIC PESTICIDE REGISTRATION ACT. REGISTERED UNDER NO. 2217. SUBJECT TO ADDITIONAL COMMENTS. 5

ST. AUGUSTINE GRASS LAWN WEEDER & FEEDER

For Use On St. Augustine Grass
 And Other Grasses For
 Control Of Growing Broadleaf Weeds

CAUTION: Keep Out Of Reach Of Children.

See Back Panel for Additional Cautions

WEEDS AS IT FEEDS

ACTIVE INGREDIENTS:

*2,4-dichlorophenoxyacetic acid	0.0820%
**2-(2-methyl-4-chlorophenoxy) propionic acid	0.2952%
***Dicamba (3,6-dichloro-o-anisic acid)	0.0295%
INERT INGREDIENTS	99.5933%
TOTAL	100.0000%

Used at 35 lbs. on 5,000 sq. ft. equivalent to:

- *2,4-D ACID 0.25 lbs/acre or 1.64 lbs/ton
- **MCPP ACID 0.90 lbs/acre or 5.904 lbs/ton
- ***Dicamba ACID 0.09 lbs/acre or 0.59 lbs/ton

U.S. Patent No. 3,284,186

E.P.A. Reg. No. 2217-594

MANUFACTURED BY
PBI - GORDON CORPORATION
 KANSAS CITY, KANSAS