

**WEEDS CONTROLLED:** Bermuda and Bentgrass. Broadleaf Herbicide is especially effective against weeds such as SHEEP SORREL, KNOTWEED, SPOTTED SPURGE, CHICKWEED, OXALIS AND HENBIT. In addition to these, it kills virtually all broadleaf weeds, some of which are listed below.

Bedstraw	Knotweed	Sheep Sorrel
Black Medic	Lambsquarter	Shepherd's Purse
Buckhorn	Lespedeza	
Burdock	Mallow	Speedwell
Chicory	Morning Glory	Spurge
Chickweed	Peppergrass	Wild Carrot
Clover	Pigweed	Wild Garlic
Dandelion	Plantains	Wild Lettuce
Dock	Poison Ivy	Wild Onion
Ground Ivy	Poison Oak	Yarrow
Heal-all	Purslane	
Henbit	Ragweed	

**CAUTION:** Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist. If contact with eyes occurs flush with plenty of water.

Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not contaminate domestic or irrigation water. Do not pour spray solutions near these plants. Do not store in contact with fertilizers, seeds, insecticides, or fungicides. Do not spray on Lippia or dichondra nor lawns or turf where desirable clovers are present. Use only lawn sprayers for this use. Coarse sprays are less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs.

When container is empty, wash thoroughly and destroy. Never re-use. Destroy by perforating or crushing, and burying in a safe place. Do not burn.

**NOTICE:** Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and conditions of application equipment will vary, Seller makes no claims other than those stated on this label.

E.P.A. Reg. No. 2217-597

866-572



**TRIMEC<sup>®</sup>**

# BERMUDA AND BENTGRASS BROADLEAF HERBICIDE

**ACCEPTED**  
 June 19, 1972  
 UNDER THE FEDERAL INSECTICIDE, FUNGICIDE AND WEEDICIDE ACT  
 FOR U.S. AND FOREIGN REGISTERED UNDER 2217-597

**Kills:** DANDELIONS KNOTWEED HENBIT  
 CHICKWEED PLANTAINS SPURGE  
 and Many Other Broadleaf Weeds in All Common Grasses

**Contains: 2,4-D, MCPP and Dicamba**

**KEEP FROM FREEZING**

<b>ACTIVE INGREDIENTS:</b>	
*Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	19.97%
**Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	35.13%
***Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid)	3.74%
<b>INERT INGREDIENTS</b>	41.16%
<b>TOTAL</b>	100.00%
*2,4-dichlorophenoxyacetic acid equivalent 1.6 pounds per gallon or 16.59%.	
**2-(2-methyl-4-chlorophenoxy) propionic acid equivalent 2.8 pounds per gallon or 29.04%.	
***3,6-dichloro-o-anisic acid equivalent 0.3 pounds per gallon or 3.11%.	

**CAUTION: Keep Out of Reach of Children**  
 See Side Panels for Additional Cautions

\*Trimec<sup>®</sup> is a registered trademark of PBI-Gordon Corporation, U.S. Patent No. 3,284,186

866-572

**CONTENTS 1 GALLON**

MANUFACTURED BY

**PBI - GORDON CORPORATION**

KANSAS CITY, KANSAS

66118



TRIMEC™

**BERMUDA AND**

**BENTGRASS**

**BROADLEAF HERBICIDE**

**Kills:** DANDELIONS KNOTWEED HENBIT  
CHICKWEED PLANTAINS SPURGE  
and Many Other Broadleaf Weeds in All Common Grasses

Contains: 2,4-D, MCPP and Dicamba

**KEEP FROM FREEZING**

**ACTIVE INGREDIENTS:**

*Dimethylamine salt of 2,4-dichloro- phenoxyacetic acid	19.97%
**Dimethylamine salt of 2-(2-methyl-4- chlorophenoxy) propionic acid	35.13%
***Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid).	3.74
INERT INGREDIENTS	41.16%
TOTAL	100.00%

\*2,4-dichlorophenoxyacetic acid equivalent 1.6 pounds per gallon or 16.59%  
\*\*2-(2-methyl-4-chlorophenoxy) propionic acid equivalent 2.8 pounds per gallon or 29.04%  
\*\*\*3,6-dichloro-o-anisic acid equivalent 0.3 pounds per gallon or 3.11%.

**CAUTION: Keep Out of Reach of Children**

See Side Panels for Additional Cautions

\*Trimec™ is a registered trademark of PBI-Gordon Corporation,  
U.S. Patent No. 3,284,186

866-572

CONTENTS 1 GALLON

MANUFACTURED BY

**PBI - GORDON CORPORATION**

KANSAS CITY, KANSAS

66118

**DIRECTIONS FOR USE**

**HOW TO USE:** Apply Bermuda and Bentgrass Broadleaf Herbicide in sufficient water for good coverage. Use higher rates when weeds are dense or in flower. If hardy weeds persist, make a second application after three weeks.

Apply when weeds are actively growing. Avoid applying during long excessively dry or hot periods.

When grasses are under stress use reduced rates to reduce potential injury. Do not irrigate within 24 hours of application. Do not apply to newly seeded turf until after the first mowing. Seed can be safely sown 3 to 4 weeks after application at recommended rate. Do not use on dichondra.

If dandelions persist as a special problem, individual spot treatment with a single nozzle hand sprayer is recommended.

**HOW MUCH TO USE:**

**Bermuda and Bentgrass**

On moderately 2,4-D tolerant grasses such as bermuda and bentgrass, use 2 pints per acre. Do not overdose closely mowed bentgrass and note cautions above.

**Bluegrass, Fescue & Zoysia**

On 2,4-D tolerant grasses such as bluegrass and fescue use 2 to 3 pints per acre.

**Centipede and St. Augustine Grass**

On 2,4-D sensitive grasses such as St. Augustine use 1 to 1½ pints per acre. Repeat after 3 weeks if necessary. Do not overdose or apply during hot summer months. Note cautions above.

866-572

ACCEPTED  
June 19, 1972  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND ROVIDENTICIDE ACT  
ED UNDER 221 (b) REGISTER  
TO