

1 of 8
Page 2
Reg # 221-597

AUG 12 1985

Stephen Johnson
FBI/Gordon Corporation
1217 West 12th Street
P.O. Box 4090
Kansas City, MO 64101-9984

Dear Mr. Johnson:

Subject: Amendments-Draft Labeling for Additional Weed and
Hose-End/Container Packaging
Gordon's Trimec 849 General Broadleaf Lawn Herbicide
EPA Registration No. 2217-597
Your Letter and Submission of August 1, 1985

The amendments referred to above, submitted in connection with registration under FIFRA, are acceptable provided that you:

1. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

a. Change the statement

Kills 33 Troublesome Weeds . . .

to read:

Kills 34 Troublesome Weeds . . .

b. Delete "SPECTRICIDE" in the heading - "HOW TO USE IN THE . . ."
Refer to 40 CFR §162.10 (a)(6)(b)(2)(ii) which states:
"No name, brand, or trademark may appear on the label which has not been approved by the Administrator through registration"

2. Submit one (1) copy of your final printed labeling before you release the product for shipment.


RD/DIS:JOB:90967:Ikeda:RD-48:Kendrick&Co:898-1270:8/9/85:8/21/85:DKD

CONCURRENTS

SYMBOL ▶									
SURNAME ▶									
DATE ▶									

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

Richard F. Mountfort 
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (TS-767)

Enclosure



TRIMEC® 849 GENERAL BROADLEAF LAWN HERBICIDE

*change to
34*

Kills ~~23~~ Troublesome Weeds In Lawns ✓

ACTIVE INGREDIENTS:

• Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	10.89%	✓
** Dimethylamine Salt of 2, 4-Dichlorophenoxyacetic acid	5.87%	✓
*** Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)	1.20%	✓
INERT INGREDIENTS	82.04%	
	TOTAL 100.00%	

This Product Contains

- 0.78 lb. 2-(2-methyl-4-chlorophenoxy) propionic acid per gallon or 9.00%
- ** 0.42 lb. 2,4-Dichlorophenoxyacetic acid per gallon or 4.88%
- *** 0.09 lb. 3,6-dichloro-o-anisic acid per gallon or 1.00%
- *** Isomer Specific by AOAC Method 6701-5

ACCEPTED
with comment
AUG 12 1985
EPA Reg. No. 2217-587

Keep Out of Reach of Children
DANGER
See side panels for additional
precautionary statements and
Statement of Practical Treatment

ONE QUART COVERS 8,000 to 16,000 SQUARE FEET

—KEEP FROM FREEZING— ✓

NET CONTENTS ONE QUART

© 1978 and 1980, PBI/GORDON CORPORATION

Change \$
34 →

General Information:

TRIMEC® 849 General Broadleaf Lawn Herbicide

kills 23 troublesome common lawn weeds, including poison ivy, in both northern and southern grasses. This concentrated formulation covers up to 16,000 square feet per quart.

Weeds Controlled:

Bedstraw
Black Medic
Buckhorn
Burdock
Chicory
Chickweed
Clover
Dandelion
Dock

Dollarweed

Ground Ivy
Heal-All
Henbit
Knotweed
Lambsquarters
Lespedeza
Mallow
Morningglory
Pepperweed
Pigweed
Plantains
Poison Ivy
Poison Oak
Purslane
Ragweed
Sheep Sorrel
Shepherds purse
Speedwell
Spurge
Wild Carrot
Wild Garlic
Wild Lettuce
Wild Onion
Yarrow
and many other similar broadleaf weeds



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

DANGER Corrosive. causes eye damage. Do not get in eyes or on skin or clothing. Wear goggles and face shield when handling. Harmful if swallowed or inhaled. Avoid exposure to spray mist. Wash thoroughly after handling.

Statement of Practical Treatment

FIRST AID — In case of eye contact, immediately flush eyes with plenty of water for 15 minutes. Consult a physician immediately.

Environmental Hazards

Do not apply directly to water. Do not apply when weather conditions favor drift from target area.

Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE Keep from freezing. Do not store near food or feedstuffs. Do not store near fertilizers, seeds, insecticides, or fungicides. Store in original container and place in locked storage area when not using.

PRODUCT DISPOSAL Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL Do not reuse can. Rinse thoroughly before discarding in trash.

EPA REG. NO. 2217-597
EPA EST. NO. 2217-KS-1
849-881



DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling

WHEN TO USE:

Spray at any time when weeds are growing. For best results, apply in spring when weeds are small and growing rapidly. Apply in fall to remove late germinating seeds.

HOW TO USE:

Hose-End Sprayer: Use 1 gallon of water for every 250 square feet to be sprayed. For concentrate sprayers, add only the required amount of TRIMEC® 849 listed in the rate chart depending on type of grass and number of square feet to be treated. Do not add water to concentrate sprayers. For dilution sprayers, add the required amount of TRIMEC® 849 and fill sprayer with water to proper level. Spray uniformly.

Pressure-Sprayer: Use 1/2 gallon of water for every 250 square feet to be sprayed. Add the required amount of TRIMEC® 849 listed in the rate chart for the type of grass to be treated and fill sprayer with water to the proper level. Adjust spray nozzle to deliver coarse spray. Spray uniformly.

Spot-Treatment: If hardy weeds persist in sensitive grasses such as St. Augustine, centipede, bermudagrass, or bentgrass, spot treat individual weeds using 1 teaspoon of TRIMEC® 849 in 1 quart of water (equivalent to 4 teaspoons per gallon of water). Minimize contact of grass.

Rates of TRIMEC® 849 for Various Lawn Grasses

Gallons Water Spray		Area to Treat	St. Augustine grass Centipede grass	Bermudagrass Bentgrass	Bluegrass Fescue, Zoysia grass
Hose End Sprayer	Pressure Sprayer				
1		250 sq. ft.	1 fl. oz.	2/3 fl. oz.	2/3 to 1 fl. oz.
4	2	1,000 sq. ft.	2 fl. oz.	2-2/3 fl. oz.	2-2/3 to 4 fl. oz.
8	4	2,000 sq. ft.	4 fl. oz.	5-1/3 fl. oz.	5-1/3 to 8 fl. oz.
20	10	5,000 sq. ft.	10 fl. oz.	13-1/3 fl. oz.	13-1/3 to 20 fl. oz.
32	16	8,000 sq. ft.	16 fl. oz.	21-1/3 fl. oz.	21-1/3 to 32 fl. oz.

Convenient Equivalent Measures

1 fl. oz. = 2 tablespoons or 6 teaspoons
2 fl. oz. = 4 tablespoons
4 fl. oz. = 1 cup
8 fl. oz. = 1 cup
16 fl. oz. = 1 pint
32 fl. oz. = 1 quart

HOW TO USE IN THE SPECTRICIDE LAWN WEED KILLER APPLICATOR:

*Connect sprayer nozzle to garden hose; remove safety cap from container, and insert sprayer unit, securing tightly.

*Turn on water; place finger over hole on right side of nozzle and product will automatically be dispersed. Spray evenly and uniformly over area requiring application. (Check application guide to determine the correct area to be covered per ounce.)

X

*APPLICATION GUIDE			
fl Ounces	Sensitive grasses such as Centipede, St. Augustine	Bermudagrass, Bentgrass	Bluegrass Fescue, Zoysia
Recommended Coverage Area Per ^{fl} Ounce (in sq. ft.)			
1	500	375	250
2	1,000	750	500
3	1,500	1,125	750
4	2,000	1,500	1,000
5	2,500	1,875	1,250
6	3,000	2,250	1,500
7	3,500	2,625	1,750
8	4,000	3,000	2,000
9	4,500	3,375	2,250
10	5,000	3,750	2,500
11	5,500	4,125	2,750
12	6,000	4,500	3,000
13	6,500	4,875	3,250
14	7,000	5,250	3,500
15	7,500	5,625	3,750
16	8,000	6,000	4,000

*Good for this pre calibrated sprayer only.

Use Precautions: Use only lawn sprayer that delivers coarse spray to reduce drift. Avoid spray drift to vegetables, flowers, ornamental plants, shrubs, trees, and other desirable plants. Do not spray roots of trees and ornamentals. Do not exceed specified dosages for any area, be particularly careful within dripline of tree and other ornamental species. Do not spray carpetgrass, dichondra, or lawns where desirable clovers are present. Do not spray while grass is emerging from dormancy. Do not apply to newly seeded lawn grasses until well established. Do not spray when air temperature exceeds 85° F. Seed can be safely sown 3 to 4 weeks after application. Do not contaminate domestic or irrigation waters. After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides. Failure to follow above precautions may result in injury.