



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

June 16, 2017

Mr. Eric Smith
Federal Registration Manager
PBI/Gordon Corporation
P.O. Box 014090
Kansas City, MO 64101

Subject: Notification per PRN 98-10 – Add marketing claims and minor updates
Product Name: Gordon's Trimec 849 Broadleaf Lawn Herbicide
EPA Registration Number: 2217-597
Application Date: January 6, 2017
Decision Number: 525043

Dear Mr. Smith:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped "Notification" and will be placed in our records.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If you have any questions, you may contact Kathryn Montague at 703-305-1243 or via email at montague.kathryn@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Kathryn V. Montague".

Kathryn Montague, Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

GORDON'S TRIMEC® 849 GENERAL BROADLEAF LAWN HERBICIDE

NOTIFICATION

2217-597

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

06/16/2017

EPA Reg. No. 2217-597

ACTIVE INGREDIENTS:

Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid	5.45%
Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	5.88%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)	1.21%
OTHER INGREDIENTS:	87.46%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- 0.39 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 4.50%
- 0.42 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 4.88%
- 0.09 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 1.00%

Isomer Specific by AOAC Methods.

Note: If the design, format, or small size of the container labels makes it impractical to present the entire ingredient statement on the front panel, then add the referral statement: "See back panel for substatement of ingredient statement" and present this substatement on the back panel:

KEEP OUT OF REACH OF CHILDREN

CAUTION

KEEP FROM FREEZING

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

See attached booklet for complete [Precautionary Statements and] Directions for Use, including Storage and Disposal, and back panel for [complete Precautionary Statements, including] First Aid. OR See back panel for complete Precautionary Statements, including First Aid, and Directions for Use, including Storage and Disposal.

Net Contents: _____
EPA Est. No. _____

Company Name and Address:

PBI/Gordon Corporation
1217 West 12th Street
Kansas City, Missouri 64101



PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Wear long-sleeved shirt, long pants, socks, shoes, and gloves. Remove and wash contaminated clothing before reuse.

First Aid	
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If on skin or on clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 - 20 minutes.• Call a poison control center or doctor for treatment advice.
If Inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

[Environmental Hazards (Alternate per PRN 2008-1)

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.]

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Precautions And Use Restrictions

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Do not apply this product through any type of irrigation system.
- Use only lawn sprayer that delivers coarse spray droplets. Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not exceed specified dosages for any area.
- Do not spray carpetgrass, dichondra, or lawn where desirable clovers are present. Do not apply this product to "Floritam" St. Augustinegrass in Florida.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply to newly seeded lawn grasses until well established. Seed can be sown 3 to 4 weeks after application.
- Do not spray when air temperatures exceed 85°F.
- After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.
- Failure to follow above precautions may result in injury.

2. Product Information

Gordon's Trimec® 849 General Broadleaf Lawn Herbicide controls troublesome common lawn weeds, including poison ivy, in both northern and southern grasses.

When to Use: Spray at any time when weeds are growing. For best results, apply in spring when weeds are small and growing rapidly. Apply in fall to remove late germinating weeds.

3. Broadcast Treatments

Limitations on broadcast applications to residential turfgrass:

The maximum application rate is 4.0 fl.oz. of product per 1,000 sq.ft. per application (0.72 lb 2,4-D ae, 0.57 lb MCP-p ae and 0.12 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 fl.oz. of product per 1,000 sq.ft. (1.14 lb 2,4-D ae, 1.06 lb MCP-p ae and 0.24 lb dicamba ae per acre).

3.1 Broadcast Treatments With Ortho Hose-End Sprayers or Gilmour® Dial-A-Mix® Lawn And Garden Sprayers

Add only Gordon's Trimec® 849 General Broadleaf Lawn Herbicide to the sprayer jar. Do not add water to the sprayer jar.

St. Augustinegrass, Centipedegrass, and Bahiagrass:

1. Set the dial - Set dial on 1 1/3 Tablespoon (Tbs.) or 4 Teaspoons (tsp.) per gallon.

2. **Fill** - Determine [Measure] the lawn area to be sprayed. Calculate square feet of rectangular areas by multiplying length by width. The application rate for this product is 2 fl.oz. per 1,000 sq.ft. of lawn. Add an adequate amount of herbicide to the spray bottle to treat the total lawn area.
Example for 5,000 sq.ft. of lawn area: Measure 10 fl.oz. into sprayer jar. Do not add water. Set dial on 4 Teaspoons (tsp.) and spray uniformly on 5,000 sq.ft. of lawn.
3. **Spray** - Spray evenly over the lawn.

Bermudagrass, Buffalograss and Bentgrass:

1. **Set the dial - Set dial on 2 Tablespoon (Tbs.) or 6 Teaspoons (tsp.) per gallon.**
2. **Fill** - Determine [Measure] the lawn area to be sprayed. Calculate square feet of rectangular areas by multiplying length by width. The application rate for this product is 3 fl.oz. per 1,000 sq.ft. of lawn. Add an adequate amount of herbicide to the spray bottle to treat the total lawn area.
Example for 5,000 sq.ft. of lawn area: Measure 15 fl.oz. into sprayer jar. Do not add water. Set dial on 2 Tablespoons (Tbs.) or 6 Teaspoons (tsp.) and spray uniformly on 5,000 sq.ft. of lawn.
3. **Spray** - Spray evenly over the lawn.

Kentucky Bluegrass, Fescue, Perennial Ryegrass and Zoysiagrass:

1. **Set the dial - Set dial on 3 Tablespoon (Tbs.) or 9 Teaspoons (tsp.) per gallon.**
2. **Fill** - Determine [Measure] the lawn area to be sprayed. Calculate square feet of rectangular areas by multiplying length by width. The application rate for this product is 4 fl.oz. per 1,000 sq.ft. of lawn. Add an adequate amount of herbicide to the spray bottle to treat the total lawn area.
Example for 5,000 sq.ft. of lawn area: Measure 20 fl.oz. into sprayer jar. Do not add water. Set dial on 3 Tablespoons (Tbs.) or 9 Teaspoons (tsp.) and spray uniformly on 5,000 sq.ft. of lawn.
3. **Spray** - Spray evenly over the lawn.

3.2 Broadcast Treatments With Hose-End Sprayers With Dilution

Spray preparation for hose-end dilution sprayers: Add the required amount of Gordon's Trimec® 849 General Broadleaf Lawn Herbicide and fill sprayer with water to proper level. Spray uniformly.

Table 1. RATES OF GORDON'S TRIMEC® 849 GENERAL BROADLEAF LAWN HERBICIDE FOR VARIOUS LAWN GRASSES:				
Gallons of Water Spray	Area to Treat	St. Augustinegrass, Centipedegrass, Bahiagrass	Bermudagrass, Buffalograss, Bentgrass	Bluegrass, Zoysiagrass, Fescue
Hose End Sprayer with dilution				
1 gal	250 sq.ft.	0.5 fl.oz.	0.67 fl.oz.	0.67 to 1 fl.oz.
4 gal	1,000 sq.ft.	2 fl.oz.	2.67 fl.oz.	2.67 to 4 fl.oz.
8 gal	2,000 sq.ft.	4 fl.oz.	5.33 fl.oz.	5.33 to 8 fl.oz.
20 gal	5,000 sq.ft.	10 fl.oz.	13.33 fl.oz.	13.33 to 20 fl.oz.
32 gal	8,000 sq.ft.	16 fl.oz.	21.33 fl.oz.	21.33 to 32 fl.oz.

EQUAL MEASURES: 1 fluid ounce (fl.oz.) = 2 tablespoons (Tbs.) or 6 teaspoons (tsp.) 2 fluid ounces = 4 tablespoons 4 fluid ounces = 1/2 cup 8 fluid ounces = 1 cup 16 fluid ounces = 1 pint (pt.) 32 fluid ounces = 1 quart (qt.)

[Application Directions for Ferti-lome Weed-Out Lawn Weed Killer with attached sprayer]

[READY-TO-SPRAY DIRECTIONS FOR USE OUTDOORS ON LAWNS

HOW TO USE THE READY-TO-SPRAY SYSTEM

Connect

1. Shake well before using.
2. Connect sprayer to hose.
3. Turn on water.

Spray

1. To begin spraying, point nozzle in the direction you want to spray.
2. [Bend small plastic tab back and] Turn knob [clockwise] to ON position.

- Spray evenly to the area to be treated. Refer to the [sight gauge] clear view strip (graduated scale) on the side of the container to determine the amount of product sprayed.

Finish

- To stop spraying, [turn knob counter clockwise] to OFF position.
- Turn off water.
- Relieve water pressure by [bending plastic tab back and] turning knob to ON position until water slows to a drip. Then turn knob back to OFF position.
- Disconnect sprayer from hose.

3.3 Broadcast Treatments With Pressure Sprayers

Spray preparation for pressure sprayers: Determine [Measure] the lawn area to be sprayed. Add an adequate amount of herbicide to the sprayer to treat the total lawn area. See Table

Gallons of Water Spray	Area to Treat	St. Augustinegrass, Centipedegrass, Bahiagrass	Bermudagrass, Buffalograss, Bentgrass	Bluegrass, Zoysiagrass, Fescue
Pressure Sprayer				
0.5 gal	250 sq.ft.	0.5 fl.oz.	0.67 fl.oz.	0.67 to 1 fl.oz.
2 gal	1,000 sq.ft.	2 fl.oz.	2.67 fl.oz.	2.67 to 4 fl.oz.
4 gal	2,000 sq.ft.	4 fl.oz.	5.33 fl.oz.	5.33 to 8 fl.oz.
10 gal	5,000 sq.ft.	10 fl.oz.	13.33 fl.oz.	13.33 to 20 fl.oz.
16 gal	8,000 sq.ft.	16 fl.oz.	21.33 fl.oz.	21.33 to 32 fl.oz.

EQUAL MEASURES: 1 fluid ounce (fl.oz.) = 2 tablespoons (Tbs.) or 6 teaspoons (tsp.) 2 fluid ounces = 4 tablespoons 4 fluid ounces = 1/2 cup 8 fluid ounces = 1 cup 16 fluid ounces = 1 pint (pt.) 32 fluid ounces = 1 quart (qt.)

3.4 Broadcast Treatments With Lawn Weed Killer Applicators

HOW TO USE IN THE LAWN WEED KILLER APPLICATOR:

- Connect sprayer nozzle to garden hose; remove safety cap from container and insert sprayer unit, securing tightly.
- Turn on water; place finger over hole on right side of nozzle and product will automatically be dispersed.
- Spray evenly and uniformly over area requiring application. (Check application guide to determine the correct area to be covered per ounce.)

Fluid Ounces	Recommended Coverage Area Per Ounce		
	Sensitive grasses such as Centipedegrass, St. Augustinegrass, and Bahiagrass (sq.ft.)	Bermudagrass, Bentgrass, and Buffalograss (sq.ft.)	Bluegrass, Fescue, and Zoysiagrass (sq.ft.)
1	500	375	250
2	100	750	500
3	150	1,125	750
4	2,000	1,500	1,000
5	2,500	1,875	1,250
6	3,000	2,250	1,500
7	3,500	2,525	1,750
8	4,000	3,000	2,000
9	4,500	3,375	2,250
10	5,000	3,750	2,500

Fluid Ounces	Recommended Coverage Area Per Ounce		
	Sensitive grasses such as Centipedegrass, St. Augustinegrass, and Bahiagrass (sq.ft.)	Bermudagrass, Bentgrass, and Buffalograss (sq.ft.)	Bluegrass, Fescue, and Zoysiagrass (sq.ft.)
11	5,500	4,125	2,750
12	6,000	4,500	3,000
13	6,500	4,875	3,250
14	7,000	5,250	3,500
15	7,500	5,625	3,750
16	8,000	6,000	4,000

*Good for this precalibrated sprayer only.

4. Spot Treatment with Pressure Sprayers

Limitations on spot treatments to residential turfgrass:

Spot treatment is defined as a treatment area no greater than 1,000 sq.ft. per acre. The maximum application rate is 4.0 fl.oz. per 1,000 sq.ft. per application (0.57 lb MCPP-p acid equivalent per acre). The maximum number of spot treatments is limited to 2 per year with a minimum of 30 days between applications.

St. Augustinegrass, Centipedegrass, and Bahiagrass:

- Mix 2/3 fl.oz. (4 teaspoons) in 1 gallon water and uniformly spray 330 sq.ft.
- Do not apply to St. Augustinegrass, centipedegrass, or bahiagrass emerging from winter dormancy; temperatures below 45°F within 72 hours of application may cause damage.
- Do not use on 'Floritam' St. Augustinegrass in Florida.

Bermudagrass, Buffalograss and Bentgrass:

- Mix 1 fl.oz. (2 Tbs) in 1 gallon water and uniformly spray 330 sq.ft. See Table 1 for additional spray preparations.

Kentucky Bluegrass, Fescue, Perennial Ryegrass and Zoysiagrass:

- Mix 1 1/3 fl.oz. (8 teaspoons) in 1 gallon water and uniformly spray 330 sq.ft. See Table 1 for additional spray preparations.

Gallons of water	St. Augustinegrass, Centipedegrass, and Bahiagrass	Bermudagrass, Buffalograss and Bentgrass	Bluegrass, Perennial ryegrass, Zoysiagrass and Fescue	Area to Treat
1	0.67 fl.oz.	1.0 fl.oz.	1.3 fl.oz.	330 sq.ft.
2	1.3 fl.oz.	2.0 fl.oz.	2.6 fl.oz.	660 sq.ft.
3	2.0 fl.oz.	3.0 fl.oz.	4.0 fl.oz.	1000 sq.ft.

EQUAL MEASURES: 1 fluid ounce (fl.oz.) = 2 tablespoons (Tbs.) or 6 teaspoons (tsp.)

OR

Gallons of water	St. Augustinegrass, Centipedegrass, and Bahiagrass	Bermudagrass, Buffalograss and Bentgrass	Bluegrass, Perennial ryegrass, Zoysiagrass and Fescue	Area to Treat
1	1.33 TBS	2 TBS	2.67 TBS	330 sq.ft.

2	2.67 TBS	4 TBS	5.33 TBS	660 sq.ft.
3	4 TBS	6 TBS	8 TBS	1000 sq.ft.

EQUAL MEASURES: 1 fluid ounce (fl.oz.) = 2 tablespoons (Tbs.) or 6 teaspoons (tsp.)

5. Broadleaf Weeds			
Annual yellow sweetclover	Cockle	Narrowleaf plantain	Spiny cocklebur
Aster	Cocklebur	Narrowleaf vetch	Spiny sowthistle
Austrian fieldcress	Common mullein	Nettle	Spotted catsear
Bedstraw	Creeping jenny	Orange hawkweed	Spotted spurge
Beggarticks	Cudweed	Oriental cocklebur	Spurweed
Betony, Florida	Curly dock	Oxalis	Stinging nettle
Bindweed, field	Daisy, English	Parsley-piert	Strawberry, India mock
Bird vetch	Daisy fleabane	Parsnip	Tall nettle
Bitter wintercress	Daisy, oxeye	Pearlwort	Tall vervain
Bittercress, hairy	Dandelion	Pennycress	Tansy ragwort
Bitterweed	Dichondra	Pennywort	Tansy mustard
Black-eyed Susan	Dogfennel	Pepperweed	Tanweed
Black medic	Dollarweed	Pigweed	Thistle
Black mustard	False dandelion	Pineywoods bedstraw	Trailing crownvetch
Blackseed plantain	Falseflax	Plains coreopsis	Tumble mustard
Blessed thistle	False sunflower	Plantain	Tumble pigweed
Bloodflower milkweed	Fiddleneck	Poison ivy	Velvetleaf
Blue lettuce	Florida pusley	Poison oak	Venice mallow
Blue vervain	Frenchweed	Pokeweed	Virginia buttonweed
Bracted plantain	Galinsoga	Poorjoe	Virginia creeper
Bristly oxtongue	Goldenrod	Prairie sunflower	Virginia pepperweed
Broadleaf dock	Ground ivy	Prickly lettuce	Wavyleaf bullthistle
Broadleaf plantain	Gumweed	Prickly sida	Western clematis
Broomweed	Hairy fleabane	Prostrate knotweed	Western salsify
Buckhorn	Hawkweed	Prostrate pigweed	White mustard
Buckhorn plantain	Healall	Prostrate spurge	Wild aster
Bulbous buttercup	Heartleaf drymary	Prostrate vervain	Wild buckwheat
Bull thistle	Heathaster	Puncturevine	Wild carrot
Bullnettle	Hedge bindweed	Purslane, common	Wild four-o'clock
Burclover	Hedge mustard	Ragweed	Wild garlic
Burdock	Henbit	Red sorrel	Wild geranium
Burning nettle	Hoary cress	Redroot pigweed	Wild lettuce
Burweed	Hoary plantain	Redstem filaree	Wild marigold
Buttercup	Hoary vervain	Rough cinquefoil	Wild mustard
Buttonweed	Horsenettle	Rough fleabane	Wild onion
Canada thistle	Jimsonweed	Russian pigweed	Wild parsnip
Carolina geranium	Knawel	Russian thistle	Wild radish
Carpetweed	Knotweed	Scarlet pimpernel	Wild rape
Catchweed bedstraw	Kochia	Scotch thistle	Wild strawberry
Catnip	Lambsquarters	Sheep sorrel	Wild sweet potato
Catsear	Lespedeza	Shepherdspurse	Wild vetch
Chickweed, common	Mallow	Slender plantain	Woodsorrel
Chickweed, mouseear	Matchweed	Smallflower galinsoga	Woolly croton
Chicory	Mexicanweed	Smooth dock	Woolly morningglory
Cinquefoil	Milk vetch	Smooth pigweed	Woolly plantain
Clover, crimson	Morningglory	Sorrel	Wormseed
Clover, hop	Mouseear hawkweed	Sowthistle	Yarrow
Clover, red	Mugwort	Spanishneedles	Yellow rocket
Clover, strawberry	Musk thistle	Speedwell	Yellowflower pepperweed
Clover, sweet	Mustard	Spiny amaranth	and other broadleaf weeds
Clover, white			

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep from freezing. Store in original container and place in locked storage area when not using.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or www.website.com for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

APPENDIX

1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Contents: _____
- EPA Est. No. _____

21. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

- One quart covers 8,000 to 16,000 square feet.
- Controls 34 troublesome weeds in lawns.
- This concentrated formulation covers up to 16,000 square feet per quart.
- One quart of product may treat up to 16,000 square feet of lawns established with St. Augustinegrass, centipedegrass, and bahiagrass.
- Kills (Controls) troublesome [broadleaf] weeds: dandelion, chickweed, clover, and many others as listed.
- Contains Trimec® Herbicide
- In the event of a medical or chemical emergency, contact ChemTel Inc North America 1-800-255-3924

DOCUMENT CONTROL INFORMATION

1. **Unique Label Identifier:** 002217-00597.20170322.notif-proposed-highlighted

2. **Reason for Issue:** ad claims, minor corrections, formatting + EPA Comments #1

Internal Usage Notes
Associated Container Label: Supplemental Distributors