



liquid sevin spray

hoppers. Do not use on Boston Ivy as injury may result. Treat at first sign of insect damage. Repeat at 7 to 10-day intervals or as necessary to maintain control. Wet all leaf surfaces thoroughly.

LAWNS: Mix 1 pint in 15 gallons water, apply to 1250 sq. ft. of turf. Mow lawn and water before spraying. Repeat in 2 to 3 weeks if necessary. For control of chinchbugs, sod webworm (lawn moths), fall armyworms, cutworms, earwigs, fleas and ants.

ADULT MOSQUITOES: Spray shrubbery, hedges, flower beds and other places where mosquitoes may rest during the day. Use 4 tsp. per gallon of spray.

lea beetles.

PEACHES AND APRICOTS: To control oriental fruit moth, plum curculio and cat facing bugs, use 4 tsp. per gallon. May be used 1 day before harvest on peaches 3 days before harvest of apricots.

CAUTION: Harmful if swallowed or inhaled. Avoid breathing of spray mist. Avoid skin contact. Wash face and hands after spraying. Avoid contamination of food and feedstuff. Keep out of reach of children and pets. Do not use in any manner other than specified on this label. Destroy this container when empty. Do not reuse container.

MANUFACTURED BY
PBI - GORDON CORPORATION
KANSAS CITY, KANSAS

ACCEPTED
MAY 1972

DIRECTIONS FOR USE

Shake well before measuring. Keep container closed to prevent caking of contents. Best results are obtained by thinning the measured dosage with water in a separate container before adding to the sprayer. Protect from freezing as bottle may break. Spray all plant surfaces thoroughly.

TREES, FLOWERS, ORNAMENTALS, SHRUBS, SHADE TREES: Use 4 tsp./gallon of finish spray. For control of Japanese beetles, bagworm, rose aphids, exposed thrips, elm leaf beetles, birchleaf miners, rose slugs, box elder bugs, tent caterpillars, lace bugs, leaf rollers, blister beetles, periodical cicada and leaf-



liquid sevin spray

during the day. Use 4 tsp./gallon of spray.

APPLES AND PEARS: To control codling moth, plum curculio, apple maggot and fruit tree leaf roller, use 4 tsp./gallon. Do not apply before second cover spray to avoid fruit thinning. Apply at 10-day intervals in July for apple maggot. May be used up to 1 day before picking. Apply at petal fall and every 10 to 14 days thereafter until control is achieved. Application of Sevin within 30 days after full bloom may also provide apple thinning. To avoid this, delay use until at least 30 days after bloom.

GRAPES: To control grape berry moth and leafhoppers, use 6 tsp./gallon. May be used up to harvest.

STRAWBERRY: Use 4 tsp./gallon up to 1 day before picking to control leaf roller, strawberry weevil and meadow spittle bug.

838-1172

Insecticide for Control of Certain Listed Pests of Shade Trees, Lawns, Ornamentals, Fruit Trees and Vegetables

Contains 2 lbs. actual Carbaryl per gallon. U.S. Patent No. 2,983,478. *Trademark of Union Carbide Corporation for the active ingredient.

ACTIVE INGREDIENT:		
Carbaryl, 1-naphthyl N-methylcarbamate	_____	22.4%
INERT INGREDIENTS	_____	77.6%
		TOTAL 100.0%

CAUTION: KEEP OUT OF REACH OF CHILDREN

See side panels for additional cautions

838-1172

NET CONTENTS ONE QUART U.S. MEASURE

VEGETABLES: Lima and green beans—use 4 tsp./gallon for Mexican bean beetles, leafhoppers, flea beetles and tarnished plant bugs. May be used up to day of harvest. Do not feed vines to dairy or meat animals. Tomato, eggplants, peppers—use 4 tsp./gallon for tomato fruitworms, stinkbugs, flea beetles, leafhoppers and fall armyworms. Apply up to day of picking. Cucumbers, melons and squash—use above dosage for squash bugs, melon worms, cucumber beetles and flea beetles. Do not apply to wet foliage or to watermelons in Florida or injury may result. Potatoes—use 4 tsp./gallon for Colorado potato beetles, leafhoppers and

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. **ENVIRONMENTAL CAUTIONS:** This product is highly toxic to bees exposed to direct treatment or residues on plants. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not spray when plants are in full bloom or when bees are active in the area. Do not contaminate any body of water.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

EPA Reg. No. 2217-600-AA

838-1172