

RAT AND MOUSE KILLER

READY TO USE

ACTIVE INGREDIENT:

Warfarin, 3-(α -acetylbenzyl)-4-hydroxycoumarin..... 0.025%

INERT INGREDIENTS 99.975%


100.000%

CAUTION: Keep Out of Reach of Children
See Back Panel for Additional Cautions

NET WEIGHT FIVE POUNDS



RAT AND MOUSE KILLER



DIRECTIONS

RATS: Place 4 to 6 ounces of bait in dry container next to burrows, in runways, or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 12 days and continue baiting until all signs of feeding have stopped. Replace stale, damp, or moldy baits with fresh bait.

MICE: Place tablespoonful amounts of bait in shallow containers where mice or signs of mice have been seen. Bait stations at intervals of 8 to 12 feet are recommended. Assure an uninterrupted supply of bait for a period of not less than 15 days.

CAUTION: Protect children, pets, wildlife, and domestic animals from bait. If swallowed by humans, pets, or domestic animals, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of dihydroxy coumarin.

DIRECTIONS FOR USE

RATS: Place 4 to 6 ounces of bait in dry container next to burrows, in runways, or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 12 days and continue baiting until all signs of feeding have stopped. Replace stale, damp, or moldy baits with fresh bait.

MICE: Place tablespoonful amounts of bait in shallow containers where mice or signs of mice have been seen. Bait stations at intervals of 8 to 12 feet are recommended. Assure an uninterrupted supply of bait for a period of not less than 15 days.

CAUTION: Protect children, pets, wildlife, and domestic animals from bait. If swallowed by humans, pets, or domestic animals, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

Baits should be placed in areas inaccessible to children, pets, wildlife, and domestic animals, or in tamper proof boxes. This rodenticide contains Warfarin, an anticoagulant chemical that reduces the clotting ability of the blood and, upon repeated feeding, causes fatal hemorrhage in rats and mice. Do not reuse container. Destroy when empty.

NOTE TO PHYSICIANS: When a human has been known to have ingested this product, blood transfusions combined with intravenous injections and oral doses of vitamin K are indicated as in the case of hemorrhage caused by overdoses of Dicumarol.

STORE IN COOL PLACE TO KEEP BAIT FRESH

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

E.P.A. Reg. No. 2217-601-AA 602-673