

ST. AUGUSTINE GRASS WEED & FEED 878

20-10-5

LAWN FERTILIZER **ACCEPTED**

PLUS

TRIMEC[®] HERBICIDAL CONCENTRATE

- KILLS WEEDS -

- FEEDS TENDER GRASSES -

Chickweed, Dandelion, Knotweed, Plantain, Henbit, Spurge, and many other weeds in St. Augustine, Centipede, Bentgrass and most other grasses.

ACTIVE INGREDIENTS:

*Dimethylamine salt of 2,4-dichlorophenoxy acetic acid .17%
**Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid .57%
***Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid) .97%

INERT INGREDIENTS: 99.19%
TOTAL 100.00%

EQUIVALENT TO:

*2,4-dichlorophenoxyacetic acid .14% or 24 lb. per acre
**2-(2-methyl-4-chlorophenoxy) propionic acid .47% or 81 lb. per acre
***3,6-dichloro-o-anisic acid .058% or 10 lbs. per acre

CAUTION: Keep Out of Reach of Children

See Back Panel for Additional Cautions

NET WEIGHT 20 POUNDS

FOR FERTILIZER GUARANTEE SEE REVERSE SIDE

E.P.A. Reg. No. 2217-603-AA

583-1072

MANUFACTURED BY

PBI - GORDON CORPORATION

KANSAS CITY, KANSAS

68118

DIRECTIONS FOR USE

GENERAL INFORMATION: This product contains TRIMEC[®] Herbicidal Concentrate, a special formulation low in 2,4-D for maximum safety to St. Augustine Grass. The patented mixture of herbicides will give maximum broadleaf weed control.

This product can be used on such sensitive grasses as St. Augustine, Bahia, Centipede, Bermuda, and Zoysia. Under conditions of adequate moisture, high temperature and high humidity, overwatering may cause minor transitory discoloration on St. Augustine. Under these conditions, however, the vigorous growth of St. Augustine will choke out most weeds and herbicide treatment would not be necessary.

WHEN TO APPLY: For best weed control, application should be made when weeds are growing vigorously. Application can be made at any time but for maximum safety to St. Augustine grass make application when temperatures are moderate.

COVERS 5,000 SQUARE FEET

HOW MUCH TO USE: This 20 lb. bag should be used on 5,000 square feet. Apply evenly with a lawn spreader either as a one-pass application or a two-pass application, crisscrossing the area at half the rate each time.

To get maximum weed killing results, do not rake, mow, or water the lawn for at least 24 hours after spreading the product.

Lawns which are heavily overrun by weeds and are in poor condition may require more than one application of this product. A suggested second application could be made in approximately 30 days, or spot treat hard-to-kill weeds with a liquid Trimec product.

CAUTION: Harmful if swallowed. Avoid inhaling dust or contact with skin, eyes and clothing. Do not contaminate water intended for domestic or irrigation purposes.

Do not use this product where desirable clovers are present or allow the product to "wind drift" on flowers, vegetables or ornamental shrubbery. Do not apply to newly seeded lawns. During application close spreader openings when turning. Apply when the air is calm. Keep away from fertilizers, seeds, insecticides or fungicides. Do not apply in areas underlain by roots of desirable trees and shrubs. Do not reuse container. Destroy by burying in a safe place. Do not burn.

Do not use on dichondra.

Spreader setting should be calibrated as follows:

Scott	6
Lawn Beauty	5
Jackson	7 1/2
Scars	4

The above spreader settings are approximate. Check your spreader on 100 sq. ft. and adjust to apply on 1/2 lb. per 125 sq. ft. Use normal rate of walk when applying to avoid an error on rate of application.

The following is a partial list of weeds controlled by this product:

Bedstraw	Grooved Ivy	Peppergress	Spindwell
Black Medick	Hedgeall	Plantain	Spurge
Buckhorn	Henbit	Plantain	Thistle
Hyssop	Knotweed	Parsley Ivy	Wild Carrot
Chicory	La. Nettle	Poison Oak	Wild Garlic
Chickweed	Lespedeza	Purple	Wild Lettuce
Clover	Mallow	Ragweed	Wild Onion
Dandelion	Morning Glory	Sheep Sorrel	Yarrow
Doan	Oxalis	Shepherd's Purse	

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

FERTILIZER GUARANTEE

Total Nitrogen (not less than)	20%
Water Insoluble Nitrogen (not less than)	5%
Available Phosphoric Acid (not less than)	10%
Soluble Potash (not less than)	5%

583-1072

BEST DOCUMENT AVAILABLE

ST. AUGUSTINE GRASS WEED & FEED 878

20-10-5

LAWN FERTILIZER

PLUS

TRIMEC[®] HERBICIDAL CONCENTRATE

- KILLS WEEDS -
- FEEDS TENDER GRASSES -

Chickweed, Dandelion, Knotweed, Plantain, Henbit, Spurge, and many other weeds in St. Augustine, Centipede, Bentgrass and most other grasses.

ACTIVE INGREDIENTS:

*Dimethylamine salt of 2,4-dichlorophenoxy acetic acid	.17%
**Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	.57%
***Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid)	.07%
INERT INGREDIENTS:	99.19%

TOTAL 100.00%

EQUIVALENT TO:

*2,4 dichlorophenoxyacetic acid	.14% or 24 lb. per acre
**2-(2-methyl-4-chlorophenoxy) propionic acid	.47% or 81 lb. per acre
***3,6-dichloro-o-anisic acid	.058% or 10 lbs. per acre

CAUTION: Keep Out of Reach of Children

See Back Panel for Additional Cautions

NET WEIGHT 20 POUNDS

FOR FERTILIZER GUARANTEE SEE REVERSE SIDE

E.P.A. Reg. No. 2217-603-AA

583 1071



MANUFACTURED BY

GORDON CORPORATION

KANSAS CITY, KANSAS
66118