Reg # 2217-603

PM-23



1217 W. 12тн STREET P.O. BOX 4090 KANSAS CITY, MO 64101

The Certification of Compliance for the 2,4-D Exposure Reduction Program

"I, being an authorized representative of PBI/Gordon Corporation, certify that all containers of Gordon's Trimec Weed and Feed for Sensitive Grasses (2217-603) produced by June 15, 1994 for products registered for use only on residential or turf sites, excluding sod farms will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products. I further certify that all containers of said product sold or distributed by this company by January 1, 1995 for products registered for use only on residential or turf sites, excluding sod farms will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products."


J. A. Runhmater

2/17/93

James A. Armbruster, Ph.D. Mgr. Regulatory Services

l

Date



816/421-4070 · TOLL-FREE 1-800-821-7925 · FAX: 816-474-0462 · TLX: 362133

GORDON'S TRIMEC(R) WEED & FEED FOR SENSITIVE GRASSES

20-10-5

TWENTY POUNDS COVERS UP TO 5,000 SQUARE FEET

CONTROLS: Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and many other broadleaf weeds in St. Augustinegrass, Centipedegrass, Bahiagrass, Bermudagrass, Zoysiagrass, Bluegrass, Perennial Ryegrass, Fescue and Bentgrass lawns.

ACTIVE INGREDIENTS:
*Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy)
propionic acid
**Dimethylamine Salt of 2,4-Dichlorophenoxyacetic
acid
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic
acid
INERT INGREDIENTS
TOTAL 100.000%

THIS PRODUCT CONTAINS:

l

^{*} 9.531 lbs. or 0.476% 2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton. 2.750 lbs. or 0.137% 2,4-Dichlorophenoxyacetic acid equivalent per ton. 1.171 lbs. or 0.059% 3,6-dichloro-o-anisic acid equivalent per ton.

Isomer Specific by AOAC Methods

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water. For eyes, flush with water for 15 minutes and get medical attention.

See back panel for additional Precautionary Statements.


TRIMEC^(R) is a registered trademark of PBI/GORDON CORPORATION

NET WEIGHT 20 POUNDS

583/1292 APX0000X

EPA REG. NO. 2217-603 EPA EST. NO. 2217-KS-1

Mid. by PBI/GORDON CORPORATION KANSAS CITY, MISSOURI 64101



STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Contraction into the second

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. Wear long pants, long-sleeved shirt and shoes when applying this product.

When using this product, wear long-sleeved shirt, long pants, socks and shoes. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and wasesed skin with soap and water.

ENVIRONMENTAL HAZARIDS: This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from target area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried or dust has settled.

STORAGE & DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PRODUCT DISPOSAL: Securely wrap bag in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

USE PRECAUTIONS: Do not use this product where desirable clovers are present or allow the material to "wind drift" onto flowers, vegetables or ornamental shrubbery. Do not apply in areas underlaid by roots of desirable trees or shrubs. Do not exceed specified dosages for any area; be particularly careful within dripline of tree and other ornamental species. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning. Do not contaminate domestic or irrigation waters.

INSTRUCTIONS: Gordon's Trimec Weed & Feed For Sensitive Grasses controls a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantains, henbit, spurge and many others commonly found in home lawns. In addition, a 20-10-5 fertilizer provides a sufficient supply of plant food needed for a normal season's growth. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may made in approximately 30 days. Do not use on dichondra. Do not overapply, Under conditions of adequate moisture, high temperature and high humidity, over-application may a state of poor growth.

- 1. Mow lawn to normal height 1 to 2 days before application.
- Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
- Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto non-target plants.
- Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

BEST AVAILABLE COPY



SPREADER SETTINGS

Scott	6	
Lawn Beauty	5	
Jackson	7	1/2
Sears	4	

The maximum application rate to turf is 0.3 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

The above spreader settings are approximate. Calibrate your spreader on 250 sq. ft. and adjust to apply at the rate of 1 lb. of product per 250 sq. ft.

20-10-5	
GUARANTEED ANALYSIS	
Total Nitrogen (N)	. 20%
Water Insoluble Nitrogen	. 5%
Available Phosphoric Acid (P205)	. 10%
Soluble Potash (K ₂ O)	. 5%

- WEEDS -

Bedstraw	Knotweed
Black Medic	Lambsquarter
Buckhorn	Lespedeza
Burdock	Mallow
Chicory	Morningglory
Chickweed	Oxalis
Clover	Peppergrass
Dandelion	Pigweed
Dock	Plantains
Ground Ivy	Poison Ivy
Heal-All	Poison Oak
Henbit	Purslane

Ragweed Sheep Sorrel Shepherdspurse Speedwell Spurge Thistle Wild Carrot Wild Garlic Wild Lettuce Wild Onion Yarrow

LIMITED WARRANTY AND DISCLAMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED MAR-RANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These lactors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal' stake(nents or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.



24

.