

**DIRECTIONS FOR USE
RESIDUAL SPRAY IN BUILDINGS**

(Approx. 5 percent). Add 1 part of concentrate to 19 parts of water (about 7 ounces for 1 gallon of emulsion). Apply as course spray on all surfaces at the rate of 10 to 15 gallons per acre or 1 pint per 125 lineal feet on walls, ceilings and other horizontal surfaces, and to runways of ants.

OUTDOOR SPRAY (approx. 1 percent). Add 1 part of concentrate, while stirring to 99 parts of water (about 7 ounces for 5 gallons of emulsion). Apply as course spray. For the control of mosquitoes, flies and sand flies, apply lightly to vegetation, tree trunks, etc. at a rate of 10 to 15 gallons per acre depending on the amount of vegetation and type of terrain to be treated.

For the control of mites (chiggers) and ticks, apply to low vegetation and to ground surface at the rate of approximately 20 gallons per acre depending on the amount of vegetation. For the control of fleas, apply at rate of 10 to 20 gallons per acre at and under buildings. For the control of ants, apply at rate of 10 to 20 gallons per acre.

Insecticide, chlordane, 72 percent, water emulsifiable concentrate

(Approx. 8 mos. chlordane/gallon)

	<i>Percent by weight</i>
Active ingredients:	
Technical chlordane ¹	72 %
Petroleum hydrocarbons	21 %
Inert ingredients-----	7 %
Total	100 %

Specification: O-I-515

Stock number:

Manufacturer's name and address:

Date of pack:

Lot number:

Contract number:

Net contents:

¹Equivalent to 45.3 octachloro-1,7-methano-2,4-dihydro-1,4-dioxin and 28.8% of related compounds.

CAUTION:


DO NOT USE WITHOUT DILUTING

CAUTION: Keep Out of Reach of Children
See Side Panel for Additional Caution
NET CONTENTS 5 GALLON

TERMITE CONTROL. Use 1 percent emulsion as prepared and described on outdoor spray. Apply as course spray on roof. Instructions for termite control. See manual "The Art and Rats Control Plan" (AFM 85-7, TM 5-622, NAVPOCKS TT-Pa-2) for details.

CAUTION:

Harmful if swallowed. Repeated or prolonged contact with skin can cause toxic symptoms. Avoid breathing spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of food and foodstuffs. Keep out of reach of children. In living quarters, use as strip treatment (not for complete coverage of all walls). To be applied by trained personnel only. When extensive use is involved, operators should wear respirators approved by the U. S. Dept. of Agriculture. See manual "The Art and Rats Control Plan" (AFM 85-7, TM 5-622, NAVPOCKS TT-Pa-2) for details.



... ..
... ..
... ..


10% chlordane dust

GARDEN SOIL: Apply one pound per 1000 square feet. Work into top 2 to 4 inches of soil while preparing seed beds to control cutworms, white grubs and wireworms. Make a second application on soil surface around cabbage and tomatoes after transplanting if cutworms are a problem. Do not use in soil where root crops or green onions are to be planted. Apply before seeding beans, peas and corn. For surface application use one pound per 400 to 500 square feet.

ORNAMENTALS AND FLOWERS: To control thrips, grasshoppers, ants and plant bugs make a light application as soon as insects are noticed. Repeat as necessary to prevent damage.

LAWNS: Apply one pound per 900 to 1000 square feet to control ants and chiggers. Repeat as necessary for control. To control white grubs, Japanese beetle larvae and sod webworms (larvae).

MANUFACTURED BY
FBI - GORDON CORPORATION
KANSAS CITY, KANSAS



10% chlordane dust

ACTIVE INGREDIENT:
Technical Chlordane* 10%
INERT INGREDIENTS 90%
*(Contains 6% Octochloro-4,7-methanotetrahydroindane and
4% related compounds)

CONTROLS
ANTS, CHIGGERS, WHITE GRUBS,
THRIPS, WIRE WORMS

CAUTION: Keep Out of Reach of Children
See Side Panel for Additional Cautions

NET WEIGHT — ONE POUND

626-273

Use 10% Chlordane Dust as it comes from the package using a hand duster.

For control of cutworms and sod webworms (lawn moth) use one pound per 400 square feet. Water well after dusting to move material into soil zone where these pests live. Keep children and pets off treated areas until insecticide has been washed into the soil and the grass is thoroughly dry. Birds feeding on treated areas may be harmed.

BULBS AND CORNS: After digging and removing excess foliage and dirt dust tulips, hyacinths, gladiolus and dahlias to control thrips before storing. Treat again just before planting.

HUMAN CAUTIONS: Harmful if swallowed. Avoid breathing of dust. Wash with soap and water immediately after use. Do not contaminate feed or foodstuffs. Do not use on or around vegetables after edible parts begin to form. Keep out of reach of children. Do not apply in any manner not specified on this label.

ENVIRONMENTAL CAUTIONS: This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds. Do not contaminate water by wastes or treat areas with heavy runoff. This product highly toxic to bees. Do not dust when bees are visiting the area.

Do not reuse this container for any purpose. Destroy when empty.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary. Seller makes no claims other than those stated on this label.

E.P.A. Reg. No. 2217-616-AA

626-273