

Gordon's TRIMEC 450 HERBICIDE

ACCEPTED
JUN 18 1981
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 249-104-0124

KILLS: DANDELIONS KNOTWEED HENBIT
CHICKWEED PLANTAINS SPURGE
and Many Other Broadleaf Weeds

Contains 2,4-D, MCPP, and Dicamba
KEEP FROM FREEZING

ACTIVE INGREDIENT
††Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid 25.0%
††Dimethylamine Salt of 2,4,5-Trichlorophenoxyacetic acid 15.0%
††Dimethylamine Salt of 2,4-Dichlorophenoxypropionic acid 10.0%
INERT INGREDIENT 50.0%

This Product Contains
††1.88 lbs. 2,4-Dichlorophenoxyacetic acid
††2.0 lbs. 2,4,5-Trichlorophenoxyacetic acid
††0.9 lb. 2,4-Dichlorophenoxypropionic acid
††Isomer Specific by AOAC Method 8011
*TRIMEC is a registered trademark of PBI/GORDON CORPORATION, Inc.
Patent No. 3,284,196

KEEP OUT OF REACH OF CHILDREN
DANGER
STATEMENT OF PRACTICAL TREATMENT
Read the entire label carefully before using this product. Do not use if the label is torn, mutilated, or missing any instructions or warnings. Do not use if the product is frozen.

NET CONTENTS ONE U.S. GALLON

Made by **G** pbi/gordon
CORPORATION
KANSAS CITY, MISSOURI

USE INSTRUCTIONS

ORNAMENTAL LAWNS AND TURF: Apply TRIMEC 450 HERBICIDE at the rate of 2.0 to 3.0 pints per 1,000 sq. ft. of water for adequate coverage. Maximum application will be 3.0 pints per 1,000 sq. ft. in early Fall application when weeds are actively growing. Avoid applying during long, excessively dry or hot periods unless adequate irrigation is available. Do not irrigate within 24 hours of application. Do not apply to newly seeded turf until after the second or third mowing. Grass seed can be safely sown 3 to 4 weeks after application at recommended rate. Do not use on dichondra.

WEEDS CONTROLLED: TRIMEC 450 HERBICIDE is especially effective against weeds such as Sheep Sorrel, Knotweed, Spotted Spurge, Chickweed, Oxalis, and Henbit. In addition to these, it kills virtually all broadleaf weeds, some of which are listed below. Repeat application if necessary.

Bedstraw	Knotweed	Sheep Sorrel
Black Medick	Lambsquarter	Shepherd's Purse
Buckhorn	Lespedeza	Speedwell
Burdock	Mallow	Spurge
Chicory	Morning Glory	Wild Carrot
Chickweed	Peppergrass	Wild Garlic
Clover	Pigweed	Wild Lettuce
Dandelion	Plantains	Wild Onion
Dock	Poison Ivy	Yarrow
Ground Ivy	Poison Oak	
Heal all	Purslane	
Henbit	Ragweed	

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no warranty other than those stated on this label.

© 1981 PBI/GORDON CORPORATION
KANSAS CITY, MISSOURI

823 581