

DIRECTIONS FOR USE

TURF: Use 2 to 3 pints Amine 2,4-D Turf Herbicide in 10 gallons of water per acre. Apply in the spring when weeds are actively growing to obtain maximum kill, and again in early summer as certain weeds do not appear until warm weather. Use 1 1/2 lbs. per gallon of water per 1,000 square feet to treat lawns. Do not apply to newly seeded areas until grass is well established. Do not spray new lawns until after first cutting. Clover may be damaged with this spray. **Do not** use on Bentgrass, St. Augustine grass, Dichondra, or carpetgrass lawns.

DON'T OVER DOSE—Read the label directions and cautions carefully. Do not let spray drift to sensitive plants such as vegetables, grapes, flowers, or other 2,4-D susceptible plants. Box Elder and Red Bud trees are also injured by spray drift. Use a coarse spray and make treatments only when there is minimum chance of "wind drift." Under certain high temperatures vapors from this product will injure nearby susceptible plants.

ROADSIDE AND GENERAL WEED CONTROL: To control broadleaf weeds along roads, right-of-ways, drainage ditch banks, fence rows, and other similar non-crop areas, mix 2 to 3 pints in 20 gallons of water with 5 to 8 oz. liquid detergent and thoroughly wet weed foliage. Do not let spray drift. Treat in the spring and again just before bloom time before weeds go to seed. Repeat the following year.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

EPA REG. NO. 2217-633-AA

EPA EST. NO. 2217 KS-1




AMINE 2,4-D TURF HERBICIDE

FOR ORNAMENTAL TURF

Contains 4 lbs. 2,4-D acid per gallon

ACTIVE INGREDIENT:

*Dimethylamine Salt of 2,4-Dichloro- phenoxyacetic acid	49.3%
INERT INGREDIENTS	50.7%
TOTAL	100.0%
*2,4-Dichlorophenoxyacetic acid equivalent	41.0%



CAUTION: Keep Out of Reach of Children

See Side Panels for Additional Caution Statements.

SUSCEPTIBLE WEEDS

(use lower dosage rate)

Annual sow thistle	Knotweed
Annual thistle	Kochia
Bindweed	Lambsquarter
Buckhorn	Pigweed
Bull thistle	Ragweed
Carpet weed	Wild Lettuce
Chicory	Wild Radish
Dandelion	Vervain

WEEDS MORE DIFFICULT TO KILL

(use higher dosage rate)

- Climbing milkweed
- Dog fennel
- Ground Ivy
- Heal-all mallow
- Mouse ear chickweed
- Nettle
- Wild Aster
- Wild Onion

CAUTION: Harmful if swallowed! Do not get in eyes, on skin, or on clothing. In case of contact, wash with soap and water. In case of contact with eyes, flush with plenty of water and get medical attention. Avoid inhalation of spray mist.

Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees, and other desirable plants. Do not pour spray solution near these plants. Do not spray on bentgrass, St. Augustine grass, Dichondra or carpetgrass lawns or turf, or where desirable clovers are present. Use coarse sprays as there is less chance of wind drift. Do not apply to areas underlaid by roots of desirable trees and shrubs. Do not contaminate domestic or irrigation waters. Do not store this product or its spray solutions near fertilizers, seeds, insecticides, or fungicides. Always use separate equipment for insecticides and fungicides.

CONTAINER DISPOSAL: Do not re-use container. Destroy when empty. When empty, wash thoroughly with water and detergent, and discard in a safe place away from water supplies, preferably by perforating or crushing.

814 775