

2217-638

RESTRICTED USE PESTICIDE

Due to Acute Toxicity

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

This product must not be used on golf courses and sod farms.

GORDON'S DIAZINON 25% EMULSIFIABLE

LAWN & GARDEN INSECT CONTROL

PROTECTS: Lawns, Trees, Shrubs, Roses, and Flowers.

CONTROLS: Aphids, Chinch bugs, Cutworms, Mites, Sod Webworms, and many others as listed.

ACTIVE INGREDIENTS:

Diazinon [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate]..... 25.0%

INERT INGREDIENTS..... 75.0%

TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN

W A R N I N G

Statement of Practical Treatment

IF SWALLOWED: Call a physician immediately. Gastric lavage is not indicated if material was taken internally. DO NOT INDUCE VOMITING unless other treatment is not available. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. Apply respiration if indicated.

IF ON SKIN: Wash with soap and water. Rinse thoroughly.

IF IN EYES: Flush eyes with plenty of water for at least 15 minutes. Get medical attention immediately.

See side panels for additional precautionary statements.

COMBUSTIBLE MIXTURE, DO NOT USE IN THE HOME.

NET CONTENTS ONE U.S. GALLON


895/489 AP

EPA REG NO 2217-638

EPA EST NO XXXX-XX-X

Mfd. by PBI/GORDON CORPORATION
KANSAS CITY, KANSAS 66118

Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.



STOP! READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

WARNING: May be fatal if swallowed. Keep out of the reach of children. May be absorbed through the skin. Do not breathe spray mist. Avoid contact with skin, eyes and clothing. Wash hands thoroughly after handling and before smoking or eating. Do not use on household pets or humans. Do not permit children or pets to go onto sprayed areas until the spray has completely dried. Do not use in the home. Do not contaminate ornamental fish ponds.

NOTE TO PHYSICIAN: If signs of cholinesterase inhibition are present, atropine is antidotal, 2-PAM is also antidotal and may be administered in conjunction with atropine.

May cause contact sensitization following repeated contact with skin in susceptible individuals. Avoid repeated contact with skin. If sensitization reactions result consult a physician.


Environmental Hazards: This pesticide is highly toxic to birds, fish and wildlife including waterfowl. Birds and waterfowl feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product in Nassau County, New York between November 1 and May 20. Do not exceed maximum permitted label rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. Where irrigation (watering) is recommended apply a minimum of 0.25 inch of water immediately after application. Stop irrigation before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries (except for effluence from treated watercress beds). Do not apply to water that will be used for recreational purpose and human and livestock consumption. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

PHYSICAL OR CHEMICAL HAZARDS: Do not use, pour, spill or store near heat or open flame.

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

215



Outdoor Domestic Use: Wear a long sleeve shirt and long legged pants. Spray with the wind to your back. Do not spray on windy days (wind speed is greater than 10 miles per hour). Remove clothing after spraying and launder before reuse. If clothing becomes wet from spray, **stop spraying**, immediately remove clothes and shower with soap and water.

Do not allow children or pets onto a treated area until the spray has dried.

STORAGE & DISPOSAL

STORAGE: Store in original container in a locked storage area inaccessible to children or pets.

PRODUCT DISPOSAL: Do not contaminate food or feedstuffs by storage or disposal. Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse container. Rinse thoroughly before discarding in trash.

GORDON'S DIAZINON 25% EMULSIFIABLE kills many outdoor insects which destroy lawns, roses, flowers, trees, and shrubs or annoy pets and people. Refer to the label directions for specific insects controlled. Diazinon insecticide provides positive control of many "hard-to-kill" sucking and chewing insects.

HOW TO MIX & APPLY

1. Determine size of spraying job. For lawns and vegetables, measure the area to be sprayed. For all other applications, apply as a thorough cover spray. Make applications when insects first appear. Repeat applications as needed or as noted in the rate chart.
2. Use clean sprayer. Carefully measure and mix the amount of Diazinon and water as suggested in the rate chart.
3. Spray as directed in rate chart. Use coarse drenching spray for lawns, soil and other surface applications. Use medium fine mist sprays for trees and plants. Thorough coverage is important. Flush sprayer with clean water after each use.

NOTE: Use the contents of this package within 6 months after opening.

See chart for complete insect list - use dosages shown unless otherwise indicated.


1 fl. oz. = 6 tsp. or 2 tbsp.

8 fl. oz. = $\frac{1}{2}$ pint or 1 cup

1 fl. oz. in 3 gals. water = 2 tsp. in 1 gal water or

8 fl. oz. in 24 gals. water

NOTE: Do not use on ferns, poinsettia, hibiscus, papaya, pilea gardenia and orchids since injury may occur.



USE DIRECTIONS

LAWNS & TURF: For use on Bentgrass, Bermuda, Bluegrass, Dichondra, Fescue, Marion, St. Augustine, and Zoysia Grass.

Ants, Armyworms, Bermudagrass Thrips, Brown Dog Ticks*, Chiggers, Crickets, Cutworms, Earwigs, Fleas, Frit Flies, Chinch Bugs, Lawn Moths (Sod Webworms), Leafhoppers, Mites (Bermuda, Clover), Sowbugs (Pillbugs), Springtails (Collembola):** Thoroughly wet down grass a few hours before applying. Apply at a rate of 1 fluid ounce in 3 gallons of water to cover 125 sq.ft. or 8 fluid ounces in 24 gallons of water to cover 1,000 sq.ft. For the Chemlawn Liquid Applicator, use 32 ounces (one quart) in 2½ gallons of spray and cover 4,000 sq.ft.

*Brown Dog Ticks - Do not apply to animals. In particular, spray under shrubbery and the grass next to house.

**In the North, if Chinch Bugs are the only lawn insect problem, a lower rate of application can be used (½ fluid ounce in 3 gallons of water covering 125 sq.ft.).

Lawn Billbugs - Apply at the rate of 1 fluid ounce in 3 gallons of water, covering 125 sq.ft. Water the lawn thoroughly immediately after treatment.

Millipedes: Use 2 fluid ounces in 3 gallons of water, covering 126 square feet. Thoroughly wet grass a few hours before applying.


Rhodengrass Scales: Use 1 fluid ounce in 3 gallons of water, covering approximately 100 sq.ft. Apply when Scale Crawlers first emerge.

Ants, Digger Wasps: Use 1 fluid ounce in 3 gallons of water, covering 125 sq.ft. Spot spray ant hills and wasp ground nest openings. Make application when the wasps are not active.

Billbugs, White Grubs, such as Japanese Beetle, European Chafer, Southern Chafer: Use 1 fluid ounce in 3 gallons of water to cover 125 sq.ft. Water grass thoroughly after application. For billbug control apply when activity is first observed or when chewed or brown grass is present indicating damage from the insect. For grub control make application anytime between late July and early October.

Hyperodes Weevil (New York only): Use ¼ fluid ounce in 3 gallons of water to cover 125 sq.ft. Water grass thoroughly after application. Treat problem areas in mid April and again in mid-May.

TREES & SHRUBS: For use on Arborvitae, Azalea, Birch, Boxwood, Camellia, Douglas Fir, Elm, Hawthorn, Holly, Juniper, Lilac, Locust, Maple, Mimosa, Oak, Pine, Plum, Poplar, Rhododendron, Spruce, Willow.



To Control Aphids, Bagworms, European Pine Shoot Moths, Holly Bud Moths, Dipterous Leaf Miners (Arborvitae, Azalea, Birch, Boxwood, Holly, Oak), Leafhoppers, Mealybugs, Mites (Clover, Cyclamen, European Red, Privet, Two-spotted Spider), Obscure Root Weevils, Omnivorous Leaf Tiers, Scale Crawlers (Cottony-cushion, Eunonymus, Locanium, Pine Needle, Pit Scale, San Jose, Soft): Use 1 fluid ounce in 3 gallons of water. Apply as a thorough foliar spray. Wet leaves and branches to the dropping point. Try to penetrate dense foliage.

Webworms (Juniper, Mimosa), **Whiteline**: Apply at a rate of 1 fluid ounce in 3 gallons of water. Apply as a thorough foliar spray. Wet leaves and branches to the dropping point. Try to penetrate dense foliage. For Mimosa Webworms, apply when adults first appear. Make two additional applications at 4-5 week intervals.

Apples and Thorn Skeletonizers, Hemlock, Charmes, Oak Loopers, Oblique Banded Leaf Rollers, Pear Slugs, Tent Caterpillars, Webworms (Cotoneaster, Fall): Apply at a rate of 3 fluid ounces in 3 gallons of water. Apply as a thorough foliar spray. Wet leaves and branches to the dropping point. Try to penetrate dense foliage.

ROSES & FLOWERS: For use on Azaleas, Camellias, Chrysanthemums, Gladioli, Lilac, Rhododendron, and Roses.

Aphids, Euonymus Scale Crawlers, Flea Beetles, Dipterous Leaf Miners, Leafhoppers, Mealybugs, Mites (Carnation Bud, Carnation Shoot, Cyclamen, European Red, Two-spotted Spider), **Thrips, Whiteflies**: Apply at the rate of 1 fluid ounce in 3 gallons of water. Apply as a thorough foliar spray, wetting the plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage.


NUISANCE PESTS IN OUTSIDE AREAS:

Ants, Boxelder Bugs, Brown Dog Ticks, Chiggers, Clover Mites, Cockroaches, Crickets, Earwigs, Fleas, Flies, Millipedes, Sowbugs (Pillbugs), Spiders, Springtails (Collembola): Apply at the rate of 8 fluid ounces in 3 gallons of water. Thoroughly spray window and door screens, sills, foundations, porch and patio walls and ceilings, entranceways, walks, garbage cans, tree trunks, and into cracks and other places where insects can hide. Do not treat plants or animals. For better control of most of these pests, spray a 5 foot band of soil around the house next to the foundation and spray the foundation of the house to a height of 2-3 feet. See lawn recommendations for other control measures for these pests.

FIRE ANTS:

To aid in the control of fire ants in lawns and other recreation areas, apply 1 fluid ounce (2 tbsp) in a minimum of 1 gallon of water as a drench. Apply over and 6 inches around each mound.

6



Apply gently to avoid disturbing the ants. Reapply as necessary. Larger quantities of solution can be prepared for multiple mound treatments. Use equipment capable of delivering the diluted product as a gentle rain at the rate of 1 gallon per mound. High pressure sprays may disturb the ants and cause migration, reducing product effectiveness. Note: For best results apply in cool weather, 65-80°F, or in early morning or late evening hours. Treat new mounds as they appear.

LIMITED WARRANTY. Manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled use when applied according to label directions. Because of widely varying use conditions, it is impossible to eliminate all risks even when label directions are followed.

EXCLUSION OF OTHER WARRANTIES AND REMEDIES. Except where such disclaimers and exclusions are specifically prohibited by applicable law. THE FOREGOING IS THE ONLY GUARANTEE OR WARRANTY APPLICABLE TO THIS PRODUCT AND IS GIVEN EXPRESSLY AND IN LIEU OF ALL OTHER WARRANTIES EXPRESSED OR IMPLIED, OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, AND ALL SUCH WARRANTIES WHICH EXCEED OR DIFFER FROM SAID LIMITED WARRANTY ARE DISCLAIMED BY MANUFACTURER, and, upon Manufacturer's compliance with said limited warranty. BUYER SHALL HAVE NO REMEDY AGAINST MANUFACTURER FOR ANY TYPE OF DAMAGE OR LOSS, and, IN NO EVENT SHALL MANUFACTURER BE LIABLE FOR CONSEQUENTIAL DAMAGE OR LOSS.

