Please read instructions on r		03/	10/200	5 Form Ann	roved	OMB No. 2070	-0060	
€ЕРА	Environme	United States ntal Protectic Ishington, DC 2046				Registration Amendmer Other	n	OPP Identifier Number
		Application fo	or Pesticio	de - Secti	ion I			
1. Company/Product Num			2. EPA	Product Mar	•		3. Prop	oosed Classification
	2217-640			Joanne	I. Mil	ler		
4. Company/Product (Name) Gordon's Trimec [®] Weeder Granules			PM# Pro	duct Manag	aer—	Team 23		None Restricted
5. Name and Address of A			6. Expedited Review. In accordance with FIFRA Section 3(c)(3)					
PBI/Gordon Corporation			(b)(i), my product is similar or identical in composition and labeling to:					
Post Office Box 0140			EPA Reg. No					
Kansas City, Missour	is is a new address		Produc	t Name				
	is is a new address		ection - II					
		3			abels i	in response to		
Amendment - Explain				Agency letter	dated		NO	FIFICATION
Resubmission in resp	onse to Agency letter d	ated		"Ме Тоо" Арр				R 1 0 2005
Notification - Explain i	below.			Other - Explai	in belo	w.	1071	102003
2. We ask to add additional claim This notification is consistent labeling or Confidential State the EPA. I further understan FIFRA and I may be subject 1. Material This Product Will Child-Resistant Packaging Yes* No * Certification must be submitted 3. Location of Net Contents In	cide Registration a third alternate set is is presented in it one advertising classing classing classing classing classing classing classing classing classing the provisions of F ment of Formula of this dithat if this notification to enforcement action a Be Packaged In: Unit Packaging Yes No If "Yes" Unit Packaging wgt. iformation ainer	Notice (PRN) 98 at of spreader set is entirety on pag aim which may be ed on page 7 of f <u>PR Notice 98-10</u> and product. I understan is not consistent with ind penalties under S Se Se No. per container 4. Size(s) Retail Co Lithograph Paper glued Stenciled	B-10: ttings which ge 4 of the en- e presented the enclosed EPA regulation d that it is a vio the terms of <u>P</u> ection 12 and 12 ection - 11 fater Soluble Pa Yes No "Yes" ackage wgt.	No. per container	eling c 52.46, .S.C Sc <u>Q</u> and 2 5. Loca	of our supple , and no other cl ec. 1001 to willfi	mental nanges h ully make this pro iner iner	distributors. This have been made to the e any false statement to duct may be in violation of
1. Contact Point (Complete it	erns directly below for id	tentification of individ			sary, to	<u> </u>		
Name James L. H	Kunstman, Ph.D.		Director of F	Regulatory	Servi		8	o. <i>(Include Area Code)</i> 16-460-6292
	knowingly false or misle	ading statement may 3. Titl D	<i>v be punishable</i> e irector of Re	by fine or imp	orisonn	nent or	6.	Date Application Rec.alved (Stamped)
4. Typed Name James L. Kunstman, Ph.D.			Februar	y 21, 2005				
EPA Form 8570-1 (Rev. 3-94	Previous editions are	obsolete.		White	- EPA	File Copy (orig	ginal) N	eilow - Applicant Copy

2/8 NOTIFICATION MAR 1 0 2005

GORDON'S TRIMEC[®] WEEDER GRANULES

ACTIVE INGREDIENTS:1.37%2,4-dichlorophenoxyacetic acid.1.37%(+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid.0.31%Dicamba: 3,6-dichloro-o-anisic acid.0.13%INERT INGREDIENTS:98.19%TOTAL100.00%

THIS PRODUCT CONTAINS:

27.4 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.

6.2 lbs. (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid equivalent per ton.

2.6 lbs. 3,6-dichloro-o-anisic acid equivalent per ton.

Isomer Specific by AOAC Methods

Contains the single isomer form of Mecoprop-p.

TRIMEC[®] is a registered trademark of PBI/Gordon Corporation.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for Precautionary Statements.

NET WEIGHT 10 OR 20 POUNDS

650/ AP022105 EPA REG. NO. 2217-640 EPA Est. No. 2217-KS-1 MANUFACTURED BY:



Telephone: 1-800-821-7925



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hand and exposed skin with soap and water.

First Aid

If in eyes:	 Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing:	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.
If swallowed:	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

ENVIRONMENTAL HAZARDS:

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater. Do not contaminate domestic or irrigation waters. Do not apply when weather conditions favor drift from target area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until dust has settled.

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs. **DISPOSAL:**

If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partially filled: Call your local solid waste agency or toll free number (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.

USE PRECAUTIONS:

Do not allow the material to drift onto flowers, vegetables or ornamental shrubbery. Apply when the air is calm. Do not apply to newly seeded lawns. Do not apply in areas underlaid by roots of desirable trees and shrubs. Do not exceed specified dosages for any area; be particularly careful within dripline of tree and other ornamental species. During application, close spreader openings when turning. Do not contaminate domestic or irrigation waters.

INSTRUCTIONS:


Gordon's Trimec[®] Weeder Granules controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low meisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days.

Established turf of warm season grasses includes bermudagrass, centipedegrass, and zoysiagrass. Established turf of cool season grasses includes Kentucky bluegrass, perennial ryegrass, tall fescue, and turf-type tall fescues. Do not apply this product on carpetgrass, dichondra, St. Augustinegrass, or established turf containing desirable clovers.

The maximum application rate to turf is 1.2 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

- 1. Mow lawn to normal height 1 to 2 days before application.
- 2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
- 3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants.
- 4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

Option 1: Coverage and Spreader Settings



The above spreader settings are approximate. Calibrate your spreader on 100 square feet and adjust to apply at the rate of 1 pound of product per 500 square feet.

Option 2: Alternate Coverage and Spreader Settings

COVERAGE: To determine how much fertilizer you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct any non-lawn areas (house, garden, etc.). One bag covers 5,000 sq.ft. [Length x Width = Total Sq.Ft. - Non-Lawn Areas = Total Sq.Ft. Coverage You Need]

SPREADER SETTINGS:

IMPORTANT: It is important to follow the directions as indicated on this quality lawn care product as over application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. Be sure to check spreader before beginning, adjusting if necessary. Consult Owner's Manual. *Never apply by hand.*

Spreading Hints: These granular settings are approximate. Calibrate your spreader on 100 sq.ft. and adjust to apply at the rate of 1 lb. of product per 500 sq.ft.. This bag covers _____ sq.ft. of lawn area.

To avoid gaps and streaks, first spread border of lawn and then fill in center with uniform, parallel passes. Walk at a steady pace so as to apply evenly throughout the lawn. [Graphics and/or diagrams may be presented on the container label].

Rotary Spreaders	Setting
Ames	see spreader instructions
EV-N-Spread	13
Ortho	3*
Precision	5
Quaker	4
Republic EZ	7 1/4
Republic EZ Grow	3
Scotts Easy Green	27 1⁄4
Scotts Speedy Green	5
Scotts Speedy Green 1000	6
Scotts Speedy Green 2000	7
Scotts Speedy Green 3000	6
True Temper	4
Vigoro	7 1/4

Drop Spreaders	Setting		
Ames	see spreader instructions		
Bandini	2		
Ortho	6 1/4		
Precision	6 1/2		
Republic EZ	6 1/4		
Scotts AccuGreen	8 3/4		
Scotts Drop	6 1/2		
Scotts Even Green	6 1/2		
Vigoro	2 1/2		

Spreader Settings

before making hard turns.

The spreader settings below are approximate. Calibrate your spreader on 100 sq. ft. and adjust to apply at the rate of 1 pound of product per 500 sq. ft. (2 lbs. of material per 1.000 sq. ft.). Rate of walk, moisture and spreader candition can vary the spreader setting necessary to apply the correct amount.

	Broadcast Spreaders	Setting			
	Apri-Feb* Rotany	5			
	Cycloner Rolery	- Ă			
	Earthway Rotary	16	~		
	Greenwiew* Jet-Sored* SS1	16			
and the second	Lesco* Rotary (dial A-S)/(number dial				
	Red Devil® Rotary (1-20)	5			
Anstron a ginale negative at a rate	Republic* Rotary	2			
Apply in a single pass at a rate	Scotte* EasyGreen*	26			
of 2 lbs. per 1,000 sq. ft. Start on	Scotta* EdgeGuard*	5			
the outside border and apply in a	Scotter SpeedyGreen* 2000, 3000	5			
circular path working toward the		3 5			
center. Overlap the coverage	Sears* Craftsmen Rotary				
	Sta-Green* Rotary (1-10)	3	-		
slightly, and shut off spreader	True Temper* Rotary Vigoro* Rotary (1-10)	4			
before making hard turns.	2.000 (1000) (100)				
	Drop Spreaders	Setting	\bigcap		
	Age-Fab* Drop	4	14		
	Greenwew* Drop-Spred* XP1	16	- F A		
	Red Dew/* Drop (1-5)		-k		
n de la constante de la constan Martin de la constante de la cons	Republic* Drop	5%			
	Scotts* Drop AccuGreen (2-18)	5 12			
	Scots* Drop PF1 (2-18)	7%			
Apply in a single pass at a rate	Sta-Green* Drop (1-5)	4			
of 2 lbs. per 1.000 sq. ft. Start on					
the outside border and apply in a rectangular path working back and forth across the length of the lawn. Overlap the coverage slightly, and shut off spreader	If your spreader model does not appear above, go to our website at www.preen.com, or call toll-free (800) 233-1067, 8 A.M 5 P.M. EDT, Monday - Friday, for additional model settings.				

[Note: Additional spreader models and settings may be added or substituted by the supplemental registrant provided they deliver the labeled rate of product.]

[Note: Spreader settings may be varied by the supplemental registrant as different grades of fertilizer are used, provided they deliver the labeled rate of pesticide.]

Annual yellow sweetclover Aster Austrian fieldcress **Bedstraw** Beggarticks Betony, Florida Bindweed, field Bird vetch **Bitter** wintercress Bittercress, hairy Bitterweed Black-eyed Susan Black medic Black mustard Blackseed plantain Blessed thistle Bloodflower milkweed Blue lettuce Blue vervain Bracted plantain Bristly oxtongue Broadleaf dock Broadleaf plantain Broomweed Buckhorn Buckhorn plantain **Bulbous** buttercup **Bull thistle** Bullnettle Burclover Burdock Burning nettle Burweed Buttercup Buttonweed Canada thistle Carolina geranium Carpetweed Catchweed bedstraw Catnip Catsear Chickweed, common Chickweed. mouseear Chicory Cinquefoil Clover, crimson Clover, hop Clover, red Clover, strawberry

Clover, white Cockle Cocklebur Common mullein Creeping jenny . Cudweed Curly dock Daisy, English Daisy fleabane Daisy, oxeye Dandelion Dichondra Dogfennel Dollarweed False dandelion Falseflax False sunflower Fiddleneck Florida pusley Frenchweed Galinsoga Goldenrod Ground ivv Gumweed Hairy fleabane Hawkweed Healall Heartleaf drymary Heath aster Hedge bindweed Hedge mustard Henbit Hoary cress Hoary plantain Hoary vervain Horsenettle Jimsonweed Knawel Knotweed Kochia Lambsquarters Lespedeza Mallow Matchweed Mexicanweed Milk vetch Morningglory Mouseear hawkweed Mugwort

Musk thistle

Mustard Narrowleaf plantain Narrowleaf vetch Nettle Orange hawkweed Oriental cocklebur Oxalis Parsley-piert Parsnip Pearlwort Pennycress Pepperweed Pigweed Pineywoods bedstraw Plains coreopsis Plantain Poison ivy Poison oak Pokeweed Poorioe Prairie sunflower Prickly lettuce Prickly sida Prostrate knotweed Prostrate pigweed Prostrate spurge Prostrate vervain Puncturevine Purslane, common Ragweed Red sorrel Redroot pigweed Redstem filaree Rough cinquefoil Rough fleabane Russian pigweed **Russian** thistle Scarlet pimpernel Scotch thistle Sheep sorrel Shepherdspurse Slender plantain Smallflower galinsoga Smooth dock Smooth pigweed Sorrel Sowthistle Spanishneedles

- WEEDS -

Spiny cocklebur Spiny sowthistle Spotted catsear Spotted spurge Spurweed Stinging nettle Strawberry, India mock Tall nettle Tall vervain Tansy ragwort Tansy mustard Tanweed Thistle Trailing crownvetch Tumble mustard Tumble pigweed Velvetleaf Venice mallow Virginia buttonweed Virginia creeper Virginia pepperweed Wavyleaf bullthistle Western clematis Western salsify White mustard Wild aster Wild buckwheat Wild carrot Wild four-o'clock Wild garlic Wild geranium Wild lettuce Wild marigold Wild mustard Wild onion Wild parsnip Wild radish Wild rape Wild strawberry Wild sweet potato Wild vetch Woodsorrel Woolly croton Woolly morningglory Woolly plantain Wormseed Yarrow Yellow rocket Yellowflower pepperweed and other broadleaf weeds

Clover, sweet

Speedwell

Spiny amaranth

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

I. Guarantee of customer satisfaction to be offered by supplemental registrants:

• The Sta-Green[®] Promise.

YOUR SATISFACTION [100%] GUARANTEED OR YOUR MONEY BACK. If you apply <u>Brand</u> <u>Name</u> as directed and are not completely satisfied, just send a proof of purchase from this package and a cash register receipt to the address below. We will promptly send your money back [refund your money].

LOOK

2.5

For other quality Sta-Green[®] Lawn and Garden products. For more information about Sta-Green[®] products, visit us on the Internet at <u>www.fertilizer.com</u> or give our customer service a call at 1-800-874-8892.

II. Advertising claims that may be used: These claims may be presented on the retail container label or on the labeling accompanying the product.

- Ten pounds covers up to 5,000 square feet.
- Twenty pounds covers up to 10,000 square feet.
- Controls dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf weeds including those listed on the back panel.
- Covers 5,000 square feet.
- America's Best Value For All Your Lawn & Garden Needs, Vigoro Lawn Weed Control.
- Try Brand Name Premium Garden Products
- Kills (Controls) these weeds & 150 others
- Visit us on the internet at www.XXXXXXX or call our customer service Hotline at (phone number).
- Controls dandelions and 200 other broadleaf weeds.
- Will not burn your lawn.

EPA Reg. No. 2217-640

- Apply anytime weeds are growing—Spring, Summer or Fall.
- Proven Trimec[®] Lawn Weed Control.
- Apply once in Spring when weeds are growing

