Please read instructions on i	reverse hefore completii	no form	1910	9006 Form A	noroved.	OMB No. 207	n-nn60.	Approval Expires 2-28-95
⊕ EPA	Environme	United States ntal Protection shington, DC 2046	~			Registration Amendme Other	n	OPP Identifier Number
,		Application fo	r Pesti	cide - Sec	ction	<u> </u>		·
1. Company/Product Num	nber 2217-640		2. E	PA Product M Joann	lanager ne I. Mil		3. Pro	oposed Classification
4. Company/Product (Nar Gordon's	ne) Trimec [®] Weeder Gr	anules	PM#	Product Mar	nager-	Team 23	X	None Restricted
5. Name and Address of A PBI/Gordon Corpora Post Office Box 0140 Kansas City, Missour Check if the	tion 090		(b)(i) EPA Prod	, my product Reg. No duct Name	is simila	ar or identical i	n comp	FRA Section 3(c)(3) position and labeling to:
			ection		ا ماماله	<u> </u>	1	NOTIFICATION
Amendment - Explain Resubmission in resp Notification - Explain I	onse to Agency letter da	ated	[_	Agency lett "Me Too" A Other - Exp	ter dated application	n.		JUN 1 9 2006
Three advertising An advisory regal An advisory state These are all presented a This notification is consistent labeling or Confidential State the EPA. I further understan-	arding the timing of ap ement to show the and is highlighted text in the twith the provisions of P ement of Formula of this d that if this notification is	nount of nitrogen p he Appendix (Page PR Notice 98-10 and I product. I understand is not consistent with nd penalties under Se	rovided wile 7) of the EPA regular that it is a the terms of the terms	hen the produ enclosed labe tions at 40 <u>CFF</u> violation of 18 of <u>PR Notice 98</u> nd 14 of FIFRA	uct is ap eling. 3 152.46 U.S.C S 3-10 and	plied as direct , and no other c ec. 1001 to willf	hanges uliv mak	e any false statement to
Material This Product Will	Be Packaged In:		<u> </u>	<u>'!!</u>				
Child-Resistant Packaging Yes* No * Certification must be submitted	Unit Packaging Yes No If "Yes" Unit Packaging wgt.	No. per	Yes No Yes" ckage wgt.	No. per container		2. Type of Conta		
Location of Net Contents In Label Cont Cont	nformation tainer	4. Size(s) Retail Cor	ntainer			ation of Label Di n Label n Labeling acco	rections	
6. Manner in Which Label is A	Affixed to Product	Lithograph Paper glued Stenciled		Other				g product
			ection -					
1. Contact Point (Complete its Name James L. F	ems directly below for id Kunstman, Ph.D.	Title		ntacted, if neco	-	Tele	phone N	n.) No. (Include Area Code) 16-460-6292
I certify that the stateme I acknowledge that any I both under applicable la	knowingly false or mislea	ading statement may	be punisha				3.	Date Application Received (Stamped)
	150		rector of l	Regulatory \$	Service	s 		
4. Typed N ame James L. K	unstman, Ph.D.	5. Date		ne 5, 2006				
EPA Form 8570-1 (Rev. 3-94)	Previous editions are o	bsolete.		Whi	te - EPA	File Copy (ori	ginal) Y	fellow - Applicant Copy

GORDON'S TRIMEC® WEEDER GRANULES

ACTIVE INGREDIENTS:

2,4-dichlorophenoxyacetic acid	1.37%
(+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid	0.31%
Dicamba: 3,6-dichloro-o-anisic acid	0.13%
INERT INGREDIENTS:	98.19%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

27.4 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.

6.2 lbs. (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid equivalent per ton.

2.6 lbs. 3,6-dichloro-o-anisic acid equivalent per ton.

Isomer Specific by AOAC Methods

Contains the single isomer form of Mecoprop-p.

TRIMEC[®] is a registered trademark of PBI/Gordon Corporation.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for Precautionary Statements.

NET WEIGHT 10 OR 20 POUNDS

650/ AP060506 EPA REG. NO. 2217-640 EPA Est. No. 2217-KS-1 MANUFACTURED BY:

An Employer-Owned Company
1217 West 12th Circuit
Kennes City, Massoul 84101

Telephone: 1-800-821-7925



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.



PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hand and exposed skin with soap and water.

First Aid

If in eyes:	 Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing:	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.
If swallowed:	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.

ENVIRONMENTAL HAZARDS:

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater. Do not contaminate domestic or irrigation waters. Do not apply when weather conditions favor drift from target area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until dust has settled.

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs. **DISPOSAL:**

If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partially filled: Call your local solid waste agency or toll free number (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.

USE PRECAUTIONS:

Do not allow the material to drift onto flowers, vegetables or ornamental shrubbery. Apply when the air is calm. Do not apply to newly seeded lawns. Do not apply in areas underlaid by roots of desirable trees and shrubs. Do not exceed specified dosages for any area; be particularly careful within dripline of tree and other ornamental species. During application, close spreader openings when turning. Do not contaminate domestic or irrigation waters.

INSTRUCTIONS:

Gordon's Trimec[®] Weeder Granules controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days.



For use on established turf of warm season grasses includes bermudagrass, centipedegrass, and zoysiagrass. Established turf of cool season grasses includes Kentucky bluegrass, perennial ryegrass, tall fescue, and turf-type tall fescues. Do not apply this product on carpetgrass, dichondra, St. Augustinegrass, or established turf containing desirable clovers.

The maximum application rate to turf is 1.2 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

- 1. Mow lawn to normal height 1 to 2 days before application.
- 2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
- Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants.
- 4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

Option 1: Coverage and Spreader Settings

Spreader Settings	
Scott	5½
Sears	6

The above spreader settings are approximate. Calibrate your spreader on 100 square feet and adjust to apply at the rate of 1 pound of product per 500 square feet.

Option 2: Alternate Coverage and Spreader Settings

COVERAGE: To determine how much fertilizer you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct any non-lawn areas (house, garden, etc.). One bag covers 5,000 sq.ft. [Length x Width = Total Sq.Ft. - Non-Lawn Areas = Total Sq.Ft. Coverage You Need]

SPREADER SETTINGS:

IMPORTANT: It is important to follow the directions as indicated on this quality lawn care product as over application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. Be sure to check spreader before beginning, adjusting if necessary. Consult Owner's Manual. *Never apply by hand.*

Spreading Hints: These granular settings are approximate. Calibrate your spreader on 100 sq.ft. and adjust to apply at the rate of 1 lb. of product per 500 sq.ft.. This bag covers _____ sq.ft. of lawn area.

To avoid gaps and streaks, first spread border of lawn and then fill in center with uniform, parallel passes. Walk at a steady pace so as to apply evenly throughout the lawn. [Graphics and/or diagrams may be presented on the container label].

Rotary Spreaders	Setting
Ames	see spreader instructions
EV-N-Spread	13
Ortho	3
Precision	5
Quaker	4
Republic EzSPRAY TM	7 1/4
Republic EZ Grow	3
Scotts Easy Green	27 ¼
Scotts Speedy Green	5
Scotts Speedy Green 1000	6
Scotts Speedy Green 2000	7
Scotts Speedy Green 3000	6
True Temper	4
Vigoro	7 1/4

Drop Spreaders	Setting		
Ames	see spreader instructions		
Bandini	2		
Ortho	6 1/4		
Precision	6 1/2		
Republic EZ	6 ¼		
Scotts AccuGreen	8 3/4		
Scotts Drop	6 1/2		
Scotts Even Green	6 1/2		
Vigoro	2 1/2		



Spreader Settings

The spreader settings below are approximate. Calibrate your spreader on 100 sq. ft. and adjust to apply at the rate of 1 pound of product per 500 sq. ft. (2 lbs. of material per 1,000 sq. ft.). Rate of walk, moisture and spreader condition can vary the spreader setting necessary to apply the correct amount.



Apply in a single pass at a rate of 2 lbs, per 1,000 sq. ft. Start on the outside border and apply in a circular path working toward the center. Overlap the coverage slightly, and shut off spreader before making hard turns.

Broadcast Spreaders	Setun
Agn-Feb Rotary	5
Cyclone' Relaty	4
Earthway* Rolary	16
Greenwew" Jel-Spred* SS1	18
Eesto: Rotary 'dial A-Symmotor dial	6.15
Red Dever Hotary (1-20)	5
Republic Rotary	5
Scotts' EasyGreen	26
Scotts* EdgeGuard*	5
Scotts" SpeedyGreen" 2000 3000	5
Sears: Crafteman Hotary	5
Sta-Green: Rotary (1-10)	3
True Temper* Rotary	4
Vigoro* Ratary (1-10)	3





Apply in a single pass at a rate of 2 lbs. per 1,000 sq. lt. Start on the outside border and apply in a rectangular path working back and forth across the length of the lawn. Overlap the coverage slightly, and shut off spreader before making hard turns.

Drop Spreaders	Setting	
Agn-Fatt Drop	4	/_#
Greenway Drop-Spred' XP1	15	- F A
Red Devs Drop (1-5)	4	_ / //
Repubse* Drop	512	
Scotts* Oron Accudingen (2-18)	5%. ◢	
Social Drop PF1 (2-18)	7',1	
Sta-Green* Drop (1-5)	4	

If your spreader model does not appear above, go to our website at www.preen.com, or call toll-free (800) 233-1067, 8 A.M. - 5 P.M. EDT. Monday - Friday, for additional model settings.

[Note: Additional spreader models and settings may be added or substituted by the supplemental registrant provided they deliver the labeled rate of product.]

[Note: Spreader settings may be varied by the supplemental registrant as different grades of fertilizer are used, provided they deliver the labeled rate of pesticide.]



	-	WEEDS -	
Annual yellow sweetclover	Clover, white	Mustard	Spiny cocklebur
Aster	Cockle	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Cocklebur	Narrowleaf vetch	Spotted catsear
Bedstraw	Common mullein	Nettle	Spotted spurge
Beggarticks	Creeping jenny	Orange hawkweed	Spurweed
Betony, Florida	Cudweed	Oriental cocklebur	Stinging nettle
Bindweed, field	Curly dock	Oxalis	Strawberry, India mock
Bird vetch	Daisy, English	Parsley-piert	Tall nettle
Bitter wintercress	Daisy fleabane	Parsnip	Tall vervain
Bittercress, hairy	Daisy, oxeye	Pearlwort	Tansy ragwort
Bitterweed	Dandelion	Pennycress	Tansy mustard
Black-eyed Susan	Dichondra	Pepperweed	Tanweed
Black medic	Dogbane	Pigweed	Thistle
Black mustard	Dogfennel	Pineywoods bedstraw	Trailing crownvetch
Blackseed plantain	Dollarweed	Plains coreopsis	Tumble mustard
Blessed thistle	False dandelion	Plantain	Tumble pigweed
Bloodflower milkweed	Falseflax	Poíson ivy	Velvetleaf
Blue lettuce	False sunflower	Poison oak	Venice mallow
Blue vervain	Fiddleneck	Pokeweed	Virginia buttonweed
Bracted plantain	Florida pusley	Poorjoe	Virginia creeper
Brassbuttons	Frenchweed	Prairie sunflower	Virginia pepperweed
Bristly oxtongue	Galinsoga	Prickly lettuce	Wavyleaf bullthistle
Broadleaf dock	Goldenrod	Prickly sida	Western clematis
Broadleaf plantain	Ground ivy	Prostrate knotweed	Western salsify
Broomweed	Gumweed	Prostrate pigweed	White mustard
Buckhorn	Hairy fleabane	Prostrate spurge	Wild aster
Buckhorn plantain	Hawkweed	Prostrate vervain	Wild buckwheat
Bulbous buttercup	Healall	Puncturevine	Wild carrot
Bull thistle	Heartleaf drymary	Purslane, common	Wild four-o'clock
Bullnettle	Heath aster	Ragweed	Wild garlic
Burclover	Hedge bindweed	Red sorrel	Wild geranium
Burdock	Hedge mustard	Redroot pigweed	Wild lettuce
Burning nettle	Hemp	Redstem filaree	Wild marigold
Burweed	Henbit	Rough cinquefoil	Wild mustard
Buttercup	Hoary cress	Rough fleabane	Wild onion
Buttonweed	Hoary plantain	Russian pigweed	Wild parsnip
Canada thistle	Hoary vervain	Russian thistle	Wild radish
Carolina geranium	Horsenettle	Scarlet pimpernel	Wild rape
Carpetweed	Jimsonweed	Scotch thistle	Wild strawberry
Catchweed bedstraw	Knawel	Sheep sorrel	Wild sweet potato
Catnip	Knotweed	Shepherdspurse	Wild vetch
Catsear	Kochia	Slender plantain	Woodsorrel
Chickweed, common	Lambsquarters	Smallflower galinsoga	Woolly croton
Chickweed, mouseear	Lespedeza	Smooth dock	Woolly morningglory
Chicory	Mallow	Smooth pigweed	Woolly plantain
Cinquefoil	Matchweed	Sorrel	Wormseed
Clover, crimson	Mexicanweed	Sowthistle	Yarrow
Clover, hop	Milk vetch	Spanishneedles	Yellow rocket
Clover, red	Morningglory	Speedwell	Yellowflower pepperweed
Clover, strawberry	Mouseear hawkweed	Spiny amaranth	and other broudleaf weeds
Clover, sweet	Mugwort		
	Musk thistle		

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

8/8

APPENDIX

- I. Guarantee of customer satisfaction to be offered by supplemental registrants:
 - The Sta-Green[®] Promise.

YOUR SATISFACTION [100%] GUARANTEED OR YOUR MONEY BACK. If you apply <u>Brand Name</u> as directed and are not completely satisfied, just send a proof of purchase from this package and a cash register receipt to the address below. We will promptly send your money back [refund your money].

LOOK

For other quality Sta-Green[®] Lawn and Garden products. For more information about Sta-Green[®] products, visit us on the Internet at www.fertilizer.com or give our customer service a call at 1-800-874-8892.

- II. Advertising claims that may be used: These claims may be presented on the retail container label or on the labeling accompanying the product.
 - Ten pounds covers up to 5,000 square feet.
 - ◆ Twenty pounds covers up to 10,000 square feet.
 - Controls dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf weeds including those listed on the back panel.
 - ♦ Covers 5,000 square feet.
 - America's Best Value For All Your Lawn & Garden Needs, Vigoro Lawn Weed Control.
 - ◆ Try Brand Name Premium Garden Products
 - ♦ Kills (Controls) these weeds & 150 others
 - Visit us on the internet at www.XXXXXXX or call our customer service Hotline at (phone number).
 - Controls dandelions and 200 other broadleaf weeds.
 - Will not burn your lawn.
 - Apply anytime weeds are growing—Spring, Summer or Fall.
 - Proven Trimec[®] Lawn Weed Control.
 - Apply once in Spring when weeds are growing
 - · Apply anytime broadleaf weeds are actively growing
 - ◆ Apply in Spring and Fall
 - Enjoy a Beautiful Weed-Free Lawn
 - Special guidelines may apply to the specific timing of fall fertilizer applications to Centipedegrass and St. Augustinegrass lawns. Contact your local Extension Service for information specific to your geographic region.
 - Typically a lawn utilizes 1-4 lb of nitrogen per 1,000 sq ft per growing season. When applied as directed this product provides 0.79 lb of nitrogen per 1,000 sq ft."