

GORDON'S TRIMEC® WEED & FEED 24B 20-10-5

ACTIVE INGREDIENTS:

*2,4-dichlorophenoxyacetic acid.....	0.5471%
**2-(2-methyl-4-chlorophenoxy)propionic acid.....	0.2448%
***Dicamba: 3,6-dichloro-o-anisic acid.....	0.0516%
INERT INGREDIENTS.....	99.1565%


TOTAL	100.0000%
-------	-----------

THIS PRODUCT CONTAINS:

- *10.942 lbs. 2,4-Dichlorophenoxyacetic acid equivalent per ton.
 - **4.896 lbs. 2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton.
 - ***1.032 lbs. 3,6-dichloro-o-anisic acid equivalent per ton.
- Isomer Specific by AOAC Methods
TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

KEEP OUT OF REACH OF CHILDREN

CAUTION




Statement of Practical Treatment

In case of contact, wash skin with plenty of soap and water.
For eyes, flush with water for 15 minutes and get medical attention.
See back panel for additional Precautionary Statements.

NET WEIGHT (16, 18, 20, 22, 25, 40, or 54) POUNDS

560/ APXXXXXX
EPA REG. NO. 2217-643
EPA EST. NO. 2217-KS-X
MANUFACTURED BY:



2 4 5



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

ENVIRONMENTAL HAZARDS: Drift or runoff may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from target area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.

PESTICIDE DISPOSAL: Securely wrap bag in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.


USE PRECAUTIONS: Do not apply this product to dichondra or carpetgrass. Do not apply this product to 'Floritam' St. Augustinegrass in Florida.

Do not use this product where desirable clovers are present or allow the material to wind drift onto flowers, vegetables or ornamental shrubbery. Do not apply in areas underlaid by roots of desirable trees or shrubs. Do not exceed specified dosages for any area; be particularly careful within dripline of tree and other ornamental species. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning to prevent over-application.

INSTRUCTIONS: Gordon's Trimec® Weed & Feed 24B controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantains, henbit, spurge and many others commonly found in home lawns. In addition, a 20-10-5 fertilizer provides plant nutrients or a generous supply of plant food. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days.

The maximum application rate to turf is 4.0 ~~1.2~~ pounds/2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.



Spreader Settings	
Scott.....	6
Lawn Beauty.....	5
Jackson.....	7½
Sears.....	4

The above spreader settings are approximate.

- 16 lb. bag: Calibrate your spreader on 310 sq.ft. and adjust to apply 1.0 lb. of product per 310 sq.ft.
- 18 lb. bag: Calibrate your spreader on 275 sq.ft. and adjust to apply 1.0 lb. of product per 275 sq.ft.
- 20 lb. bag: Calibrate your spreader on 250 sq.ft. and adjust to apply 1.0 lb. of product per 250 sq.ft.
- 22 lb. bag: Calibrate your spreader on 227 sq.ft. and adjust to apply 1.0 lb. of product per 227 sq.ft.
- 25 lb. bag: Calibrate your spreader on 200 sq.ft. and adjust to apply 1.0 lb. of product per 200 sq.ft.
- 40 lb. bag: Calibrate your spreader on 250 sq.ft. and adjust to apply 1.0 lb. of product per 250 sq.ft.
- 54 lb. bag: Calibrate your spreader on 275 sq.ft. and adjust to apply 1.0 lb. of product per 275 sq.ft.


- WEEDS -			
Bedstraw	Ground Ivy	Peppergrass	Spurge
Black Medic	Healall	Pigweed	Thistle
Buckhorn	Henbit	Plantains	Wild Carrot
Burdock	Knotweed	Poison Ivy	Wild Garlic
Chicory	Lambsquarters	Poison Oak	Wild Lettuce
Chickweed	Lespedeza	Purslane	Wild Onion
Clover	Mallow	Ragweed	Yarrow
Dandelion	Momingglory	Sheep Sorrel	
Dock	Oxalis	Speedwell	

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.



495

APPENDIX

I. Container Packaging and Size:

Note to Supplemental Registrants: Select the appropriate bag size and present only the information provided for that bag size on your product labeling.

Bag Size, Pounds	Calibration Rate, Pounds/Sq.Ft.	Application Rate, Pounds/1000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.	
			Cool Season Grass	St. Augustinegrass
16	1/310	3.23	5,000	5,000
18	1/275	3.64	5,000	5,000
20	1/250	4.00	5,000	-----
22	1/227	4.40	5,000	-----
25	1/200	5.00	5,000	-----
40	1/250	4.00	10,000	-----
54	1/275	3.64	15,000	-----

II. Previously Approved Fertilizer Grades:

Fertilizer grades with nutrient contents expressed as the percentages of N, P₂O₅, and K₂O include the following:

15-5-7	18-4-6	20-10-5	23-4-5	26-3-3	29-3-4
15-5-8	18-6-12	22-3-16	25-3-3	26-5-3	30-2-4
16-3-5	20-4-4	22-4-4	25-3-5	28-2-14	31-4-4
16-6-8	20-4-10	23-3-3	25-6-4	28-3-3	

The fertilizer grade guaranteed analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.

III. Advertising claims appropriate for the bag size:

- ◆ For fertilizers containing iron, supplemental registrants may add one of the following two statements:
 - a) "This product may stain concrete and stone, where spilling occurs, sweep off immediately, then flush with water."
 - b) Sweep all residue from concrete surfaces immediately after application. The iron content in this product will stain sidewalks, patios, driveways, swimming pools and deck coatings if not removed before watering. Seller nor distributor assumes any responsibility for damage incurred to any surface other than product's intended use.
- ◆ ___ pounds of product cover up to ___ square feet of turf
- ◆ Controls Dandelion, Chickweed, Knotweed, Plantains, Henbit, Spurge and many other broadleaf weeds; including those listed on the back panel.
- ◆ Contains Polyon® Controlled-Release Nitrogen.
- ◆ Greens 3X Longer.
- ◆ Premium Quality
- ◆ What makes Pursell's Polyon® controlled-release nitrogen so unique? Each granule of Polyon® controlled-release nitrogen is encapsulated by our exclusive ultra-thin polymer coating which steadily releases nitrogen into your lawn as your grass needs it. This means your lawn receives a gradual and precise amount of nitrogen. Polyon® controlled-release nitrogen releases in response to temperature rather than moisture so when used as directed, there is virtually no risk of burning or surge growth following heavy rainfall. Your lawn steadily becomes thicker, sturdier and greener.
- ◆ Feeds and greens lawns quickly.
- ◆ Three types of nitrogen for long-lasting results.
- ◆ Satisfaction Guaranteed!
- ◆ Greens Quick, Stays Green.

- ◆ Promotes Longer-Lasting Winter Hardiness & Early Spring Green-Up.
- ◆ This product contains a highly effective weed killer and a winter lawn conditioner.
- ◆ Controls a wide spectrum of broadleaf weeds.
- ◆ Provides controlled-release feeding and promotes longer-lasting winter hardiness and drought tolerance.
- ◆ A _____ (Grade ex. 22-3-16) fertilizer helps condition lawns to prevent winter injury.
- ◆ A lawn typically utilizes 1 to 4 lbs. of Nitrogen per 1,000 sq.ft. per growing season. When applied as directed, this product will supplement your lawn maintenance program by providing ___ lbs. of nitrogen per 1,000 sq.ft.
- ◆ Kills Over 30 Varieties of Weeds While Feeding Your Lawn!
- ◆ Premier formula for weed and feed.
- ◆ A balanced formula for overall lawn health.
- ◆ Long lasting with 2% iron.
- ◆ Controls (kills) dandelions, plantains, and other weeds.
- ◆ Feeds the lawn with two types of nitrogen.
- ◆ Easy application with any type of spreader.
- ◆ Combines a highly effective lawn weed killer and winter lawn conditioner.
- ◆ Always apply with a properly calibrated rotary or drop type fertilizer spreader to distribute the granules evenly. Never apply by hand.
- ◆ Place your spreader on the driveway or other paved surface when filling. This guards against spilling fertilizer on grass. For even coverage, begin walking before you turn spreader to the open position to avoid dropping fertilizer in one spot when the spreader is opened. Be sure the spreader is closed when filling it, when turning to backtrack, or when stopping on the lawn.
- ◆ To avoid gaps and streaks, first spread the border and fill in center with uniform, parallel passes.
- ◆ Equipment settings may vary and may be presented on the container labeling of supplemental registrations.
- ◆ [Optional Spreader Listing]

Rotary Spreaders	Ortho Rotary	Ortho Drop
Republic EZ	True Temper	Scotts Drop (PF)
EV N Spread	Scotts Speedy Green	Scotts Even Green
Scotts Easy Green	Republic EZ Grow	Garden Plus
Precision	Garden Plus	Scotts AccuGreen
Quaker	Drop Spreaders	

