

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXI, SUBSTANCES り

FEB 17 2009

James L. Kunstman, Ph. D. Product Registration PBI/Gordon, Corporation P.O. Box 014090 Kansas City, Missouri 64101

 SUBJECT: Application for Pesticide Notification (PRN 98-10 and 2008-1) Request General Label Change(s) Additional Marketing Claims & Table Rate Change, Directions for Use and Environmental Hazards Statement EPA Reg. No. 2217-643 Application Dated January 6, 2008

Dear Registrant:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 and 2008-1 dated 01/06/09 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

Linda Arrington Notifications & Minor Formulations Team Leader Registration Division (7505P) Office of Pesticide Programs

F structions on re	verse before completin	g form.	Form	Approve	d. OMB No. 207	70-0060. Approval Expires 2-28-95
€ЕРА		United States ntal Protection shington, DC 20460	Agency] Registratio	
	4	pplication for	Pesticide - Se	ection	1	
1. Company/Product Num			2. EPA Product		er	3. Proposed Classification
4. Company/Product (Nam Gordon's Tri		eed 24B	PM# Product Ma		· · · ·	None Restricted
5. Name and Address of A	pplicant (Include ZIP	? Code)	6. Expedited R	eview.	In accordance	with FIFRA Section 3(c)(3)
PBI/Gordon Corporati Post Office Box 0140			(b)(i), my produc EPA Reg. No.	ct is simi		in and labeling to:
Kansas City, Missour	64101		Product Name		FEB 17	2009
	s is a new address		tion II		<u></u>	
		500	tion – II			· · · · · · · · · · · · · · · · · · ·
Amendment - Explain below. Resubmission in response to Agency letter dated Notification - Explain below.			Agency I	Final printed labels in response to Agency letter dated "Me Too" Application.		
 Clarify spread Add one fertili Add one adve Add one adve This notification is consistent labeling or Confidential State 	er patterns (single p. zer grade of 23-0-7 / rtising claim <i>[end of</i> with the provisions of <u>F</u> ment of Formula of this I that if this notification	ass, double pass, crò [page 6] Appendix, page 8]. 2 <u>R Notice 98-10</u> and EP product. I understand th is not consistent with th	oss hatching). <i>[pag</i> PA regulations at 40 <u>C</u> hat it is a violation of he terms of <u>PR Notice</u>	FR 152.4 18 U.S.C 98-10 ar	16, and no other Sec. 1001 to wi	Notice (PRN) 2008-1. [page 2] changes have been made to the lifully make any false statement to 6, this product may be in violation of
FIFRA and I may be subject t	o enforcement action a			HA.		
1. Material This Product Will	Be Packaged In:					
Child-Resistant Packaging Yes*	Unit Packaging Yes No	Wate	er Soluble Packaging Yes No		2. Type of Cor Metal Plastic	itainer .
* Certification must be submitted	If "Yes" Unit Packaging wgt.		If "Yes" No. per Package wgt. container Image: Container Other (Specify)		Specify)	
3. Location of Net Contents In		4. Size(s) Retail Conta	ainer	5. Lo	ocation of Label On Label	
6. Manner in Which Label is A	fixed to Product	Lithograph Paper glued	Oth	ner	•	
		Stenciled			•	
			tion – IV			
1. Contact Point (Complete ite	ems directly below for it		I to be contacted, if n	ecessary		
Name James L. K	Kunstman, Ph.D.		irector of Regulat	ory Ser		elephune No. (Inc!ude Area Code) 816-460-6292
I certify that the stateme I acknowledge that any I both under applicable la	knowingly false or misle	eading statement may b				6. Date Application Received (Stamped)
2. Signature	Ç., E		ector of Regulator	y Servi	ces	
4. Typed Name James L. K	unstman, Ph.D.	5. Date	January 6, 20	009		
EPA Form 8570-1 (Rev. 3-94) Previous editions are	obsolete.	. 1	White - E	PA File Copy (original) Yellow - Applicant Copy

Certification of label change relative to PR Notice 2008-1

Relative to the changes made to the label pertaining to the environmental hazards section of this label, the registrant certifies the following:

"Notification of label change relative to PR Notice 2008-1. This notification is consistent with the guidance in PR Notice 2008-1 and the requirements of EPA's regulations at 40 CFR 156.10, 40 CFR 156.80 and 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the guidance of PR Notice 2008-1 and the requirements of 40 CFR 156.10, 40 CFR 156.80 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA."

James L. Kunstman, PhD Director, Regulatory Services PBI/Gordon Corporation

10 - Feb - 2009

Date

Submitter

Certification with Respect to Label Integrity

I certify that the information (including, but not limited to, text, tables, and graphics) contained in the electronic file identified below by file name and submitted with this certification is the same information as that on the paper copies of these documents included with this submission.

PROPOSED LABEL					
EPA Registration #	Date Submitted to EPA	Electronic file name			
2217-643	1/6/09	002217-00643.20081223			

I certify that the statements that I have made on this form are true, accurate, and complete. I acknowledge that any knowingly false or misleading statements may be punishable by fine or imprisonment or both under applicable law.

Signature

James L. Kunstman, Ph.D. Name (typed)

Director, Regulatory Services Title 01/06/2009

Date

GORDON'S TRIMEC[®] WEED & FEED 24B 20-10-5

ACTIVE INGREDIENTS:

2,4-dichlorophenoxyacetic acid	0.55%
(+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid	0.12%
Dicamba: 3,6-dichloro-o-anisic acid	0.05%
INERT INGREDIENTS:	<u>99.28%</u>
TOTAL	100.00%

THIS PRODUCT CONTAINS:	
11.0 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.	
2.4 lbs. (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid equivalent per ton.	
1.0 lbs. 3,6-dichloro-o-anisic acid equivalent per ton.	NOTHICATION
Isomer Specific by AOAC Methods	
Contains the single isomer form of Mecoprop-p.	
TRIMEC [®] is a registered trademark of PBI/GORDON CORPORATION.	FEB 1 7 2009

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for Precautionary Statements.

NET WEIGHT (16, 17, 18, 20, 22, 25, 40, 50 or 54) POUNDS

560/ AP122308 EPA Reg. No. 2217-643 EPA Est. No. 2217-KS-1 MANUFACTURED BY:

An Employto-Owned Company 1217 West 12th Street Kanasa City, Mesouri 64101 Telephone: 1-800-821-7925

Page 1 of 8 EPA Reg. No. 2217-643 002217-00643 20081223



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

First Aid

	ter en la construction de la construction d
If in eyes:	 Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing:	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.
If swallowed:	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
lf inhaled:	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.
	ct container or label with you when calling a poison control center or doctor or going for additional information in case of emergency, call toll free 1-877-800-5556.

ENVIRONMENTAL HAZARDS:

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater. Do not apply when weather conditions favor drift from target area.

Environmental Hazards (alternate per PRN 2008-1)

Drift or runoff may adversely affect nontarget plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street; back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems!

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Jo not enter treatment area until spray has dried or dust has settled.

Page 2 of 8

EPA Reg. No. 2217-643 002217-00643.20081223

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs. **DISPOSAL:**

If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partially filled: Call your local solid waste agency or toll free number (1-800-XXX-XXX) for disposal instructions. Never place unused product down any indoor or outdoor drain.

USE PRECAUTIONS:

Do not apply this product to dichondra or carpetgrass. Do not apply this product to 'Floratam' St. Augustinegrass in Florida.

Do not use this product where desirable clovers are present or allow the material to drift onto flowers, vegetables or ornamental shrubbery. Do not apply in areas underlaid by roots of desirable trees or shrubs. Do not exceed specified dosages for any area; be particularly careful within dripline of tree and other ornamental species. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning to prevent over-application.

INSTRUCTIONS:

Gordon's Trimec[®] Weed & Feed 24B controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a 20-10-5 fertilizer provides plant nutrients or a supply of plant food. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days.

For state registrations in Kentucky, add the following paragraph:

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing (*Number of pounds) of nitrogen per 1,000 square feet. Please consult your local extension office for questions regarding a regular fertilizer program suitable for your geographic region.

Fertilizer Labeling Requirements:

- All fertilizer products distributed in Washington at a minimum, must have the following labeling statement: "This product has been registered with the Washington State Department of Agriculture. When applied as directed, this fertilizer meets the Washington standards for arsenic, cadmium, cobalt, mercury, molybdenum, lead, nickel, selenium, and zinc. You have the right to receive specific information about Washington standards from the distributor of this product."
- <u>After July 1, 1999</u>, the label must also state: "Information received by the Washington State Department of Agriculture regarding the components in this product is available on the internet at http://www.wa.gov/agr/."
- "OR" Information regarding the contents and levels of metals in this product is available on the Internet at <u>http://www.aapfco.org/metal.htm</u>.

The maximum application rate to turf is 1.2 pounds of 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

- 1. Mow lawn to normal height 1 to 2 days before application.
- 2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
- 3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants.
- 4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

Page 3 of 8 0081223

EPA Reg. No. 2217-643 002217-00643 20081223

Spreader Settings	
Scott	6
Lawn Beauty	5
Jackson	71⁄2
Sears	4

As a guide, set spreader at the setting indicated by the above chart before applying. Remember the settings provided are only a guide and depend on the age, condition and speed of operation of your spreader. Check spreader calibration often and make adjustments as needed. Keep this product off sidewalks, patios, driveways, etc. This product contains materials which will stain concrete and painted surfaces. If it comes in contact with a concrete surface, sweep off immediately before finishing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter product beyond lawn edge and onto concrete.

Calibration Instructions: To provide proper distribution, calibrate spreader before application.

The above spreader settings are approximate.

- 16 lb. bag: Calibrate your spreader on 310 sq.ft. and adjust to apply 1.0 lb. of product per 310 sq.ft.
- 17 lb. bag: Calibrate your spreader on 300 sq.ft. and adjust to apply 1.0 lb. of product per 300 sq.ft.
- 18 lb. bag: Calibrate your spreader on 275 sq.ft. and adjust to apply 1.0 lb. of product per 275 sq.ft.
- 20 lb. bag: Calibrate your spreader on 250 sq.ft. and adjust to apply 1.0 lb. of product per 250 sq.ft.
- 22 lb. bag: Calibrate your spreader on 227 sq.ft. and adjust to apply 1.0 lb. of product per 227 sq.ft.
- 25 lb. bag: Calibrate your spreader on 200 sq.ft. and adjust to apply 1.0 lb. of product per 200 sq.ft.
- 40 lb. bag: Calibrate your spreader on 250 sq.ft. and adjust to apply 1.0 lb. of product per 250 sq.ft.
- 50 lb. bag: Calibrate your spreader on 300 sq.ft. and adjust to apply 1.0 lb. of product per 300 sq.ft.
- 54 lb. bag: Calibrate your spreader on 275 sq.ft. and adjust to apply 1.0 lb. of product per 275 sq.ft.

[Optional

Choose One Of These Spreader Patterns

- 1 Single Pass Operate the equipment at intervals corresponding to the width or swath of the spreader. For drop spreaders, overlap the wheel tracks to ensure complete coverage. Close the spreader openings when turning to change directions.
- 2 Double Pass Reduce the application rate by one-half of the recommended rate, calibrate the spreader, and adjust the settings for the desired application rate. Operate the spreader over the lawn two (2) times. Close the spreader openings when turning to change directions.
- 3: Cross Hatching Reduce the application rate by one-half of the recommended rate calibrate the spreader, and adjust the settings for the desired application rate. Operate the spreader over the lawn two (2) times. For the second trip, operate the spreader perpendicular to the wheel tracks of the spreader pattern of the first trip. Close the spreader opening when turning to change directions.]

- WEEDS -					
Bedstraw	Ground ivy	Peppergrass	Spurge		
Black medic	Healall	Pigweed	Thistle		
Buckhorn	Henbit	Plantain	Wild carrot		
Burdock	Knotweed	Poison ivy	Wild garlic		
Chicory	Lambsquarters	Poison oak	Wild lett-ice		
Chickweed	Lespedeza	Purslane	Wild onion		
Clover	Mallow	Ragweed	Yarrow		
Dandelion	Morningglory	Sheep sorrel			
Dock	Oxalis	Speedwell			

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

I. Container Packaging and Size:

Note to Supplemental Registrants: Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, equipment settings of drop spreaders, rotary spreaders, and other granular applicators (pictures or drawings) may be presented on the container labels.

Bag Size, Calibration Rate,		Application Rate,	Treated Area for This Bag Size, Sq.Ft.		
Pounds	Pounds/Sq.Ft.	Pounds/1000 Sq.Ft.	Cool Season Grass	St. Augustinegrass	
160.00	1/310	3.23	5,000	5,000	
170.00	1/300	3.33	5,100	5,100	
18	1/275	3.64	5,000	5,000	
20	1/250	4.00	5,000		
22	1/227	4.40	5,000		
25	1/200	5.00	5,000		
40	1/250	4.00	10,000		
50	1/300	3.33	15,000		
54	1/275	3.64	15,000		

II. Previously Approved Fertilizer Grades:

Fertilizer grades with nutrient contents expressed as the percentages of N, P₂O₅, and K₂O include the following:

12-5-5	18-4-6	20-4-4	23-0-7	25-3-5	28-2-14
15-5-7	18-5-10	20-4-10	23-3-3	25-4-8	28-3-3
15-5-8	18-6-12	20-10-5	23-3-7	25-6-4	29-3-4
16-3-5	18-8-0	22-3-16	23-4-5	26-3-3	30-2-4
16-6-8	19-4-8	22-4-4	25-3-3	26-5-3	31-4-4

The fertilizer grade guaranteed analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.

Ill. Advertising claims appropriate for the bag size:

- Prepares your lawn for winter dormancy.
- The key to a beautiful lawn next spring.
- Promotes deeper green lawns and early rapid growth.
- For long lasting results.
- Controls (Kills) dandelion and other tough broadleaf weeds.
- Advisories for fertilizers containing iron:
 - a) "This product may stain concrete and stone, where spilling occurs, sweep off immediately, then flush with water."
 - b) Sweep all residue from concrete surfaces immediately after application. The iron content in this product will stain sidewalks, patios, driveways, swimming pools and deck coatings if not removed before watering. Seller nor distributor assumes any responsibility for damage incurred to any surface other than product's intended use.
- ____ pounds of product cover up to ____ square feet of turf
- Controls (Kills) (desired weed species from attached listing may be inserted here e.g. dandelicn, chickweed, knotweed, plantain, henbit, spurge) and many other broadleaf weeds; including these listed on the back panel.

Page 6 of 8

EPA Reg. No. 2217-643 002217-00643.20081223

- Contains Polyon[®] controlled-release nitrogen.
- Greens 3X longer.
- Premium quality
- What makes Pursell's Polyon® controlled-release nitrogen so unique? Each granule of Polyon® controlled-release nitrogen is encapsulated by our exclusive ultra-thin polymer coating which steadily releases nitrogen into your lawn as your grass needs it. This means your lawn receives a gradual and precise amount of nitrogen. Polyon® controlled-release nitrogen releases in response to temperature rather than moisture so when used as directed, there is virtually no risk of burning or surge growth following heavy rainfall. Your lawn steadily becomes thicker, sturdier and greener.
- Feeds and greens lawns quickly.
- Three types of nitrogen for long-lasting results.
- Satisfaction guaranteed!
- Greens quick, stays green.
- Promotes longer-lasting winter hardiness & early spring green-up.
- This product contains a highly effective weed killer and a winter lawn conditioner.
- Controls (Kills) a wide spectrum of broadleaf weeds.
- Provides controlled-release feeding and promotes longer-lasting winter hardiness and drought tolerance.
- A (Grade ex. 22-3-16) fertilizer helps condition lawns to prevent winter injury.
- Kills (Controls) over (Number) varieties of weeds while feeding your lawn!
- Premier formula for weed and feed.
- A balanced formula for overall lawn health.
- Long lasting with 2% iron.
- Controls (kills) dandelion, plantain, and other weeds.
- Feeds the lawn with two types of nitrogen.
- Easy application with any type of spreader.
- Combines a highly effective lawn weed killer and winter lawn conditioner.
- Always apply with a properly calibrated rotary or drop type fertilizer spreader to distribute the granules evenly. Never apply by hand.
- Place your spreader on the driveway or other paved surface when filling. This guards against spilling fertilizer on grass. For even coverage, begin walking before you turn spreader to the open position to avoid dropping fertilizer in one spot when the spreader is opened. Be sure the spreader is closed when filling it, when turning to backtrack, or when stopping on the lawn.
- To avoid gaps and streaks, first spread the border and fill in center with uniform, parallel passes.
- [Spreader Listing and Graphics]

Rotary Spreaders	Ortho Rotary	Ortho Drop
Republic EZ	True Temper	Scotts Drop (PF)
Ev N Spread	Scotts Speedy Green	Scotts Even Green
Scotts Easy Green	Republic EZ Grow	Garden Plus
Precision	Garden Plus	Scotts AccuGreen
Quaker	Drop Spreaders	

- Greens up to 3 months.
- Greens rapidly.
- 10% slow release nitrogen.
- 2% iron for dark green turf.
- 1% iron for greener lawns
- ♦ 12% sulfur
- ♦ 1% iron
- ♦ 1% zinc
- Homogeneous pellets
- Non-burning formula
- Pelletized for easy application.
- Contains Trimec[®] herbicide to control (kill) the toughest weeds.
- Controls broadleaf weeds
- Slow-release formula builds healthier, thicker lawns

Page 7 of 8

EPA Reg. No. 2217-643 002217-00643.20081223

- Controls weeds-Late Sping and/or Early Fall—Feeds turf growth
- For a beautiful lawn all year long, choose a GroundWork Feitilizer for every season.
- Controls (Number) weeds! (Listing of weeds may include the following.)

Coverage up to XXXX square feet

Annual yellow sweetclover Aster Austrian fieldcress Bedstraw Beggarticks Betony, Florida Bindweed, field Bird vetch Bitter wintercress Bittercress, hairy Bitterweed Black-eyed Susan Black medic Black mustard Blackseed plantain Blessed thistle Bloodflower milkweed Blue lettuce Blue vervain Bracted plantain Bristly oxtongue Broadleaf dock Broadleaf plantain Broomweed Buckhorn Buckhorn plantain Bulbous buttercup Bull thistle Bullnettle Burclover Burdock Burning nettle Burweed Buttercup Buttonweed Canada thistle Carolina geranium Carpetweed Catchweed bedstraw Catnip Catsear Chickweed, common Chickweed, mouseear Chicory Cinquefoil Clover, crimson Clover, hop Clover, red Clover, strawberry Clover, sweet

Clover, white Cockle Cocklebur Common mullein Creeping jenny Cudweed Curly dock Daisy, English Daisy fleabane Daisy, oxeye Dandelion Dichondra Dogfennel Dollarweed False dandelion Falseflax False sunflower Fiddleneck Florida pusley Frenchweed Galinsoga Goldenrod Ground ivy Gumweed Hairy fleabane Hawkweed Healall Heartleaf drymary Heath aster Hedge bindweed Hedge mustard Henbit Hoary cress Hoary plantain Hoary vervain Horsenettle Jimsonweed Knawel Knotweed Kochia Lambsquarters Lespedeza Mallow Matchweed Mexicanweed Milk vetch Morningglory Mouseear hawkweed Mugwort Musk thistle

- WEEDS -Mustard Narrowleaf plantain Narrowleaf vetch Nettle Orange hawkweed Oriental cocklebur Oxalis Parsley-piert Parsnip Pearlwort Pennycress Pepperweed Pigweed Pineywoods bedstraw Plains coreopsis Plantain Poison ivv Poison oak Pokeweed Poorjoe Prairie sunflower Prickly lettuce Prickly sida Prostrate knotweed Prostrate pigweed Prostrate spurge Prostrate vervain Puncturevine Purslane, common Ragweed Red sorrel Redroot pigweed Redstem filaree Rough cinquefoil Rough fleabane Russian pigweed Russian thistle Scarlet pimpernel Scotch thistle Sheep sorrel Shepherdspurse Slender plantain Smallflower galinsoga Smooth dock Smooth pigweed Sorrel Sowthistle Spanishneedles Speedwell

Spiny amaranth

Spiny cocklebur Spiny sowthistle Spotted catsear Spotted spurge Spurweed Stinging nettle Strawberry, India mock Tall nettle Tall vervain Tansy ragwort Tansy mustard Tanweed ' Thistle Trailing crownvetch Tumble mustard Tumble pigweed Velvetleaf Venice mallow Virginia buttonweed Virginia creeper Virginia pepperweed Wavyleaf bullthistle Western clematis Western salsify White mustard Wild aster Wild buckwheat Wild carrot Wild four-o'clock Wild garlic Wild geranium Wild lettuce Wild marigold Wild mustard Wild onion Wild parsnip Wild radish Wild rape Wild strawberry Wild sweet potato Wild vetch Woodsorrel Woolly croton Woolly morningglony Woo'ly plantain Wornseed Yarrev Yellow locket Yellowflower pepperweed and other broadleaf weeds

Page 8 of 8 EPA Reg. No. 2217-643 002217-00643.20081223

NOTHICATION

FEB 1 7 2009