

2817-655  
PJ 182

BEST COPY AVAILABLE

ACCEPTED

NOV 20 1987

2817-655

**GORDON'S**  
PROFESSIONAL  
TURF & ORNAMENTAL PRODUCTS



# TRIMEC<sup>®</sup>

**SOUTHERN**

## BROADLEAF HERBICIDE

FOR SENSITIVE SOUTHERN GRASSES  
BERMUDA & BENTGRASS, BLUEGRASS,  
FESCUE & ZOYSIA, CENTIPEDE & ST. AUGUSTINE GRASSES  
— KEEP FROM FREEZING —



ACTIVE INGREDIENTS:


***Dimethylamine Salt of 2 (2-methyl-4-chlorophenoxy) propionic acid .....	34.73%
***Dimethylamine Salt of 2, 4-Dichlorophenoxyacetic acid .....	18.74%
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid) .....	3.85%
INERT INGREDIENTS .....	42.88%
<b>TOTAL</b>	<b>100.00%</b>

THIS PRODUCT CONTAINS:  
 \*\*2.77 lbs 2 (2-methyl-4-chlorophenoxy) propionic acid per gallon or 28.70%  
 \*\*1.88 lbs 2,4-Dichlorophenoxyacetic acid per gallon or 19.57%  
 \*\*0.34 lb 3,6-dichloro-o-anisic acid per gallon or 3.20%  
 \*\*Isomet Specific by AOAC Method

**KEEP OUT OF REACH OF CHILDREN**  
**DANGER**  
 See Side Panels for Precautionary Statements and  
 Statement of Practical Treatment.

TRIMEC<sup>®</sup> is a registered trademark of FMC GORDON CORPORATION

**NET CONTENTS 2 1/2 U.S. GALLONS (9.46 liters)**



READ THE ENTIRE LABEL FIRST.  
OBSERVE ALL PRECAUTIONS AND  
FOLLOW DIRECTIONS CAREFULLY.

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans & Domestic Animals

**DANGER:** Corrosive, causes damage to eyes. Do not get into eyes, on skin or clothing. Wear goggles or face shield when handling. Hazard if swallowed or inhaled. Avoid exposure to spray mist. Wash thoroughly after handling.

#### Statement of Practical Treatment

**If in Eyes:** In case of eye contact, immediately flush eyes with plenty of water for 15 minutes. Call a physician at once.

**If Swallowed:** Drink one or two glasses of water. Induce vomiting by leaning back of throat with finger. Call a physician at once. Do not induce vomiting or give anything by mouth to an unconscious person.

**If Inhaled:** Remove victim to fresh air and apply respiration if indicated.

**If On Skin:** Wash promptly with soap and water. Rinse thoroughly.

#### Environmental Hazards

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift away from target area.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### STORAGE & DISPOSAL

**STORAGE:** Store in original container in a locked storage area. Keep from freezing. To prevent cross contamination, do not store near other pesticides, fertilizers, seeds, feed or feed.

**PRODUCT DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes are toxic. Irrigation disposal of excess pesticide, spray mixture, or residue is a violation of Federal law. If those wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the nearest Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed, by state and local authorities, by burning. If burned, stay out of smoke.

### OTHER USE PRECAUTIONS

#### OBSERVE CAREFULLY.

1. Avoid drift or spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants.
2. Do not pour spray solutions near these plants.
3. Do not use on Lilies or daisies nor on lawns or turf where desirable clovers are present.
4. Use only 1,1,1-type sprayers.
5. **NOTE:** Coarse sprays are less likely to wind drift.
6. Do not spray roots of trees and ornamentals. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species.
7. Do not spray while grass is emerging from dormancy.
8. Do not apply to newly sown lawns until well established.
9. Do not spray when air temperature exceeds 85°F.
10. Seed can be safely sown 2 to 4 weeks after application.
11. Do not contaminate domestic or irrigation waters.
12. Be careful not to overdose on Carfopate or St. Augustine grass lawns. These species are sensitive to 2,4-D, a component of this product.
13. After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides.
14. Failure to observe the above precautions may result in injury.

#### How to Use and Where:

**Bermuda & Bahiagrass** — Use one quart per acre. Do not overdose closely mowed bahiagrass. Read the use precautions and observe carefully. Do not use more than the recommended rate. Bermuda and bahiagrass are moderately sensitive to 2,4-D.

**Bluegrass, Fescue & Zoysia** — Use one quart to three pints per acre. These species are tolerant to 2,4-D. However, do not overdose.

**St. Augustine & Centipede** — Use 1 to 1½ pints per acre. Do not apply after 3 weeks if necessary. Do not overdose. Do not spray when grass is in, or emerging from, dormancy. Do not make applications when ambient temperature is below 50°F or above 90°F, or expected to be within 22 hours of application. Do not make applications when grass is under stress from drought, insect attack, or other herbicides. St. Augustine and Centipede turfgrasses are sensitive to 2,4-D and great care should be taken to prevent overdosing.

#### General:

Mix in 5 to 200 gallons of water per acre. Use higher rates when weeds are dense or in flower. If hardy weeds persist, make a second application after three weeks. Apply when weeds are young and actively growing. Avoid applications during long excessively dry or hot periods. When grasses are under stress use reduced rates to reduce potential injury. Do not irrigate within 24 hours of application. Do not apply to newly seeded turf until after the first mowing. Seed can be safely sown 2 to 4 weeks after application at recommended rates.

#### WEEDS CONTROLLED

Bedstraw	Knotweed	Sheep Sorrel
Black Medic	Lambsquarters	Shepherd's Purse
Buckhorn	Lespedeza	Spandell
Burdock	Mallow	Sourgrass
Chicory	Morning Glory	Wild Carrot
Chickweed	Peppercorn	Wild Garlic
Clover	Pigweed	Wild Onion
Dandelion	Plantain	Yarrow
Deck	Polygon Ivy	and many
Ground Ivy	Polygon Oak	other broadleaf
Heal-All	Rutabaga	weeds in lawns
Horsetail	Ragweed	and turf.

#### WARRANTY

The manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled uses when applied according to label directions. Because of safety varying use conditions, it is impossible to eliminate all risks even when label directions are followed. Therefore, the manufacturer makes no other implied or express warranty nor is any agent of the manufacturer allowed to do so. Upon purchase of this product, the buyer assumes all risks associated with use of this product. In the event of damage resulting from a breach of warranty, the buyer agrees to accept a refund of the purchase price of the product as full discharge of the manufacturer's liability.

898487  
EPA REG. NO. 2217-855  
EPA EST. NO. 2217-KS-1



ACCEPTED

NOV 20 1987

U.S. Federal Register

2217-6557

**GORDON'S**  
PROFESSIONAL  
TURF & ORNAMENTAL PRODUCTS



# TRIMEC<sup>®</sup>

**SOUTHERN**

## BROADLEAF HERBICIDE

FOR SENSITIVE SOUTHERN GRASSES  
BERMUDA & BENTGRASS, BLUEGRASS,  
FESCUE & ZOYSIA, CENTIPEDE & ST. AUGUSTINE GRASSES  
— KEEP FROM FREEZING —



ACTIVE INGREDIENTS:

*Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid .....	34.73%
**Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid .....	18.74%
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid) .....	3.85%
INERT INGREDIENTS .....	42.68%

TOTAL 100.00%

THIS PRODUCT CONTAINS:

- \*\*2.77 lbs. 2-(2-methyl-4-chlorophenoxy) propionic acid per gallon or 28.70%
- \*\*1.50 lbs. 2,4-Dichlorophenoxyacetic acid per gallon or 15.57%
- \*\*\*0.31 lb. 3,6-dichloro-o-anisic acid per gallon or 3.20%
- \*\*Isomer Specific by AOAC Method.

KEEP OUT OF REACH OF CHILDREN  
**DANGER**

See Side Panels for Precautionary Statements and  
Statement of Practical Treatment.

TRIMEC<sup>®</sup> is a registered trademark of PBMGORDON CORPORATION.

**BEST AVAILABLE COPY**

**NET CONTENTS 2½ U.S. GALLONS (9.46 liters)**

581 2/28  
552-1128  
P M 23

BEST AVAILABLE COPY

### STORAGE & DISPOSAL

**STORAGE:** Store in original container in a locked storage area. Keep from freezing. To prevent cross-contamination, do not store near other pesticides, fertilizers, seeds, food or feed.

**PRODUCT DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or residue is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed, by state and local authorities, by burning. If burned, stay out of smoke.

### OTHER USE PRECAUTIONS.

#### OBSERVE CAREFULLY.

1. Avoid drift or spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants.
2. Do not pour spray solutions near these plants.
3. Do not use on Lippia or dichondra nor on lawns or turf where desirable clovers are present.
4. Use only lawn-type sprayers.
5. NOTE: Coarse sprays are less likely to wind-drift.
6. Do not spray roots of trees and ornamentals. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species.
7. Do not spray while grass is emerging from dormancy.
8. Do not apply to newly seeded lawns until well established.
9. Do not spray when air temperature exceeds 85°F.
10. Seed can be safely sown 3 to 4 weeks after application.
11. Do not contaminate domestic or irrigation waters.
12. Be careful not to overdose on Centipede or St. Augustine grass lawns. These species are sensitive to 2,4-D, a component of this product.
13. After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides.
14. Failure to observe the above precautions may result in injury.

#### How Much to Use and Where:

**Bermuda & Bentgrass** — Use one quart per acre. Do not overdose closely mowed bentgrass. Read the use precautions and observe carefully. Do not use more than the recommended rate. Bermuda and bentgrass are moderately sensitive to 2,4-D.

**Bluegrass, Fescue & Zoysia** — Use one quart to three pints per acre. These species are tolerant to 2,4-D. However, do not overdose.

**St. Augustine & Centipede** — Use 1 to 1½ pints per acre. Repeat after 3 weeks if necessary. Do not overdose. Do not spray when grass is in, or emerging from, dormancy. Do not make applications when ambient temperature is below 50°F or above 90°F, or expected to be within 72 hours of application. Do not make applications when grass is under stress from drought, insect attack, or other herbicides. St. Augustine and Centipede turfgrasses are sensitive to 2,4-D and great care should be taken to prevent overdosing.

#### General:

Mix in 5 to 300 gallons of water per acre. Use higher rates when weeds are dense or in flower. If hardy weeds persist, make a second application after three weeks. Apply when weeds are young and actively growing. Avoid applications during long excessively dry or hot periods. When grasses are under stress use reduced rates to reduce potential injury. Do not irrigate within 24 hours of application. Do not apply to newly seeded turf until after the first mowing. Seed can be safely sown 3 to 4 weeks after application at recommended rates.

#### WEEDS CONTROLLED

Bedstraw	Knotweed	Sheep Sorrel
Black Medic	Lambsquarter	Shepherd's Purse
Buckhorn	Lespedeza	Speedwell
Burdock	Mallow	Spurge
Chicory	Morning Glory	Wild Carrot
Chickweed	Peppergrass	Wild Garlic
Clover	Pigweed	Wild Onion
Dandelion	Plantains	Yarrow
Dock	Poison Ivy	and many
Ground Ivy	Poison Oak	other broadleaf
Heal-All	Purslane	weeds in lawns
Horbit	Ragweed	and turf.

#### WARRANTY:

The manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled uses when applied according to label directions. Because of widely varying use conditions, it is impossible to eliminate all risks even when label directions are followed. Therefore the manufacturer makes no other implied or express warranty nor is any agent of the manufacturer allowed to do so. Upon purchase of this product, the buyer assumes all risks associated with use of this product. In the event of damage resulting from a breach of warranty the buyer agrees to accept a refund of the purchase price of the product as full discharge of the manufacturer's liability.

896/507  
EPA REG NO 2217-655  
EPA EST. NO. 2217-KS-1

 **gbl/gordon**  
corporation  
ESTD 1917, HANCOCK, MAINE



**READ THE ENTIRE LABEL FIRST.  
OBSERVE ALL PRECAUTIONS AND  
FOLLOW DIRECTIONS CAREFULLY.**

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans & Domestic Animals

**DANGER:** Corrosive, causes damage to eyes. Do not get into eyes, on skin or clothing. Wear goggles or face shield when handling. Harmful if swallowed or inhaled. Avoid exposure to spray mist. Wash thoroughly after handling.

#### Statement of Practical Treatment

**If In Eyes:** In case of eye contact, immediately flush eyes with plenty of water for 15 minutes. Call a physician at once.

**If Swallowed:** Drink one or two glasses of water. Induce vomiting by touching back of throat with finger. Call a physician at once. Do not induce vomiting or give anything by mouth to an unconscious person.

**If Inhaled:** Remove victim to fresh air and apply respiration if indicated.

**If On Skin:** Wash promptly with soap and water. Rinse thoroughly.

#### Environmental Hazards

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift away from target area.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**BEST AVAILABLE COPY**

### STORAGE & DISPOSAL

**STORAGE:** Store in original container in a locked storage area. Keep from freezing. To prevent cross-contamination, do not store near other pesticides, fertilizers, seeds, food or feed.

**PRODUCT DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If those wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed, by state and local authorities, by burning. If burned, stay out of smoke.

#### OTHER USE PRECAUTIONS. OBSERVE CAREFULLY.

1. Avoid drift or spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants.
2. Do not pour spray solutions near these plants.
3. Do not use on Lippia or dichondra nor on lawns or turf where desirable clovers are present.
4. Use only lawn-type sprayers.
5. NOTE: Coarse sprays are less likely to wind-drift.
6. Do not spray roots of trees and ornamentals. Do not exceed specified dosages for any area; be particularly careful within the drip-line of trees and other ornamental species.
7. Do not spray while grass is emerging from dormancy.
8. Do not apply to newly seeded lawns until well established.
9. Do not spray when air temperature exceeds 85°F.
10. Seed can be safely sown 3 to 4 weeks after application.
11. Do not contaminate domestic or irrigation waters.
12. Be careful not to overdose on Centipede or St. Augustine grass lawns. These species are sensitive to 2,4-D, a component of this product.
13. After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides.
14. Failure to observe the above precautions may result in injury.

#### How Much to Use and Where:

**Bermuda & Bentgrass** — Use one quart per acre. Do not overdose closely-mowed bentgrass. Heed the use precautions and observe carefully. Do not use more than the recommended rate. Bermuda and bentgrass are moderately sensitive to 2,4-D.

**Bluegrass, Fescue & Zoysia** — acre. These species are tolerant dose.

**St. Augustine & Centipede** — Use after 3 weeks if necessary. Do not apply if grass is in, or emerging from, dormancy when ambient temperature is below 60°F. Do not apply when grass is under stress from drought or herbicides. St. Augustine and Centipede require special care should be taken.

#### General:

Mix in 5 to 300 gallons of water. Do not apply if weeds are dense or in flower. If weeds are in flower, apply after three weeks. Do not apply to actively growing grasses. Avoid application during hot periods. When grasses are dormant, reduce potential injury. Do not irrigate immediately after application. Do not apply to newly seeded lawns. It can be safely sown 3 to 4 weeks after application.

#### WEEDS CONTROLLED

Bedstraw	Knotweed
Black Medic	Lambsquarters
Buckhorn	Lespedeza
Burdock	Mallow
Chicory	Morning Glory
Chickweed	Peppergrass
Clover	Pigweed
Dandelion	Plantain
Dock	Poison Ivy
Ground Ivy	Poison Oak
Heat-All	Purslane
Henbit	Ragweed

#### WARRANTY:

The manufacturer warrants that the product will perform as stated in the product statement given on the label if used when applied according to label conditions. It is impossible to eliminate all weeds. Therefore the manufacturer does not warrant that the product will eliminate all weeds. The manufacturer is not responsible for any damage resulting from a label error or a refund of the purchase price of the product.

896/587

EPA REG. NO. 2217-655

EPA EST. NO. 2217-KS-1