

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval Expires 2-28-95



United States Environmental Protection Agency Washington, DC 20460

Registration [ ] Amendment [ ] Other [X]

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number: 2217-657
2. EPA Product Manager: Joanne I. Miller
3. Proposed Classification: [X] None [ ] Restricted
4. Company/Product (Name): Gordon's Trimec® Weeder Granules L
5. Name and Address of Applicant: PBI/Gordon Corporation, Post Office Box 014090, Kansas City, Missouri 64101
6. Expedited Review: In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. Product Name

Section - II

Amendment - Explain below. [ ] Resubmission in response to Agency letter dated. [ ] Notification - Explain below. [X] Final printed labels in response to Agency letter dated. [ ] "Me Too" Application. [ ] Other - Explain below. [ ] NOTIFICATION - JAN 30 2006

Explanation: Use additional page(s) if necessary. (For section I and Section II.)
NOTIFICATION: Labeling Notification per PRN 98-10. We ask to add advertising claims that are consistent with the Directions for Use previously-approved. These claims may be presented on container labels, advertisement brochures, and other marketing/sales promotional materials. Please refer to page 11 of the appendix of the labeling.
One (1) copy of the proposed labeling is enclosed for your review.
This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 USC Sec. 1001 to willfully make any false statement to the EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Section 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In: Child-Resistant Packaging, Unit Packaging, Water Soluble Packaging, 2. Type of Container: Metal, Plastic, Glass, Paper, Other (Specify)
3. Location of Net Contents Information: Label, Container
4. Size(s) Retail Container
5. Location of Label Directions: On Label, On Labeling accompanying product
6. Manner in Which Label is Affixed to Product: Lithograph, Paper glued, Stenciled, Other

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)
Name: James L. Kunstman, Ph.D. Title: Director of Regulatory Services Telephone No. (Include Area Code): 816-430-6292
Certification: I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.
2. Signature: [Signature] 3. Title: Director of Regulatory Services
4. Typed Name: James L. Kunstman, Ph.D. 5. Date: January 17, 2006
6. Date Application Received (Stamped)

2712

NOTIFICATION

JAN 30 2006

# GORDON'S TRIMEC® WEEDER GRANULES L

Controls dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf weeds.

**ACTIVE INGREDIENTS:**

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid .....	1.65%
Dimethylamine salt of (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid.....	0.40%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid) .....	0.18%

INERT INGREDIENTS: .....	<u>97.77%</u>
<b>TOTAL</b>	<b>100.00%</b>

**THIS PRODUCT CONTAINS:**

- 27.4 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton or 1.37%.
- 6.6 lbs. (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.33%.
- 3.0 lbs. 3,6-dichloro-o-anisic acid equivalent per ton or 0.15%.

Isomer Specific by AOAC Methods  
 Contains the single isomer form of Mecoprop-p.  
 TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

## KEEP OUT OF REACH OF CHILDREN

## CAUTION

See back panel for Precautionary Statements.

**NET WEIGHT: 10 POUNDS**

651/ AP011706  
 EPA REG NO 2217-657  
 EPA EST NO 2217-KS-1  
 MANUFACTURED BY:



2157 Zeeb, 21619 usil  
 P.O. Box 100, Meriden, CT 06450  
 Telephone: 1-800-821-7925

3 7-12

**SUBLABEL 1). OUTDOOR DOMESTIC USE WITH APPLICATIONS BY HOMEOWNERS ONLY.**



**READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans & Domestic Animals**

**CAUTION:** Harmful if swallowed. Causes moderate eye irritation. Avoid breathing dust. Avoid contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

**First Aid**

<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment.</p> <p>You may also call toll-free, 1-877-800-5556 for emergency medical treatment information.</p>	

**ENVIRONMENTAL HAZARDS:**

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until dust has settled.

**STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.

**PESTICIDE DISPOSAL:** Securely wrap bag in several layers of newspaper and discard in trash.

**DISPOSAL:**

**If Empty —** Do not reuse this container. Place in trash or offer for recycling if available.

**If Partly Filled—**Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

4 8 12

**USE PRECAUTIONS:**

Do not apply this product to St. Augustinegrass, carpetgrass, dichondra, ornamentals, or lawns containing desirable clovers.

Avoid spreading granules onto landscape, plantings, flowers, vegetables or ornamentals. Do not spread over exposed roots of trees and ornamentals. Do not exceed specified dosages for any area; be particularly careful within dripline of tree and other ornamental species. Treated areas may be seeded 3 to 4 weeks after application. Application of this product to newly seeded areas should be delayed until the third mowing.

**WHERE TO USE:**

Gordon's Trimec® Weeder Granules L is a Trimec® complex, a combination of three proven weed killers - 2,4-D, Mecoprop-p and Dicamba. Together, they control a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantains, henbit, spurge and many others found in home lawns. Established turf of Kentucky bluegrass, bentgrass, bahiagrass, centipedegrass, fescues, bermudagrass, zoysiagrass, and perennial ryegrass may be treated.

**HOW MUCH TO APPLY:**

**For Kentucky bluegrass, bentgrass, bahiagrass, centipedegrass, fescues, perennial ryegrass, bermudagrass, and zoysiagrass:** Apply 2.0 pounds of product per 1,000 square feet of turfgrass. A second or follow-up application may be required for dense infestations of weeds or for adverse seasonal conditions. The second application of 2.0 pounds of product per 1,000 square feet should be used 30 days after the first application.

The maximum application rate to turf is 1.2 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

**CALIBRATION INSTRUCTIONS:**

A uniform distribution of product on the treated area is essential. Choose a calm day to ensure uniform coverage and to avoid spreading granules onto flowers, vegetables, and ornamental shrubbery. Do not exceed the application rate for any area and use care in the application of this product within the dripline of trees, shrubs, and other ornamentals.

Drop spreaders and rotary spreaders should be calibrated for uniform applications of this product.

For 2.0 lbs. of product per 1,000 sq.ft. of turfgrass: Check your spreader on 500 sq.ft. and adjust to apply at the rate of 1 lb. of product per 500 sq.ft.

Spreader Settings	
Scott.....	XX
Sears.....	XX

- BROADLEAF WEEDS -

Annual yellow sweetclover	Clover, white	Mustard	Spiny cocklebur
Aster	Cockle	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Cocklebur	Narrowleaf vetch	Spotted catsear
Bedstraw	Common mullein	Nettle	Spotted spurge
Beggarticks	Creeping jenny	Orange hawkweed	Spurweed
Betony, Florida	Cudweed	Oriental cocklebur	Stinging nettle
Bindweed, field	Curly dock	Oxalis	Strawberry, India mock
Bird vetch	Daisy, English	Parsley-piert	Tall nettle
Bitter wintercress	Daisy fleabane	Parsnip	Tall vervain
Bittercress, hairy	Daisy, oxeye	Pearlwort	Tansy ragwort
Bitterweed	Dandelion	Pennycress	Tansy mustard
Black-eyed Susan	Dichondra	Pepperweed	Tanweed
Black medic	Dogfennel	Pigweed	Thistle
Black mustard	Dollarweed	Pineywoods bedstraw	Trailing crownvetch
Blackseed plantain	False dandelion	Plains coreopsis	Tumble mustard
Blessed thistle	Falseflax	Plantain	Tumble pigweed
Bloodflower milkweed	False sunflower	Poison ivy	Velvetleaf
Blue lettuce	Fiddleneck	Poison oak	Venice mallow
Blue vervain	Florida pusley	Pokeweed	Virginia buttonweed
Bracted plantain	Frenchweed	Poorjoe	Virginia creeper
Bristly oxtongue	Galinsoga	Prairie sunflower	Virginia pepperweed
Broadleaf dock	Goldenrod	Prickly lettuce	Wavyleaf bullthistle
Broadleaf plantain	Ground ivy	Prickly sida	Western clematis
Broomweed	Gumweed	Prostrate knotweed	Western salsify
Buckhorn	Hairy fleabane	Prostrate pigweed	White mustard
Buckhorn plantain	Hawkweed	Prostrate spurge	Wild aster
Bulbous buttercup	Healall	Prostrate vervain	Wild buckwheat
Bull thistle	Heartleaf drymary	Puncturevine	Wild carrot
Bullnettle	Heath aster	Purslane, common	Wild four-o'clock
Burclover	Hedge bindweed	Ragweed	Wild garlic
Burdock	Hedge mustard	Red sorrel	Wild geranium
Burning nettle	Henbit	Redroot pigweed	Wild lettuce
Burweed	Hoary cress	Redstem filaree	Wild marigold
Buttercup	Hoary plantain	Rough cinquefoil	Wild mustard
Buttonweed	Hoary vervain	Rough fleabane	Wild onion
Canada thistle	Horsenettle	Russian pigweed	Wild parsnip
Carolina geranium	Jimsonweed	Russian thistle	Wild radish
Carpetweed	Knawel	Scarlet pimpernel	Wild rape
Catchweed bedstraw	Knotweed	Scotch thistle	Wild strawberry
Catnip	Kochia	Sheep sorrel	Wild sweet potato
Catsear	Lambsquarters	Shepherdspurse	Wild vetch
Chickweed, common	Lespedeza	Slender plantain	Woodsorrel
Chickweed, mouseear	Mallow	Smallflower galinsoga	Woolly croton
Chicory	Matchweed	Smooth dock	Woolly morningglory
Cinquefoil	Mexicanweed	Smooth pigweed	Woolly plantain
Clover, crimson	Milk vetch	Sorrel	Wormseed
Clover, hop	Morningglory	Sowthistle	Yarrow
Clover, red	Mouseear hawkweed	Spanishneedles	Yellow rocket
Clover, strawberry	Mugwort	Speedwell	Yellowflower pepperweed
Clover, sweet	Musk thistle	Spiny amaranth	and other broadleaf weeds

**SUBLABEL 2). DOMESTIC AND NONDOMESTIC USE PATTERNS.**



**READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans & Domestic Animals**

**CAUTION:** Harmful if swallowed. Causes moderate eye irritation. Avoid breathing dust. Avoid contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

**First Aid**

<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment.</p> <p>You may also call toll free, 1-877-800-5556 for emergency medical treatment information.</p>	

**ENVIRONMENTAL HAZARDS:**

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until dust has settled.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

**STORAGE & DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** For Paper and Plastic Bags - Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or if allowed by state and local authorities by burning. If burned stay out of smoke.

**WHERE TO USE:**

Gordon's Trimec® Weeder Granules L is a Trimec® herbicide containing fertilizer, 2,4-D, mecoprop-p, and dicamba. This granular weed and feed provides plant nutrients and controls a broad spectrum of broadleaf weeds in turfgrass. Supplemental nitrogen, phosphorous, and potash should be applied according to soil tests or State Extension Service recommendations.

Gordon's Trimec® Weeder Granules L is intended to be applied by commercial applicators, maintenance applicators, service technicians, lawn care operators, and landscape personnel. Cool season and warm season grasses established in institutional, ornamental, and residential sites may be treated.

Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations; areas around hospitals, nursing homes, schools, museums, libraries, sport facilities, and office buildings may be treated.

Ornamental sites include turfgrass established around residences, parks, streets, retail outlets, cemeteries, and institutional buildings. Finally, residential sites are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.

Established turf of Kentucky bluegrass, fescues, and perennial ryegrass may be treated. Established turf of bermudagrass, centipedegrass, and zoysiagrass may be treated. Do not apply this product to St. Augustinegrass, carpetgrass, dichondra, ornamentals, or lawns containing desirable clovers.

Applications of Gordon's Trimec® Weeder Granules L to newly seeded cool season grass are recommended after the third mowing or after the grass seedlings have tillered. Applications to newly established warm season turfgrass are recommended at 3 to 4 weeks after the sodding, sprigging, or plugging operations.

**APPLICATION INSTRUCTIONS:**

1. Mow the turfgrass 1 to 2 days before application.
2. Water lawn thoroughly ( $\geq \frac{1}{2}$  inch) at least 1 day before application to sustain moisture until the next watering.
3. Apply when weeds are young and growing actively. Schedule applications in the morning when dew is present. Or, sprinkle or irrigate lightly ( $\leq \frac{1}{4}$  inch) before application to provide maximum control.
4. Rainfall or irrigation within 24 hours after application may reduce the effectiveness. Do not irrigate or water the turfgrass within 24 hours after application.

**HOW MUCH TO APPLY:**

***For Kentucky bluegrass, bentgrass, bahiagrass, centipedegrass, fescues, perennial ryegrass, bermudagrass, and zoysiagrass:*** Apply 1.5 to 2.0 pounds of product per 1,000 square feet of turfgrass. A second or follow-up application may be required for dense infestations of weeds or for adverse seasonal conditions. The second application of 1.5 to 2.0 pounds of product per 1,000 square feet should be used 30 days after the first application.

The maximum application rate to turf is 1.2 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

A uniform distribution of product on the treated area is essential. Choose a calm day to ensure uniform coverage and to avoid spreading granules onto flowers, vegetables, and ornamental shrubbery. Do not exceed the application rate for any area and use care in the application of this product within the dripline of trees, shrubs, and other ornamentals.

**CALIBRATION INSTRUCTIONS:**

1. Drop spreaders, centrifugal (rotary) spreaders, and other granular equipment should be calibrated for accurate and uniform applications of this product.
2. Calibrate the equipment by applying the product over a measured or known area and adjust the equipment according to the recommendations of the equipment manufacturer.
3. Collect the product output over a known distance and swath width. Many drop spreaders are equipped with catch pans.
4. Adjust the spreader opening in the hopper bottom until the desired application rate is obtained.

Equipment settings of drop spreaders, rotary spreaders, and other granular applicators may be presented on the container labels as requested by supplemental registrants.

**Note to Supplemental Registrants:** Refer to the rate table for the appropriate bag sizes and the area to be treated. Select the appropriate bag size and present only the information provided for that bag size on your product labeling.



9 872

Bag Size, Pounds	Desired Rate Per Acre, Pounds	Desired Rate Per 1,000 Sq. Ft., Pounds	Treated Area For This Bag Size	
			Acres	Sq. Ft.
10	65	1.5	-----	6,667
	87	2.0	-----	5,000
12	65	1.5	-----	8,000
	87	2.0	-----	6,000
15	65	1.5	-----	10,000
	87	2.0	-----	7,500
16	65	1.5	-----	10,667
	87	2.0	-----	8,000
18	65	1.5	-----	12,000
	87	2.0	-----	9,000
20	65	1.5	-----	13,333
	87	2.0	-----	10,000
24	65	1.5	-----	16,000
	87	2.0	-----	12,000
25	65	1.5	-----	16,667
	87	2.0	-----	12,500
28	65	1.5	-----	18,667
	87	2.0	-----	14,000
30	65	1.5	0.46	20,000
	87	2.0	-----	15,000
32	65	1.5	0.49	21,333
	87	2.0	-----	16,000
35	65	1.5	0.54	23,333
	87	2.0	-----	17,500
36	65	1.5	0.55	24,000
	87	2.0	-----	18,000
40	65	1.5	0.61	26,667
	87	2.0	0.46	20,000
44	65	1.5	0.67	29,333
	87	2.0	0.51	22,000
48	65	1.5	0.73	32,000
	87	2.0	0.55	24,000
50	65	1.5	0.77	33,333
	87	2.0	0.57	25,000
52	65	1.5	0.80	34,667
	87	2.0	0.60	26,000
56	65	1.5	0.86	37,333
	87	2.0	0.64	28,000
60	65	1.5	0.92	40,000
	87	2.0	0.69	30,000

- BROADLEAF WEEDS -

Annual yellow sweetclover	Clover, white	Mustard	Spiny cocklebur
Aster	Cockle	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Cocklebur	Narrowleaf vetch	Spotted catsear
Bedstraw	Common mullein	Nettle	Spotted spurge
Beggarticks	Creeping jenny	Orange hawkweed	Spurweed
Betony, Florida	Cudweed	Oriental cocklebur	Stinging nettle
Bindweed, field	Curly dock	Oxalis	Strawberry, India mock
Bird vetch	Daisy, English	Parsley-piert	Tall nettle
Bitter wintercress	Daisy fleabane	Parsnip	Tail vervain
Bittercress, hairy	Daisy, oxeye	Pearlwort	Tansy ragwort
Bitterweed	Dandelion	Pennycress	Tansy mustard
Black-eyed Susan	Dichondra	Pepperweed	Tanweed
Black medic	Dogfennel	Pigweed	Thistle
Black mustard	Dollarweed	Pineywoods bedstraw	Trailing crownvetch
Blackseed plantain	False dandelion	Plains coreopsis	Tumble mustard
Blessed thistle	Falseflax	Plantain	Tumble pigweed
Bloodflower milkweed	False sunflower	Poison ivy	Velvetleaf
Blue lettuce	Fiddleneck	Poison oak	Venice mallow
Blue vervain	Florida pusley	Pokeweed	Virginia buttonweed
Bracted plantain	Frenchweed	Poorjoe	Virginia creeper
Bristly oxtongue	Galinsoga	Prairie sunflower	Virginia pepperweed
Broadleaf dock	Goldenrod	Prickly lettuce	Wavyleaf bullthistle
Broadleaf plantain	Ground ivy	Prickly sida	Western clematis
Broomweed	Gumweed	Prostrate knotweed	Western salsify
Buckhorn	Hairy fleabane	Prostrate pigweed	White mustard
Buckhorn plantain	Hawkweed	Prostrate spurge	Wild aster
Bulbous buttercup	Healall	Prostrate vervain	Wild buckwheat
Bull thistle	Heartleaf drymary	Puncturevine	Wild carrot
Bullnettle	Heath aster	Purslane, common	Wild four-o'clock
Burclover	Hedge bindweed	Ragweed	Wild garlic
Burdock	Hedge mustard	Red sorrel	Wild geranium
Burning nettle	Henbit	Redroot pigweed	Wild lettuce
Burweed	Hoary cress	Redstem filaree	Wild marigold
Buttercup	Hoary plantain	Rough cinquefoil	Wild mustard
Buttonweed	Hoary vervain	Rough fleabane	Wild onion
Canada thistle	Horsenettle	Russian pigweed	Wild parsnip
Carolina geranium	Jimsonweed	Russian thistle	Wild radish
Carpetweed	Knawel	Scarlet pimpernel	Wild rape
Catchweed bedstraw	Knotweed	Scotch thistle	Wild strawberry
Catnip	Kochia	Sheep sorrel	Wild sweet potato
Catsear	Lambsquarters	Shepherdspurse	Wild vetch
Chickweed, common	Lespedeza	Slender plantain	Woodsorrel
Chickweed, mouseear	Mallow	Smallflower galinsoga	Woolly croton
Chicory	Matchweed	Smooth dock	Woolly morningglory
Cinquefoil	Mexicanweed	Smooth pigweed	Woolly plantain
Clover, crimson	Milk vetch	Sorrel	Wormseed
Clover, hop	Morningglory	Sowthistle	Yarrow
Clover, red	Mouseear hawkweed	Spanishneedles	Yellow rocket
Clover, strawberry	Mugwort	Speedwell	Yellowflower pepperweed
Clover, sweet	Musk thistle	Spiny amaranth	and other broadleaf weeds

11-8-12

**LIMITED WARRANTY AND DISCLAIMER.**

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

-12 8 12

**APPENDIX**

***i. Advertising claims that may be presented on the retail containers:***

- ◆ Controls dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf weeds (some of which are listed on the back panel).
- ◆ Controls dandelion, plantain, clover, chickweed and many over (Number) other weeds.
- ◆ For lawns
- ◆ Controls \_\_\_ (number) broadleaf weeds in home lawns.
- ◆ Fewer weeds mean a more luxurious lawn.

