

2217-657

7/23/2010

1/4



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number:
2217-657

Date of Issuance:
JUL 23 2010

NOTICE OF PESTICIDE:

Registration
[X] Reregistration
(under FIFRA, as amended)

Term of Issuance:

Name of Pesticide Product:

Gordon's Trimec Weeder
Granules L

Name and Address of Registrant (include ZIP Code):

PBI/Gordon Corporation
1217 West 12th Street
Kansas City, MO 64101

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is reregistered in accordance with FIFRA section 4(g)(2)(C) provided you:

- 1. Submit and/or cite all data required for registration review/reregistration of your product when the Agency requires all registrants of similar products to submit data.
2. Make the following revisions to sublabel 1:
a. Revise "Inert Ingredients" to "Other Ingredients."
b. Per the acute toxicity review for this product, the Hazards to Humans and Domestic Animals must be revised to read:
"CAUTION
Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water before eating, drinking, chewing gum, using tobacco, or using the toilet."

Continued on Page 2

Signature of Approving Official:

FOR Kathryn V. Montague
Project Manager 23
Herbicide Branch
Registration Division (7505P)

Date:

JUL 23 2010

- c. The groundwater advisory text (“This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater”) appearing in the 2,4-D RED label table must be added to the Environmental Hazards section described as ‘alternate per PRN 2008-1.’
- d. Remove the word “General” from the “General Precautions and Restrictions” heading. This word is an implied safety claim that makes all associated text unenforceable.
- e. Revise the ‘Pesticide Disposal’ statement to read as follows:
“Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.”

3. Make the following revisions to sublabel 2:

- a. Revise “Inert Ingredients” to “Other Ingredients.”
- b. Per the acute toxicity review, the Hazards to Humans and Domestic Animals section must be revised to read:

“CAUTION
Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing.”

- c. Per the 2,4-D, Dicamba, and MCPP-p REDs, the following User Safety Recommendations must be added to this section of the label and the text must be placed in a box:

<p>“USER SAFETY RECOMMENDATIONS</p> <p>User should:</p> <ul style="list-style-type: none">-Wash hands before eating, drinking, chewing gum or tobacco, or using the toilet.-Remove clothing/PPE immediately if pesticide gets inside. The wash thoroughly and put on clean clothing.-Remove clothing/PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.”

- d. Revise the first sentence of the Environmental Hazards Statement to read as follows:
“This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants.”

- e. Remove the word "General" from the "General Precautions and Restrictions" heading. This word is an implied safety claim that makes all associated text unenforceable.

A stamped copy of your labeling is enclosed for your records. You must submit one (1) copy of the final printed label before you release the product for shipment or the next printing of the label whichever occurs first. Products shipped after 12 months from the date of this Notice must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Enclosure

Stamped 11/7/08

4/14

Language Common to All Sublabels

GORDON'S TRIMEC® WEEDER GRANULES L

ACCEPTED with COMMENTS In EPA Letter Dated: JUL 23 2010

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

2217-657

EPA Reg. No. 2217-657

ACTIVE INGREDIENT:

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	1.65%
Dimethylamine salt of (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid	0.40%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)	0.18%

INERT INGREDIENTS:	97.77%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- 27.4 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton or 1.37%.
- 6.6 lbs. (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.33%.
- 3.0 lbs. 3,6-dichloro-o-anisic acid equivalent per ton or 0.15%.


Isomer Specific by AOAC Methods

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.



SUBLABEL 1: OUTDOOR DOMESTIC USE WITH APPLICATIONS BY HOMEOWNERS ONLY

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear long-sleeved shirt, long pants, socks and shoes. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or on clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call toll-free, 1-877-800-5556 for emergency medical treatment information.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate. This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

[Environmental Hazards (alternate per PRN 2008-1)

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.]

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

- Do not apply this product to St. Augustinegrass, carpetgrass, dichondra, ornamentals, or lawns containing desirable clovers.
- Do not exceed specified dosages for any area. Treated areas may be seeded 3 to 4 weeks after application. Application of this product to newly seeded areas should be delayed until the third mowing.
- Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

WHERE TO USE:

Gordon's Trimec® Weeder Granules L is a Trimec® complex, a combination of three proven weed killers - 2,4-D, Mecoprop-p and Dicamba. Together, they control a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantains, henbit, spurge and many others found in home lawns. Established turf of Kentucky bluegrass, bentgrass, bahiagrass, centipedegrass, fescues, bermudagrass, zoysiagrass, and perennial ryegrass may be treated.

HOW MUCH TO APPLY:

For Kentucky bluegrass, bentgrass, bahiagrass, centipedegrass, fescues, perennial ryegrass, bermudagrass, and zoysiagrass: Apply 2.0 pounds of product per 1,000 square feet of turfgrass. A second or follow-up application may be required for dense infestations of weeds or for adverse seasonal conditions. The second application of 2.0 pounds of product per 1,000 square feet should be used 30 days after the first application.

The maximum broadcast application rate is 2.0 pounds of product per 1,000 sq.ft. per application (1.2 lbs 2,4-D ae, 0.29 lb MCP-P ae and 0.13 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 4.0 pounds of product per 1,000 sq.ft. (2.4 lbs 2,4-D ae, 0.58 lb MCP-P ae and 0.26 lb dicamba ae per acre).

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

CALIBRATION INSTRUCTIONS:

A uniform distribution of product on the treated area is essential. Choose a calm day to ensure uniform coverage. Do not exceed the application rate for any area.

Drop spreaders and rotary spreaders should be calibrated for uniform applications of this product.

For 2.0 lbs. of product per 1,000 sq.ft. of turfgrass: Check your spreader on 500 sq.ft. and adjust to apply at the rate of 1 lb. of product per 500 sq.ft.

Spreader Settings	
Scotts	XX
Sears	XX

BROADLEAF WEEDS

Annual yellow sweetclover	Clover, white	Mustard	Spiny cocklebur
Aster	Cockle	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Cocklebur	Narrowleaf vetch	Spotted catsear
Bedstraw	Common mullein	Nettle	Spotted spurge
Beggarticks	Creeping jenny	Orange hawkweed	Spurweed
Betony, Florida	Cudweed	Oriental cocklebur	Stinging nettle
Bindweed, field	Curly dock	Oxalis	Strawberry, India mock
Bird vetch	Daisy, English	Parsley-piert	Tall nettle
Bitter wintercress	Daisy fleabane	Parsnip	Tall vervain
Bittercress, hairy	Daisy, oxeye	Pearlwort	Tansy ragwort
Bitterweed	Dandelion	Pennycress	Tansy mustard
Black-eyed Susan	Dichondra	Pepperweed	Tanweed
Black medic	Dogfennel	Pigweed	Thistle
Black mustard	Dollarweed	Pineywoods bedstraw	Trailing crownvetch
Blackseed plantain	False dandelion	Plains coreopsis	Tumble mustard
Blessed thistle	Falseflax	Plantain	Tumble pigweed
Bloodflower milkweed	False sunflower	Poison ivy	Velvetleaf
Blue lettuce	Fiddleneck	Poison oak	Venice mallow
Blue vervain	Florida pusley	Pokeweed	Virginia buttonweed
Bracted plantain	Frenchweed	Poorjoe	Virginia creeper
Bristly oxtongue	Galinsoga	Prairie sunflower	Virginia pepperweed
Broadleaf dock	Goldenrod	Prickly lettuce	Wavyleaf bullthistle
Broadleaf plantain	Ground ivy	Prickly sida	Western clematis
Broomweed	Gumweed	Prostrate knotweed	Western salsify
Buckhorn	Hairy fleabane	Prostrate pigweed	White mustard
Buckhorn plantain	Hawkweed	Prostrate spurge	Wild aster
Bulbous buttercup	Healall	Prostrate vervain	Wild buckwheat
Bull thistle	Heartleaf drymary	Puncturevine	Wild carrot
Bullnettle	Heath aster	Purslane, common	Wild four-o'clock
Burclover	Hedge bindweed	Ragweed	Wild garlic
Burdock	Hedge mustard	Red sorrel	Wild geranium
Burning nettle	Henbit	Redroot pigweed	Wild lettuce
Burweed	Hoary cress	Redstem filaree	Wild marigold
Buttercup	Hoary plantain	Rough cinquefoil	Wild mustard
Buttonweed	Hoary vervain	Rough fleabane	Wild onion
Canada thistle	Horsenettle	Russian pigweed	Wild parsnip
Carolina geranium	Jimsonweed	Russian thistle	Wild radish
Carpetweed	Knawel	Scarlet pimpernel	Wild rape
Catchweed bedstraw	Knotweed	Scotch thistle	Wild strawberry
Catnip	Kochia	Sheep sorrel	Wild sweet potato
Catsear	Lambsquarters	Shepherdspurse	Wild vetch
Chickweed, common	Lespedeza	Slender plantain	Woodsorrel
Chickweed, mouseear	Mallow	Smallflower galinsoga	Woolly croton
Chicory	Matchweed	Smooth dock	Woolly morningglory
Cinquefoil	Mexicanweed	Smooth pigweed	Woolly plantain
Clover, crimson	Milk vetch	Sorrel	Wormseed
Clover, hop	Morningglory	Sowthistle	Yarrow
Clover, red	Mouseear hawkweed	Spanishneedles	Yellow rocket
Clover, strawberry	Mugwort	Speedwell	Yellowflower pepperweed
Clover, sweet	Musk thistle	Spiny amaranth	and other broadleaf weeds

8/2

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.

PESTICIDE DISPOSAL: Securely wrap bag in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: If Empty - Do not reuse this container. Place in trash or offer for recycling if available. **If Partly Filled** - Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

SUBLABEL 2: DOMESTIC AND NONDOMESTIC USE PATTERNS

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are natural rubber. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All loaders, applicators, and other handlers must wear:

- long-sleeved shirt and long pants,
- shoes plus socks.
- chemical-resistant gloves

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

First Aid	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call toll free, 1-877-800-5556 for emergency medical treatment information.	

Environmental Hazards

This pesticide may be toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL PRECAUTIONS AND RESTRICTIONS:

- Do not enter or allow people (or pets) to enter the treated area until dusts have settled.
- Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

WHERE TO USE:

Gordon's Trimec® Weeder Granules L is a Trimec® herbicide containing fertilizer, 2,4-D, mecoprop-p, and dicamba. This granular weed and feed provides plant nutrients and controls a broad spectrum of broadleaf weeds in turfgrass. Supplemental nitrogen, phosphorous, and potash should be applied according to soil tests or State Extension Service recommendations.

Gordon's Trimec® Weeder Granules L is intended to be applied by commercial applicators, maintenance applicators, service technicians, lawn care operators, and landscape personnel. Cool season and warm season grasses established in institutional, ornamental, and residential sites may be treated.

Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations; areas around hospitals, nursing homes, schools, museums, libraries, sport facilities, and office buildings may be treated.

Ornamental sites include turfgrass established around residences, parks, streets, retail outlets, cemeteries, and institutional buildings. Finally, residential sites are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.

Established turf of Kentucky bluegrass, fescues, and perennial ryegrass may be treated. Established turf of bermudagrass, centipedegrass, and zoysiagrass may be treated. Do not apply this product to St. Augustinegrass, carpetgrass, dichondra, ornamentals, or lawns containing desirable clovers.

Applications of Gordon's Trimec® Weeder Granules L to newly seeded cool season grass are recommended after the third mowing or after the grass seedlings have tillered. Applications to newly established warm season turfgrass are recommended at 3 to 4 weeks after the sodding, sprigging, or plugging operations.

APPLICATION INSTRUCTIONS:

1. Mow the turfgrass 1 to 2 days before application.
2. Water lawn thoroughly ($\geq 1/2$ inch) at least 1 day before application to sustain moisture until the next watering.
3. Apply when weeds are young and growing actively. Schedule applications in the morning when dew is present. Or, sprinkle or irrigate lightly ($\leq 1/4$ inch) before application to provide maximum control.
4. Rainfall or irrigation within 24 hours after application may reduce the effectiveness. Do not irrigate or water the turfgrass within 24 hours after application.

HOW MUCH TO APPLY:

For Kentucky bluegrass, bentgrass, bahiagrass, centipedegrass, fescues, perennial ryegrass, bermudagrass, and zoysiagrass: Apply 1.5 to 2.0 pounds of product per 1,000 square feet of turfgrass. A second or follow-up application may be required for dense infestations of weeds or for adverse seasonal conditions. The second application of 1.5 to 2.0 pounds of product per 1,000 square feet should be used 30 days after the first application.

The maximum broadcast application rate is 2.0 pounds of product per 1,000 sq.ft. per application (1.2 lbs 2,4-D ae, 0.29 lb MCPP-p ae and 0.13 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 4.0 pounds of product per 1,000 sq.ft. (2.4 lbs 2,4-D ae, 0.58 lb MCPP-p ae and 0.26 lb dicamba ae per acre).

A uniform distribution of product on the treated area is essential. Choose a calm day to ensure uniform coverage. Do not exceed the application rate for any area.

Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result.

CALIBRATION INSTRUCTIONS:

1. Drop spreaders, centrifugal (rotary) spreaders, and other granular equipment should be calibrated for accurate and uniform applications of this product.
2. Calibrate the equipment by applying the product over a measured or known area and adjust the equipment according to the recommendations of the equipment manufacturer.
3. Collect the product output over a known distance and swath width. Many drop spreaders are equipped with catch pans.
4. Adjust the spreader opening in the hopper bottom until the desired application rate is obtained.
5. Equipment settings of drop spreaders, rotary spreaders, and other granular applicators may be presented on the container labels as requested by supplemental registrants.

[Note to Supplemental Registrants: Refer to the rate table for the appropriate bag sizes and the area to be treated. Select the appropriate bag size and present only the information provided for that bag size on your product labeling.]

Bag Size, Pounds	Desired Rate Per Acre, Pounds	Desired Rate Per 1,000 Sq.Ft., Pounds	Treated Area For This Bag Size	
			Acres	Sq. Ft.
10	65	1.5	----	6,667
	87	2.0	----	5,000
12	65	1.5	----	8,000
	87	2.0	----	6,000
15	65	1.5	----	10,000
	87	2.0	----	7,500
16	65	1.5	----	10,667
	87	2.0	----	8,000
18	65	1.5	----	12,000
	87	2.0	----	9,000
20	65	1.5	----	13,333
	87	2.0	----	10,000
24	65	1.5	----	16,000
	87	2.0	----	12,000
25	65	1.5	----	16,667
	87	2.0	----	12,500
28	65	1.5	----	18,667
	87	2.0	----	14,000
30	65	1.5	0.46	20,000
	87	2.0	----	15,000
32	65	1.5	0.49	21,333
	87	2.0	----	16,000
35	65	1.5	0.54	23,333

Bag Size, Pounds	Desired Rate Per Acre, Pounds	Desired Rate Per 1,000 Sq.Ft., Pounds	Treated Area For This Bag Size	
			Acres	Sq. Ft.
	87	2.0	----	17,500
36	65	1.5	0.55	24,000
	87	2.0	----	18,000
40	65	1.5	0.61	26,667
	87	2.0	0.46	20,000
44	65	1.5	0.67	29,333
	87	2.0	0.51	22,000
48	65	1.5	0.73	32,000
	87	2.0	0.55	24,000
50	65	1.5	0.77	33,333
	87	2.0	0.57	25,000
52	65	1.5	0.80	34,667
	87	2.0	0.60	26,000
56	65	1.5	0.86	37,333
	87	2.0	0.64	28,000
60	65	1.5	0.92	40,000
	87	2.0	0.69	30,000

BROADLEAF WEEDS			
Annual yellow sweetclover	Clover, white	Mustard	Spiny cocklebur
Aster	Cockle	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Cocklebur	Narrowleaf vetch	Spotted catsear
Bedstraw	Common mullein	Nettle	Spotted spurge
Beggarticks	Creeping jenny	Orange hawkweed	Spurweed
Betony, Florida	Cudweed	Oriental cocklebur	Stinging nettle
Bindweed, field	Curly dock	Oxalis	Strawberry, India mock
Bird vetch	Daisy, English	Parsley-piert	Tall nettle
Bitter wintercress	Daisy fleabane	Parsnip	Tall vervain
Bittercress, hairy	Daisy, oxeye	Pearlwort	Tansy ragwort
Bitterweed	Dandelion	Pennycress	Tansy mustard
Black-eyed Susan	Dichondra	Pepperweed	Tanweed
Black medic	Dogfennel	Pigweed	Thistle
Black mustard	Dollarweed	Pineywoods bedstraw	Trailing crownvetch
Blackseed plantain	False dandelion	Plains coreopsis	Tumble mustard
Blessed thistle	Falseflax	Plantain	Tumble pigweed
Bloodflower milkweed	False sunflower	Poison ivy	Velvetleaf
Blue lettuce	Fiddleneck	Poison oak	Venice mallow
Blue vervain	Florida pusley	Pokeweed	Virginia buttonweed
Bracted plantain	Frenchweed	Poorjoe	Virginia creeper
Bristly oxtongue	Galinsoga	Prairie sunflower	Virginia pepperweed
Broadleaf dock	Goldenrod	Prickly lettuce	Wavyleaf bullthistle
Broadleaf plantain	Ground ivy	Prickly sida	Western clematis
Broomweed	Gumweed	Prostrate knotweed	Western salsify
Buckhorn	Hairy fleabane	Prostrate pigweed	White mustard
Buckhorn plantain	Hawkweed	Prostrate spurge	Wild aster
Bulbous buttercup	Healall	Prostrate vervain	Wild buckwheat
Bull thistle	Heartleaf drymary	Puncturevine	Wild carrot
Bullnettle	Heath aster	Purslane, common	Wild four-o'clock
Burclover	Hedge bindweed	Ragweed	Wild garlic
Burdock	Hedge mustard	Red sorrel	Wild geranium
Burning nettle	Henbit	Redroot pigweed	Wild lettuce
Burweed	Hoary cress	Redstem filaree	Wild marigold
Buttercup	Hoary plantain	Rough cinquefoil	Wild mustard
Buttonweed	Hoary vervain	Rough fleabane	Wild onion
Canada thistle	Horsenettle	Russian pigweed	Wild parsnip

BROADLEAF WEEDS			
Carolina geranium	Jimsonweed	Russian thistle	Wild radish
Carpetweed	Knawel	Scarlet pimpernel	Wild rape
Catchweed bedstraw	Knotweed	Scotch thistle	Wild strawberry
Catnip	Kochia	Sheep sorrel	Wild sweet potato
Catsear	Lambsquarters	Shepherdspurse	Wild vetch
Chickweed, common	Lespedeza	Slender plantain	Woodsorrel
Chickweed, mouseear	Mallow	Smallflower galinsoga	Woolly croton
Chicory	Matchweed	Smooth dock	Woolly morningglory
Cinquefoil	Mexicanweed	Smooth pigweed	Woolly plantain
Clover, crimson	Milk vetch	Sorrel	Wormseed
Clover, hop	Morningglory	Sowthistle	Yarrow
Clover, red	Mouseear hawkweed	Spanishneedles	Yellow rocket
Clover, strawberry	Mugwort	Speedwell	Yellowflower pepperweed and other broadleaf weeds
Clover, sweet	Musk thistle	Spiny amaranth	

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: For Paper and Plastic Bags - Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or if allowed by state and local authorities by burning. If burned stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

APPENDIX

1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Weight: _____
- EPA Est. No. _____

2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

- Controls dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf weeds (some of which are listed on the back panel).
- Controls dandelion, plantain, clover, chickweed and many over (number) other weeds.
- For lawns
- Controls ___ (number) broadleaf weeds in home lawns.
- Fewer weeds mean a more luxurious lawn.
- Controls dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf weeds.

DOCUMENT CONTROL INFORMATION

1. **Unique Label Identifier:** 002217-00657.20081105.proposed.doc

2. **Reason for Issue:** Various changes pertaining to re-registration. Re-format label.