

# †TRIMEC®

## WEED & FEED 33 1/2

24-4-8



18 Pounds Covers up to 5,000 Square Feet

**KILLS:** Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and many other broadleaf weeds; some of which are listed on the back panel.

#### ACTIVE INGREDIENTS:

- \*2,4-Dichlorophenoxyacetic acid
- \*\*2-(2-methyl-4-chlorophenoxy) propionic acid
- \*\*\*Dicamba (3,6-dichloro-o-anisic acid)

#### INERT INGREDIENTS

#### THIS PRODUCT CONTAINS:

- \*15.22 lbs 2,4-Dichlorophenoxyacetic acid per ton
- \*\*6.82 lbs 2-(2-methyl-4-chlorophenoxy) propionic acid per ton
- \*\*\*1.44 lbs 3,6-dichloro-o-anisic acid per ton

0.781%  
0.741%  
0.072%  
98.826%

TOTAL 100.000%

APR 29 1987

2217-666

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

#### STATEMENT OF PRACTICAL TREATMENT


In case of contact, wash skin with plenty of soap and water, for eyes, flush with water for 15 minutes and get medical attention.

See back panel for additional precautionary statements

TRIMEC® is a registered trademark of GORDON CORPORATION

U.S. Patent No. 3,664,386

**NET WEIGHT 18 POUNDS**



GORDON CORPORATION



READ THE ENTIRE LABEL FIRST  
OBSERVE ALL CAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

#### PRECAUTIONARY STATEMENTS

##### Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. Wash thoroughly after handling.

##### Environmental Hazards

Do not apply directly to water. Do not apply when weather conditions favor drift from target area.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs. To prevent cross contamination, do not store near other fertilizers, or near herbicides, insecticides, fungicides, or seeds. **PRODUCT DISPOSAL:** To dispose of bags which contain unused product, securely wrap bag in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse empty bag. Fold empty bag and discard in trash.

#### USE PRECAUTIONS

Do not use this product where desirable clovers are present or allow the material to wind drift onto flowers, vegetables or ornamental shrubbery. Do not apply in areas underlain by roots of desirable trees or shrubs. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning. Do not contaminate domestic or irrigation waters.

#### INSTRUCTIONS

Gordon's Weed & Feed 33 1/2 is a TRIMEC® complex, a combination of three proven weed killers — 2,4-D, Mecoprop and Dicamba. Together, they control a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a 24-4-8 fertilizer provides a generous supply of plant food needed for a normal season's growth. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days. Do not use on dichondra, carpetgrass, closely mowed bentgrass, putting greens or St. Augustine lawns.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to help the material and prevent dust drift onto non-target plants.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

Be patient, TRIMEC works slowly and gently and is friendly to your lawn.

#### SPREADER SETTINGS

|             |       |
|-------------|-------|
| Scott       | 6     |
| Lawn Beauty | 5     |
| Jackson     | 7 1/2 |
| Sears       | 4     |

The above spreader settings are approximate. Check your spreader on 275 sq. ft. and adjust to apply at the rate of 1 lb. per 275 sq. ft.

#### FERTILIZER GUARANTEE

|                                           |     |
|-------------------------------------------|-----|
| Total Nitrogen (not less than)            | 24% |
| Water Soluble Nitrogen (not less than)    | 5%  |
| Available Phosphoric Acid (not less than) | 4%  |
| Soluble Potash (not less than)            | 8%  |

| WEEDS       |              |               |                 |
|-------------|--------------|---------------|-----------------|
| REDSTRAW    | DANDELION    | LESPEDEZA     | POISON IVY      |
| BLACK MEDIC | DOCK         | MALLOW        | POISON OAK      |
| BURKHORN    | GROUND IVY   | MORNING GLORY | PURSLANE        |
| BURDOCK     | HEAL ALL     | OXALIS        | RAGWEED         |
| CHICORY     | HENBIT       | PEPPERGRASS   | SHEEP SOREL     |
| CHICKWEED   | KNOTWEED     | PIGWEEED      | SHEPHERDS PURSE |
| CLOVER      | LANDSQUARTER | PLANTAINS     | SPEEDWELL       |
|             |              |               | YARROW          |

**NOTICE:** Seller warrants that this product conforms to the important statement on the label. Such conditions of use, such as weather, compatibility with other chemicals, and conditions of application, equipment will vary. Seller makes no claims other than those stated on this label.

EPA REG. No. 2217-660

EPA EST. No. 2217-RS-1

578 1293