

2917-660

3-21-2003

1123

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval Expires 2-28-95

	United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
--	---	--	-----------------------

Application for Pesticide - Section I

1. Company/Product Number 2217-660	2. EPA Product Manager Joanne I. Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Gordon's Trimec® Weed and Feed 33 1/2	PM# Product Manger, Team 23	
5. Name and Address of Applicant (Include ZIP Code) PBI/Gordon Corporation Post Office Box 014090 Kansas City, Missouri 64101 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	NOTIFICATION MAR 21 2003
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)
NOTIFICATION: 03/11/03 Minor revisions to Directions for Use per PRN 98-10.
 1. We ask to list one advertising claim that may be presented on the container labeling of Supplemental Registrants. Please refer to the appendix on pages 22 for this request.
 This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or Confidential Statement of Formula of this product. I understand that it is a violation of 18 U.S.C Sec. 1001 to willfully make any false statement to the EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Section 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged in:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____		
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per Container		
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product		<input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled <input type="checkbox"/> Other _____			

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Craig Martens	Title Manager of Regulatory Services	Telephone No. (Include Area Code) 816-460-6287
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Manager of Regulatory Services	
4. Typed Name Craig Martens	5. Date March 11, 2003	

2/23

GORDON'S TRIMEC[®] WEED & FEED 33½ 24-4-8

ACTIVE INGREDIENTS:

2,4-D	0.76%
Mecoprop-p (MCP-p).....	0.17%
Dicamba	0.07%
OTHER INGREDIENTS:	99.00%
TOTAL	100.000%

THIS PRODUCT CONTAINS:

15.2 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton
 3.4 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton
 1.4 lbs. 3,6-dichloro-o-anisic acid equivalent per ton
 Isomer Specific by AOAC Methods
 Contains single isomer form of Mecoprop-p
 TRIMEC[®] is a registered trademark of PBI/GORDON CORPORATION

CAS Registry Numbers:

2,4-D CAS# 94-75-7
 Mecoprop-p CAS# 16484-77-8
 Dicamba CAS# 1918-00-9

KEEP OUT OF REACH OF CHILDREN

CAUTION

**NOTIFICATION
MAR 2 1 2003**

See back panel for additional Precautionary Statements.

NET WEIGHT: ____ POUNDS

578 AP031103
 EPA REG NO 2217-660
 EPA EST NO 2217-KS-1
 MANUFACTURED BY:



3/23



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

First Aid	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-xxx-xxxx for emergency medical treatment information.</p>	

ENVIRONMENTAL HAZARDS:

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater. Do not apply when weather conditions favor drift from target area.

0
 1
 2
 3
 4
 5
 6
 7
 8
 9
 10
 11
 12
 13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65
 66
 67
 68
 69
 70
 71
 72
 73
 74
 75
 76
 77
 78
 79
 80
 81
 82
 83
 84
 85
 86
 87
 88
 89
 90
 91
 92
 93
 94
 95
 96
 97
 98
 99

OPTION 1.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

CONTAINER DISPOSAL: **If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partially filled:** Call your local solid waste agency or toll free number (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.

USE PRECAUTIONS:

Do not use on dichondra, carpetgrass or St. Augustinegrass.

Do not use this product where desirable clovers are present or allow the material to drift onto flowers, vegetables or ornamental shrubbery. Do not apply in areas underlaid by roots of desirable trees or shrubs. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning.

INSTRUCTIONS:

Gordon's Trimec® Weed & Feed 33½ is a Trimec® complex, a combination of three proven weed killers -- 2,4-D, Mecoprop-p and Dicamba. Together, they control a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a 24-4-8 fertilizer provides a supply of plant nutrients. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth.

Apply 2.4 to 3.6 pounds of product per 1,000 square feet of turfgrass. A second application may be made in approximately 30 days.

The maximum application rate to turf is 1.2 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

WHEN TO APPLY:

Apply when broadleaf weeds are actively growing in spring or early fall.

WHAT TO EXPECT:

Weed stems and leaves will begin to twist and curl in a few days, but a month or more may be required for complete control. This product will not prevent weed seeds from producing new plants.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants
4. Avoid applying when temperatures are consistently above 90°F.
5. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.


CALIBRATION INSTRUCTIONS:

To provide proper distribution, calibrate spreader before application. [Refer to table for appropriate bag sizes.]

- ◆ For 2.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 420 square feet and adjust to apply at the rate of 1 pound of product per 420 square feet.
- ◆ For 2.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 385 square feet and adjust to apply at the rate of 1 pound of product per 385 square feet.
- ◆ For 2.8 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 360 square feet and adjust to apply at the rate of 1 pound of product per 360 square feet.
- ◆ For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 333 square feet and adjust to apply at the rate of 1 pound of product per 333 square feet.
- ◆ For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.

Spreader settings for 1.0 pound of product per 310 square feet	
Scotts EasyGreen® Rotary	26
Scotts Drop.....	5
Scotts SpeedyGreen® Rotary.....	3½
GreenView Drop (Spred II).....	4
Central Drop (#205).....	5½
Sears Drop (#19810).....	4
Ortho Drop (#5000).....	5
Precision Drop (Red Devil).....	4

- ◆ For 3.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 300 square feet and adjust to apply at the rate of 1 pound of product per 300 square feet.
- ◆ For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.



Note to Supplemental Registrants: Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, equipment settings of drop spreaders, rotary spreaders, and other granular applicators may be presented on the container labels.

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.
12	2.4	5,000
	2.6	4,600
	2.8	4,300
	3.0	4,000
	3.2	3,800
	3.4	-----
	3.6	-----
13	2.4	5,400
	2.6	5,000
	2.8	4,600
	3.0	4,300
	3.2	4,100
	3.4	3,800
	3.6	3,600
14	2.4	5,800
	2.6	5,400
	2.8	5,000
	3.0	4,700
	3.2	4,400
	3.4	4,100
	3.6	3,900
15	2.4	6,300
	2.6	5,800
	2.8	5,400
	3.0	5,000
	3.2	4,700
	3.4	4,400
	3.6	4,200
16	2.4	6,700
	2.6	6,200
	2.8	5,700
	3.0	5,300
	3.2	5,000
	3.4	4,700
	3.6	4,400
17	2.4	7,100
	2.6	6,500
	2.8	6,100
	3.0	5,700
	3.2	5,300
	3.4	5,000
	3.6	4,700
18	2.4	7,500
	2.6	6,900
	2.8	6,400
	3.0	6,000
	3.2	5,600
	3.4	5,300
	3.6	5,000

7/23

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.
24	2.4	10,000
	2.6	9,200
	2.8	8,600
	3.0	8,000
	3.2	7,500
	3.4	7,100
	3.6	6,700
25	2.4	10,400
	2.6	9,600
	2.8	8,900
	3.0	8,300
	3.2	7,800
	3.4	7,400
	3.6	6,900
26	2.4	10,800
	2.6	10,000
	2.8	9,300
	3.0	8,700
	3.2	8,100
	3.4	7,600
	3.6	7,200
28	2.4	11,700
	2.6	10,800
	2.8	10,000
	3.0	9,300
	3.2	8,800
	3.4	8,200
	3.6	7,800
36	2.4	15,000
	2.6	13,800
	2.8	12,900
	3.0	12,000
	3.2	11,300
	3.4	10,500
	3.6	10,000
37	2.4	15,400
	2.6	14,200
	2.8	13,200
	3.0	12,300
	3.2	11,600
	3.4	10,900
	3.6	10,300
39	2.4	16,300
	2.6	15,000
	2.8	13,900
	3.0	13,000
	3.2	12,200
	3.4	11,500
	3.6	10,800
42	2.4	17,500
	2.6	16,200
	2.8	15,000
	3.0	14,000
	3.2	13,100
	3.4	12,400
	3.6	11,700

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.
43	2.4	17,900
	2.6	16,500
	2.8	15,400
	3.0	14,300
	3.2	13,400
	3.4	12,600
	3.6	11,900
48	2.4	20,000
	2.6	18,500
	2.8	17,100
	3.0	16,000
	3.2	15,000
	3.4	14,100
	3.6	13,300
49	2.4	20,400
	2.6	18,800
	2.8	17,500
	3.0	16,300
	3.2	15,300
	3.4	14,400
	3.6	13,600

DISTRIBUTION AND SPREADER PATTERNS:

Always apply with a properly calibrated rotary or drop type fertilizer spreader to distribute the granules evenly. Never apply by hand.

Coverage - To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to XXX square feet. *[Graphics Optional]*

SINGLE PASS DISTRIBUTION:

First spread the borders of your lawn. Then spread the center by spreading in straight line passes. Spread at a uniform walking speed to insure even coverage *[Graphics Optional]*. To avoid gaps and streaks, first spread the border and fill in center with uniform, parallel passes.

Advisory For Fertilizers Containing Iron:

Keep this product off sidewalks, patios, driveways, etc. This fertilizer contains iron which will stain concrete and painted surfaces. If it comes in contact with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.

24-4-8 GUARANTEED ANALYSIS	
Total Nitrogen (N).....	24%
Water Insoluble Nitrogen.....	5%
Available Phosphate (P ₂ O ₅).....	4%
Soluble Potash (K ₂ O).....	8%

Fertilizer Labeling Requirements:

- All fertilizer products distributed in Washington at a minimum, must have the following labeling statement: "This product has been registered with the Washington State Department of Agriculture. When applied as directed, this fertilizer meets the Washington standards for arsenic, cadmium, cobalt, mercury, molybdenum, lead, nickel, selenium, and zinc. You have the right to receive specific information about Washington standards from the distributor of this product."
- After July 1, 1999, the label must also state: "Information received by the Washington State Department of Agriculture regarding the components in this product is available on the internet at <http://www.wa.gov/ag/>."
- Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.xxxxx.com>.

Referrals to other sequentially used products:

- For best results, apply (brand name) products.
- Apply using the 4-Step Program
- **4-Step Program:**
 - STEP 1:* Menards Premium Crabgrass Preventor - Apply March - early May - Prevents crabgrass, foxtail, goosegrass, *poa annua* and other annual weed grasses. - Nutralene feeds your turfgrass for up to 2 months with long lasting deep greening.
 - STEP 2:* Menards Premium Weed & Feed - Apply May - June - Kills dandelion, clover, plantain and over XXX other broadleaf weeds. - Nutralene feeds your turfgrass for up to 2 months with long lasting deep greening.
 - STEP 3:* Menards Premium Lawn Food - Apply June - August - Nourishes your lawn throughout the summer months with extended feeding of Nitrogen. - Nutralene feeds your turfgrass for up to 2 months with long lasting deep greening.
 - STEP 4:* Menards Premium Winterizer & Weed Control - Apply late August - mid October - Kills dandelion, clover, plantain and over XXX other broadleaf weeds. - Maximizes root building. - Nutralene feeds your turfgrass for up to 2 months with long lasting deep greening.
- OR**
- STEP 4:* Menards Premium Starter & Winterizer - Apply late August - November - Maximizes root building. - Nutralene feeds your turfgrass for up to 2 months with long lasting deep greening. - Use with new seeding and sodding of turfgrass.

10/23

- BROADLEAF WEEDS -

Annual yellow sweetclover	Clover, white	Mustard	Spiny cocklebur
Aster	Cockle	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Cocklebur	Narrowleaf vetch	Spotted catsear
Bedstraw	Common mullein	Nettle	Spotted spurge
Beggarticks	Creeping jenny	Orange hawkweed	Spurweed
Betony, Florida	Cudweed	Oriental cocklebur	Stinging nettle
Bindweed, field	Curly dock	Oxalis	Strawberry, India mock
Bird vetch	Daisy, English	Parsley-piert	Tall nettle
Bitter wintercress	Daisy fleabane	Parsnip	Tall vervain
Bittercress, hairy	Daisy, oxeve	Pearlwort	Tansy ragwort
Bitterweed	Dandelion	Pennycress	Tansy mustard
Black-eyed Susan	Dichondra	Pepperweed	Tanweed
Black medic	Dogfennel	Pigweed	Thistle
Black mustard	Dollarweed	Pineywoods bedstraw	Trailing crownvetch
Blackseed plantain	False dandelion	Plains coreopsis	Tumble mustard
Blessed thistle	Falseflax	Plantain	Tumble pigweed
Bloodflower milkweed	False sunflower	Poison ivy	Velvetleaf
Blue lettuce	Fiddleneck	Poison oak	Venice mallow
Blue vervain	Florida pusley	Pokeweed	Virginia buttonweed
Bracted plantain	Frenchweed	Poorjoe	Virginia creeper
Bristly oxtongue	Galinsoga	Prairie sunflower	Virginia pepperweed
Broadleaf dock	Goldenrod	Prickly lettuce	Wavyleaf bullthistle
Broadleaf plantain	Ground ivy	Prickly sida	Western clematis
Broomweed	Gumweed	Prostrate knotweed	Western salsify
Buckhorn	Hairy fleabane	Prostrate pigweed	White mustard
Buckhorn plantain	Hawkweed	Prostrate spurge	Wild aster
Bulbous buttercup	Healall	Prostrate vervain	Wild buckwheat
Bull thistle	Heartleaf drymary	Puncturevine	Wild carrot
Bullnettle	Heath aster	Purslane, common	Wild four-o'clock
Burclover	Hedge bindweed	Ragweed	Wild garlic
Burdock	Hedge mustard	Red sorrel	Wild geranium
Burning nettle	Henbit	Redroot pigweed	Wild lettuce
Burweed	Hoary cress	Redstem filaree	Wild marigold
Buttercup	Hoary plantain	Rough cinquefoil	Wild mustard
Buttonweed	Hoary vervain	Rough fleabane	Wild onion
Canada thistle	Horsenettle	Russian pigweed	Wild parsnip
Carolina geranium	Jimsonweed	Russian thistle	Wild radish
Carpetweed	Knawel	Scarlet pimpernel	Wild rape
Catchweed bedstraw	Knotweed	Scotch thistle	Wild strawberry
Catnip	Kochia	Sheep sorrel	Wild sweet potato
Catsear	Lambsquarters	Shepherdspurse	Wild vetch
Chickweed, common	Lespedeza	Slender plantain	Woodsorrel
Chickweed, mouseear	Mallow	Smallflower galinsoga	Woolly croton
Chicory	Matchweed	Smooth dock	Woolly morningglory
Cinquefoil	Mexicanweed	Smooth pigweed	Woolly plantain
Clover, crimson	Milk vetch	Sorrel	Wormseed
Clover, hop	Morningglory	Sowthistle	Yarrow
Clover, red	Mouseear hawkweed	Spanishneedles	Yellow rocket
Clover, strawberry	Mugwort	Speedwell	Yellowflower pepperweed
Clover, sweet	Musk thistle	Spiny amaranth	and other broadleaf weeds

11/23

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

— END OF OPTION 1 —

OPTION 2: OPTIONAL USE DIRECTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until dust has settled.

INSTRUCTIONS:

Gordon's Trimec® Weed & Feed 33½ controls a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a 24-4-8 fertilizer provides a supply of plant nutrients.

KEYS FOR SUCCESS	CALIBRATION																
◆ Use when weeds are young and actively growing.	Measure your lawn. For example 20' X 50' equals 1,000 square feet.																
◆ Mow lawn to normal height 1 to 2 days before application. Water thoroughly before application to sustain moisture until the next watering.	Make sure your spreader is dry. If spreader is not listed below, adjust to apply 1 pound of product per 280 square feet (28' X 10"). 18 pounds will cover 5,000 square feet.																
◆ Apply preferably in the morning, when grass is moist from dew, or sprinkle lightly. Water will hold the particles on the foliage and prevent dust drift on to nontarget plants. Apply when air is calm. During application, close spreader openings when turning.	<p>The spreader settings are only approximate.</p> <p><i>Drop Spreader Settings:</i></p> <table style="width: 100%; border: none;"> <tr> <td>Republic EZ Drop/</td> <td style="text-align: right;">4¾</td> </tr> <tr> <td>Scotts</td> <td style="text-align: right;">5¾</td> </tr> <tr> <td>Sears, K-Gro</td> <td style="text-align: right;">4¾</td> </tr> <tr> <td>True Temper</td> <td style="text-align: right;">3¾</td> </tr> <tr> <td>Red Devil</td> <td style="text-align: right;">8½</td> </tr> </table> <p>Old or rusty spreaders will discharge products at different rates.</p>	Republic EZ Drop/	4¾	Scotts	5¾	Sears, K-Gro	4¾	True Temper	3¾	Red Devil	8½						
Republic EZ Drop/	4¾																
Scotts	5¾																
Sears, K-Gro	4¾																
True Temper	3¾																
Red Devil	8½																
◆ Do not water treated area for 1 to 2 days after application. Then water thoroughly.	<p>The spreader settings are only approximate.</p> <p><i>Rotary Spreader Settings</i></p> <table style="width: 100%; border: none;"> <tr> <td>Republic EZ Rotary</td> <td style="text-align: right;">2¾</td> </tr> <tr> <td>Scotts Easy Green</td> <td style="text-align: right;">26</td> </tr> <tr> <td>Scotts Speedy Green</td> <td style="text-align: right;">5½</td> </tr> <tr> <td>Republic EZ Grow</td> <td style="text-align: right;">5¼</td> </tr> <tr> <td>Cyclone, True Temper</td> <td style="text-align: right;">3¾</td> </tr> <tr> <td>Red Devil</td> <td style="text-align: right;">5</td> </tr> <tr> <td>Sears, K-Gro</td> <td style="text-align: right;">2¾</td> </tr> <tr> <td>Ortho Whirlybird</td> <td style="text-align: right;">2½</td> </tr> </table>	Republic EZ Rotary	2¾	Scotts Easy Green	26	Scotts Speedy Green	5½	Republic EZ Grow	5¼	Cyclone, True Temper	3¾	Red Devil	5	Sears, K-Gro	2¾	Ortho Whirlybird	2½
Republic EZ Rotary	2¾																
Scotts Easy Green	26																
Scotts Speedy Green	5½																
Republic EZ Grow	5¼																
Cyclone, True Temper	3¾																
Red Devil	5																
Sears, K-Gro	2¾																
Ortho Whirlybird	2½																
◆ Use Precautions: Do not use on desirable clovers, dichondra, carpetgrass, closely mowed bentgrass or St. Augustinegrass. Do not use or allow the material to wind drift onto flowers, vegetables or ornamental shrubbery. Do not apply in areas over root systems of desirable trees or shrubs. Do not apply to newly seeded lawns until after the third mowing.																	

If necessary, reapply in approximately 30 days, if lawn is heavily overrun by weeds, or if adverse low moisture conditions prevail.

The maximum application rate to turf is 1.2 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

- BROADLEAF WEEDS -

Annual yellow sweetclover	Clover, white	Mustard	Spiny cocklebur
Aster	Cockle	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Cocklebur	Narrowleaf vetch	Spotted catsear
Bedstraw	Common mullein	Nettle	Spotted spurge
Beggarticks	Creeping jenny	Orange hawkweed	Spurweed
Betony, Florida	Cudweed	Oriental cocklebur	Stinging nettle
Bindweed, field	Curly dock	Oxalis	Strawberry, India mock
Bird vetch	Daisy, English	Parsley-piert	Tall nettle
Bitter wintercress	Daisy fleabane	Parsnip	Tall vervain
Bittercress, hairy	Daisy, oxeye	Pearlwort	Tansy ragwort
Bitterweed	Dandelion	Pennycress	Tansy mustard
Black-eyed Susan	Dichondra	Pepperweed	Tanweed
Black medic	Dogfennel	Pigweed	Thistle
Black mustard	Dollarweed	Pineywoods bedstraw	Trailing crownvetch
Blackseed plantain	False dandelion	Plains coreopsis	Tumble mustard
Blessed thistle	Falseflax	Plantain	Tumble pigweed
Bloodflower milkweed	False sunflower	Poison ivy	Velvetleaf
Blue lettuce	Fiddleneck	Poison oak	Venice mallow
Blue vervain	Florida pusley	Pokeweed	Virginia buttonweed
Bracted plantain	Frenchweed	Poorjoe	Virginia creeper
Bristly oxtongue	Galinsoga	Prairie sunflower	Virginia pepperweed
Broadleaf dock	Goldenrod	Prickly lettuce	Wavyleaf bullthistle
Broadleaf plantain	Ground ivy	Prickly sida	Western clematis
Broomweed	Gumweed	Prostrate knotweed	Western salsify
Buckhorn	Hairy fleabane	Prostrate pigweed	White mustard
Buckhorn plantain	Hawkweed	Prostrate spurge	Wild aster
Bulbous buttercup	Healall	Prostrate vervain	Wild buckwheat
Bull thistle	Heartleaf drymary	Puncturevine	Wild carrot
Bullnettle	Heath aster	Purslane, common	Wild four-o'clock
Burclover	Hedge bindweed	Ragweed	Wild garlic
Burdock	Hedge mustard	Red sorrel	Wild geranium
Burning nettle	Henbit	Redroot pigweed	Wild lettuce
Burweed	Hoary cress	Redstem filaree	Wild marigold
Buttercup	Hoary plantain	Rough cinquefoil	Wild mustard
Buttonweed	Hoary vervain	Rough fleabane	Wild onion
Canada thistle	Horsenettle	Russian pigweed	Wild parsnip
Carolina geranium	Jimsonweed	Russian thistle	Wild radish
Carpetweed	Knawel	Scarlet pimpernel	Wild rape
Catchweed bedstraw	Knotweed	Scotch thistle	Wild strawberry
Catnip	Kochia	Sheep sorrel	Wild sweet potato
Catsear	Lambsquarters	Shepherdspurse	Wild vetch
Chickweed, common	Lespedeza	Slender plantain	Woodsorrel
Chickweed, mouseear	Mallow	Smallflower galinsoga	Woolly croton
Chicory	Matchweed	Smooth dock	Woolly morningglory
Cinquefoil	Mexicanweed	Smooth pigweed	Woolly plantain
Clover, crimson	Milk vetch	Sorrel	Wormseed
Clover, hop	Morningglory	Sowthistle	Yarrow
Clover, red	Mouseear hawkweed	Spanishneedles	Yellow rocket
Clover, strawberry	Mugwort	Speedwell	Yellowflower pepperweed
Clover, sweet	Musk thistle	Spiny amaranth	and other broadleaf weeds

24-4-8 GUARANTEED ANALYSIS	
Total Nitrogen (N).....	24%
Water Insoluble Nitrogen.....	5%
Available Phosphate (P ₂ O ₅).....	4%
Soluble Potash (K ₂ O).....	8%

Fertilizer Labeling Requirements:

- All fertilizer products distributed in Washington at a minimum, must have the following labeling statement: "This product has been registered with the Washington State Department of Agriculture. When applied as directed, this fertilizer meets the Washington standards for arsenic, cadmium, cobalt, mercury, molybdenum, lead, nickel, selenium, and zinc. You have the right to receive specific information about Washington standards from the distributor of this product."
- After July 1, 1999, the label must also state: "Information received by the Washington State Department of Agriculture regarding the components in this product is available on the internet at <http://www.wa.gov/agr/>."
- Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.xxxxx.com>.

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

CONTAINER DISPOSAL: **If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partially filled:** Call your local solid waste agency or toll free number (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

----- **END OF OPTION 2** -----

OPTION 3: OPTIONAL USE DIRECTION FOR THE SCOTTS COMPANY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

CONTAINER DISPOSAL: **If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partially filled:** Call your local solid waste agency or toll free number (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.

USE PRECAUTIONS:

Do not use on dichondra, carpetgrass or St. Augustinegrass.

Do not use this product where desirable clovers are present or allow the material to drift onto flowers, vegetables or ornamental shrubbery. Do not apply in areas underlaid by roots of desirable trees or shrubs. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning.

INSTRUCTIONS:

Gordon's Trimec® Weed & Feed 33½ is a Trimec® complex, a combination of three proven weed killers -- 2,4-D, Mecoprop-p and Dicamba. Together, they control a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a 24-4-8 fertilizer provides a supply of plant nutrients. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth.

Apply 2.4 to 3.6 pounds of product per 1,000 square feet of turfgrass. A second application may be made in approximately 30 days.

The maximum application rate to turf is 1.2 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

HOW AND WHEN TO APPLY:

Apply when broadleaf weeds are actively growing in spring or early fall.

WHAT TO EXPECT:

Weed stems and leaves will begin to twist and curl in a few days, but a month or more may be required for complete control. This product will not prevent weed seeds from producing new plants.

- Mow lawn to normal height 1 to 2 days before application.
- Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see 4th bullet).
- Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants
- Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.
- Avoid applying when temperatures are consistently above 90°F.

16/23

Note to Supplemental Registrants: Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, equipment settings of drop spreaders, rotary spreaders, and other granular applicators may be presented on the container labels.

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.
12	2.4	5,000
	2.6	4,600
	2.8	4,300
	3.0	4,000
	3.2	3,800
	3.4	-----
	3.6	-----
13	2.4	5,400
	2.6	5,000
	2.8	4,600
	3.0	4,300
	3.2	4,100
	3.4	3,800
	3.6	3,600
14	2.4	5,800
	2.6	5,400
	2.8	5,000
	3.0	4,700
	3.2	4,400
	3.4	4,100
	3.6	3,900
15	2.4	6,300
	2.6	5,800
	2.8	5,400
	3.0	5,000
	3.2	4,700
	3.4	4,400
	3.6	4,200
16	2.4	6,700
	2.6	6,200
	2.8	5,700
	3.0	5,300
	3.2	5,000
	3.4	4,700
	3.6	4,400
17	2.4	7,100
	2.6	6,500
	2.8	6,100
	3.0	5,700
	3.2	5,300
	3.4	5,000
	3.6	4,700
18	2.4	7,500
	2.6	6,900
	2.8	6,400
	3.0	6,000
	3.2	5,600
	3.4	5,300
	3.6	5,000

17/23

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.
24	2.4	10,000
	2.6	9,200
	2.8	8,600
	3.0	8,000
	3.2	7,500
	3.4	7,100
	3.6	6,700
25	2.4	10,400
	2.6	9,600
	2.8	8,900
	3.0	8,300
	3.2	7,800
	3.4	7,400
	3.6	6,900
26	2.4	10,800
	2.6	10,000
	2.8	9,300
	3.0	8,700
	3.2	8,100
	3.4	7,600
	3.6	7,200
28	2.4	11,700
	2.6	10,800
	2.8	10,000
	3.0	9,300
	3.2	8,800
	3.4	8,200
	3.6	7,800
36	2.4	15,000
	2.6	13,800
	2.8	12,900
	3.0	12,000
	3.2	11,300
	3.4	10,500
	3.6	10,000
37	2.4	15,400
	2.6	14,200
	2.8	13,200
	3.0	12,300
	3.2	11,600
	3.4	10,900
	3.6	10,300
39	2.4	16,300
	2.6	15,000
	2.8	13,900
	3.0	13,000
	3.2	12,200
	3.4	11,500
	3.6	10,800
42	2.4	17,500
	2.6	16,200
	2.8	15,000
	3.0	14,000
	3.2	13,100
	3.4	12,400
	3.6	11,700

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.
43	2.4	17,900
	2.6	16,500
	2.8	15,400
	3.0	14,300
	3.2	13,400
	3.4	12,600
	3.6	11,900
48	2.4	20,000
	2.6	18,500
	2.8	17,100
	3.0	16,000
	3.2	15,000
	3.4	14,100
	3.6	13,300
49	2.4	20,400
	2.6	18,800
	2.8	17,500
	3.0	16,300
	3.2	15,300
	3.4	14,400
	3.6	13,600

SPREADING TIPS:

To provide proper distribution, calibrate spreader before application.

For best results, use a Scotts® Spreader.

Spreader Settings		DIRECTIONS
Scotts AccuGreen® Drop Spreaders	X	Mow lawn first: apply 2 header strips; apply longest direction of lawn, overlapping wheel marks.
Scotts PrecisionGreen® Drop Spreader	X x/x	Mow lawn first: apply 2 header strips; apply longest direction of lawn, overlapping wheel marks.
Scotts SpeedyGreen® & EdgeGuard® Rotary Spreaders	X	Make each pass about 5 feet from previous pass to overlap spread width.
Scotts EasyGreen® Rotary	X	Make each pass about 30 inches from previous pass to overlap spread width.

[Optional Direction Illustrations May Be Presented]

Additional Spreader Settings:

- EZ Drop, Republic (# 530)..... X
- EZ Broadcast, Republic (# 540)..... X x/x
- Scotts Broadcast, Tractor Spreader..... X

(Non-Scotts spreader settings are approximate. You may need to adjust for proper coverage. Check your spreader on xxx sq.ft. and adjust to apply at the rate of 1 pound of product per xxx sq.ft.)

[Note to Supplemental Registrants: Refer to table for appropriate bag sizes.]

- For 2.4 lbs. of product per 1,000 sq.ft. of turfgrass: Check your spreader on 420 sq.ft. and adjust to apply at the rate of 1 lb. of product per 420 sq.ft.
- For 2.6 lbs. of product per 1,000 sq.ft. of turfgrass: Check your spreader on 385 sq.ft. and adjust to apply at the rate of 1 lb. of product per 385 sq.ft.
- For 2.8 lbs. of product per 1,000 sq.ft. of turfgrass: Check your spreader on 360 sq.ft. and adjust to

- apply at the rate of 1 lb. of product per 360 sq.ft.
- For 2.8 lbs. of product per 1,000 sq.ft. of turfgrass: Check your spreader on 360 sq.ft. and adjust to apply at the rate of 1 lb. of product per 360 sq.ft.
- For 3.0 lbs. of product per 1,000 sq.ft. of turfgrass: Check your spreader on 333 sq.ft. and adjust to apply at the rate of 1 lb. of product per 333 sq.ft.
- For 3.2 lbs. of product per 1,000 sq.ft. of turfgrass: Check your spreader on 310 sq.ft. and adjust to apply at the rate of 1 lb. of product per 310 sq.ft.
- For 3.4 lbs. of product per 1,000 sq.ft. of turfgrass: Check your spreader on 300 sq.ft. and adjust to apply at the rate of 1 lb. of product per 300 sq.ft.
- For 3.6 lbs. of product per 1,000 sq.ft. of turfgrass: Check your spreader on 275 sq.ft. and adjust to apply at the rate of 1 lb. of product per 275 sq.ft.

GUARANTEED ANALYSIS XX-X-X	
Total Nitrogen (N)	XX%
x.x% ammoniacal nitrogen	
x.x% urea nitrogen*	
Available Phosphate (P ₂ O ₅)	X%
Soluble Potash (K ₂ O)	X%
Sulfur (S)	X%
x.x% free sulfur (S)	
Derived from: polymer-coated sulfur-coated urea, urea, ammonium phosphate, potassium chloride.	
*A portion of the urea has been coated to provide x.x% coated slow-release nitrogen.	
This product and/or its manufacture is covered by one or more of the following U.S. patents and international counterparts thereof: US Patent Nos: 5,219,465; 5,300,135; 5,405,426; 5,466,274.	

Fertilizer Labeling Requirements:

- All fertilizer products distributed in Washington at a minimum, must have the following labeling statement: "This product has been registered with the Washington State Department of Agriculture. When applied as directed, this fertilizer meets the Washington standards for arsenic, cadmium, cobalt, mercury, molybdenum, lead, nickel, selenium, and zinc. You have the right to receive specific information about Washington standards from the distributor of this product."
- After July 1, 1999, the label must also state: "Information received by the Washington State Department of Agriculture regarding the components in this product is available on the internet at <http://www.wa.gov/agr/>."
- Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.xxxxx.com>.

- BROADLEAF WEEDS -

Annual yellow sweetclover	Clover, white	Mustard	Spiny cocklebur
Aster	Cockle	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Cocklebur	Narrowleaf vetch	Spotted catsear
Bedstraw	Common mullein	Nettle	Spotted spurge
Beggarticks	Creeping jenny	Orange hawkweed	Spurweed
Betony, Florida	Cudweed	Oriental cocklebur	Stinging nettle
Bindweed, field	Curly dock	Oxalis	Strawberry, India mock
Bird vetch	Daisy, English	Parsley-piert	Tall nettle
Bitter wintercress	Daisy fleabane	Parsnip	Tall vervain
Bittercress, hairy	Daisy, oxeye	Pearlwort	Tansy ragwort
Bitterweed	Dandelion	Pennycress	Tansy mustard
Black-eyed Susan	Dichondra	Pepperweed	Tanweed
Black medic	Dogfennel	Pigweed	Thistle
Black mustard	Dollarweed	Pineywoods bedstraw	Trailing crownvetch
Blackseed plantain	False dandelion	Plains coreopsis	Tumble mustard
Blessed thistle	Falseflax	Plantain	Tumble pigweed
Bloodflower milkweed	False sunflower	Poison ivy	Velvetleaf
Blue lettuce	Fiddleneck	Poison oak	Venice mallow
Blue vervain	Florida pusley	Pokeweed	Virginia buttonweed
Bracted plantain	Frenchweed	Poorjoe	Virginia creeper
Bristly oxtongue	Galinsoga	Prairie sunflower	Virginia pepperweed
Broadleaf dock	Goldenrod	Prickly lettuce	Wavyleaf bullthistle
Broadleaf plantain	Ground ivy	Prickly sida	Western clematis
Broomweed	Gumweed	Prostrate knotweed	Western salsify
Buckhorn	Hairy fleabane	Prostrate pigweed	White mustard
Buckhorn plantain	Hawkweed	Prostrate spurge	Wild aster
Bulbous buttercup	Healall	Prostrate vervain	Wild buckwheat
Bull thistle	Heartleaf drymary	Puncturevine	Wild carrot
Bullnettle	Heath aster	Purslane, common	Wild four-o'clock
Burclover	Hedge bindweed	Ragweed	Wild garlic
Burdock	Hedge mustard	Red sorrel	Wild geranium
Burning nettle	Henbit	Redroot pigweed	Wild lettuce
Burweed	Hoary cress	Redstem filaree	Wild marigold
Buttercup	Hoary plantain	Rough cinquefoil	Wild mustard
Buttonweed	Hoary vervain	Rough fleabane	Wild onion
Canada thistle	Horsenettle	Russian pigweed	Wild parsnip
Carolina geranium	Jimsonweed	Russian thistle	Wild radish
Carpetweed	Knawel	Scarlet pimpernel	Wild rape
Catchweed bedstraw	Knotweed	Scotch thistle	Wild strawberry
Catnip	Kochia	Sheep sorrel	Wild sweet potato
Catsear	Lambsquarters	Shepherdspurse	Wild vetch
Chickweed, common	Lespedeza	Slender plantain	Woodsorrel
Chickweed, mouseear	Mallow	Smallflower galinsoga	Woolly croton
Chicory	Matchweed	Smooth dock	Woolly morningglory
Cinquefoil	Mexicanweed	Smooth pigweed	Woolly plantain
Clover, crimson	Milk vetch	Sorrel	Wormseed
Clover, hop	Morningglory	Sowthistle	Yarrow
Clover, red	Mouseear hawkweed	Spanishneedles	Yellow rocket
Clover, strawberry	Mugwort	Speedwell	Yellowflower pepperweed
Clover, sweet	Musk thistle	Spiny amaranth	and other broadleaf weeds

----- END OPTION 3 -----

APPENDIX

I. Container Packaging and Size:

Approved retail container package sizes include: 12, 13, 14, 15, 16, 17, 18, 24, 25, 26, 28, 34, 36, 37, 39, 42, 43, 48, and 49 pounds.

II. Previously Approved Fertilizer Grades: Fertilizer grades with nutrient contents expressed as the percentages of N, P₂O₅, and K₂O include the following:

19-6-8	23-3-7	26-4-4	30-4-4
22-3-3	25-3-3	26-4-6	32-3-3
22-3-6	25-4-8	28-0-3	32-4-8
22-3-17	26-3-6	29-3-4	24-4-8
22-4-11		30-3-3	

The fertilizer grade guaranteed analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.

23-3-7 GUARANTEED ANALYSIS	
Total Nitrogen (N).....	23%
Water Insoluble Nitrogen.....	5%
Available Phosphate (P ₂ O ₅).....	3%
Soluble Potash (K ₂ O).....	7%

28-0-3 GUARANTEED ANALYSIS	
Total Nitrogen (N).....	28%
28% urea nitrogen*.....	
Available Phosphate (P ₂ O ₅).....	0%
Soluble Potash (K ₂ O).....	3%
Sulfur (S) (Free).....	4%
Derived from: polymer-coated sulfur-coated urea, urea and potassium chloride.	
*A portion of the urea has been coated to provide 9.6% coated slow-release nitrogen.	

22-4-11 GUARANTEED ANALYSIS	
Total Nitrogen (N).....	22%
1.6% ammoniacal nitrogen	
20.4% urea nitrogen*	
Available Phosphate (P ₂ O ₅).....	4%
Soluble Potash (K ₂ O).....	11%
Sulfur (S) (Free).....	2.2%
Derived from: sulfur-coated polymer-coated urea, urea, ammonium phosphate, potassium chloride.	
*A portion of the urea has been coated to provide 5.5% coated slow-release nitrogen.	

III. The addition of advertising claims by supplemental registrants that may be presented on the retail containers:

- (Number) pounds covers up to (Number) square feet.
- Controls (Kills) (desired weed species from attached listing may be inserted here e.g. dandelion, chickweed, knotweed, plantain, henbit, spurge) and many other broadleaf weeds including those listed on the back panel.
- Highly effective pre-emergent control of crabgrass and other grassy weeds.
- Long lasting, lush green lawn.
- Stimulates deep greening.
- Stimulates quick, deep greening.
- Stimulates early, vigorous root growth.
- Stimulates thick, lush grass blades.
- Thickens and greens lawns.
- Ortho product quality you've come to trust.
- One-step application for a lush green lawn.
- Fast deep greening balanced with long-term feeding by patented Polyon® coated fertilizer technology.
- Small particle size for uniform coverage to control a wide range of weeds.
- Patented Polyon® coated fertilizer technology for timed release of fertilizer.
- Polyon® coating prevents burning.
- Scientific blend of small particles promotes fast deep greening and balanced long term feeding.
- Outdoor use only
- Satisfaction guaranteed* or your purchase price refunded with proof-of-purchase and original cash register receipt.
- Controls (Kills) broadleaf lawn weeds.
- Controls (Kills) (Number) common lawn weeds including dandelion and clover.
- Feeds, greens and thickens lawns while controlling weeds.
- Poly-S® Technology.
- Scotts® slow-release polymer-encapsulated nitrogen fertilizer for uniform growth with extended feeding.
- Provides a regular turf feeding.
- Feeds up to 8 weeks.
- These products are the quality choice in complete lawn care. You determine what your lawn needs, and you'll find a product to help you. Fertilizers, lawn starters, crabgrass prevention, weed and insect control - everything for a lush green lawn and year-round protection. Each one is easy to apply and guaranteed to produce quick, long-lasting results.
- This product is designed to be used in the spring, when weeds are actively growing.
- Apply anytime weeds are actively growing.
- The fertilizer combination of nitrogen, phosphate, and potash gives your lawn strength above and below the surface. This product promotes root and rhizome development, helping the grass to produce more blades. The result is a thicker, greener lawn.
- Phosphorous-Free Formulation
- Scotts® Phosphorous-Free fertilizer provides the best formula for soils already containing a high natural presence of available phosphorous. A soil test can help determine if your soil will support normal turf growth without a phosphorous supplement.
 - *Nitrogen for Color*—Contributes to thicker, greener growth.
 - *Phosphorous for strength*—Stimulates root growth. Test your soil regularly to see if phosphorous should be applied.
 - *Potassium for Protection*—Protects turf against disease and drought damage.
- Scotts Phosphorous-Free products are specifically formulated with Scotts Poly-S® timed-release fertilizer. The unique Poly-S® coating provides your lawn with uniform growth and an extended feeding for up to two months.
- With the right weather conditions and correct application, your lawn will be greener and thin areas of turf will fill in, making the density of your lawn more uniform.
- An easy guide to timing your application is to look for dandelions on your lawn.

- This product can be applied in the fall when weeds are actively growing. Because plants are taking nutrients into their roots at this time of year, it's easier to control some weeds in the fall than in the spring.
- This product makes it easier to have a lush, green lawn. Our four (4) simple applications eliminate the guesswork and give you professional lawn care results. Each stage combines just the right amount of fertilizer with the right amount of control and prevention. Just match each stage to the correct season and you'll prevent crabgrass, control weeds and insects, and prepare your lawn for the winter. It's what your lawn needs for strong growth.
 - Stage One (February—April) is a pre-emergent -- *Scotts Phosphorous-Free fertilizer with Crabgrass Preventer*
 - Stage Two is a broadleaf postemergent -- *Scotts Phosphorous-Free fertilizer with Weed Control*
 - Stage Three is an insecticide.
 - Stage Four is a fall broadleaf postemergent.
- Specially formulated with Scotts Poly S[®] timed-release fertilizer [as used by landscape professionals.]
- Specially formulated with Scotts[®] Poly-S[®] timed-release fertilizer to provide uniform growth and extended feeding.
- Weed control penetrates down to the root to kill dandelion, clover and other broadleaf weeds.
- Contains 40% of total guarantee of nitrogen as controlled release nitrogen for uniform growth and extended greening.
- (Number) pounds treats and feeds (Number) square feet of turfgrass.
- For best results, [always] use a Scotts[®] spreader.
- Visit our website www.xxxxx.com.
- Scotts No Quibble Guarantee™. If for any reason you are not satisfied with results after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.
- Answers to FAQ's on the internet at www.xxxxx.com
- Feeds up to 2 months.
- Contains Nutralene[®] Controlled Release Nitrogen.
- Produce superior results ... every time.
- Nutralene is a controlled release homogenous proven technology. It releases nitrogen gradually through hydrolysis and microbial action in the soil for a sustained feeding. Nutralene is unlike other nitrogen sources that depend on coatings, particle size, and temperature for their nitrogen release.
- The ultimate timed-released fertilizer as used by professional golf courses.
- Contains both readily available nitrogen for quick green up and controlled release nitrogen for uniform growth and extended feeding.
- Treats and feeds ____ sq.ft.
- Proudly produced in the U.S.A.
- Can we help you? Call toll-free: 1-XXX-XXX-XXXX or visit our website at www.XXXXXXXX.com.
- Clears out dandelions and other broadleaf weeds.
- Feeds, greens and thickens lawns.
- Apply in late fall.
- Feeds & treats 5,000 sq. ft.
- ~~Knocks out weeds.~~

IV. Guarantees offered by supplemental registrants may include the following:

- CAN'T FAIL GUARANTEE. If you are not completely satisfied with the results after using this product, (Company Name) will refund your purchase price in full. Simply send proof of purchase and register receipt to (Company Name and Address).

V. Addition of a referral claim:

- To spot treat individual weeds, use Ortho Weed-B-Gon Weed Killer[®].