



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

May 17, 2017

Eric D. Smith
PBI/Gordon Corporation
P.O. Box 014090
Kansas City, Missouri 64101

Subject: Notification per PRN 98-10 – Minor Label changes and adding advertising statements
Product Name: GORDON'S TRIMEC WEED & FEED 33 1/2
EPA Registration Number: 2217-660
Application Date: January 13, 2017
Decision Number: 525421

Dear Mr. Smith:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped "Notification" and will be placed in our records.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If you have any questions, you may contact Grant Rowland at 703-347-0254 or via email at rowland.grant@epa.gov

Sincerely,

A handwritten signature in black ink, appearing to read "Kathryn V. Montague for".

Kathryn V. Montague, Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

GORDON'S TRIMEC® WEED & FEED 33½

EPA Reg. No. 2217-660

Three sublabels represent the entire master label of Gordon's Trimec Weed & Feed 33 1/2. Please refer to the appropriate section of the labeling as shown as follows:

Section
Language Common to All Sublabels
Sublabel 1
Sublabel 2
Sublabel 3
Appendix
Document Control Information

Language Common to All Sublabels

GORDON'S TRIMEC® WEED & FEED 33½

EPA Reg. No. 2217-660

ACTIVE INGREDIENTS:

2,4-D	0.76%
Mecoprop-p (MCP-p)	0.17%
Dicamba	0.07%
OTHER INGREDIENTS:	<u>99.00%</u>
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- 15.2 lb 2,4-dichlorophenoxyacetic acid equivalent per ton
- 3.4 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton
- 1.4 lb 3,6-dichloro-o-anisic acid equivalent per ton

Isomer Specific by AOAC Methods

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

CAS Registry Numbers: 2,4-D: 94-75-7, Mecoprop-p: 16484-77-8, Dicamba: 1918-00-9

KEEP OUT OF REACH OF CHILDREN

CAUTION



NOTIFICATION

2217-660

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

05/17/17

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

See attached booklet for complete [Precautionary Statements and] Directions for Use, including Storage and Disposal, and back panel for [complete Precautionary Statements, including] First Aid. OR See back panel for complete Precautionary Statements, including First Aid, and Directions for Use, including Storage and Disposal.

Net Contents: _____
EPA Est. No. _____

Company Name and Address:

PBI/Gordon Corporation
1217 West 12th Street
Kansas City, Missouri 64101



PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves made of any waterproof material. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid	
If on skin or on clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 - 20 minutes.• Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferable mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-xxx-xxxx for emergency medical treatment information.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

[Environmental Hazards (alternate per PRN 2008-1)]

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.]

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

SUBLABEL 1

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Do not use on dichondra, carpetgrass or St. Augustinegrass.
- Do not use this product where desirable clovers are present. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning.
- Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

INSTRUCTIONS:

Gordon's Trimec® Weed & Feed 33½ is a Trimec® complex, a combination of three proven weed killers - 2,4-D, Mecoprop-p and Dicamba. Together, they control a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others **as listed which are** commonly found in home lawns. In addition, a 24-4-8 fertilizer provides a supply of plant nutrients. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth.

Apply 2.4 to 3.6 **pounds lb** of product per 1,000 **square feet sq.ft.** of turfgrass. A second application may be made in 30 days.

The maximum broadcast application rate is 3.6 **pounds lb** of product per 1,000 sq.ft. per application [1.2 lb 2,4-D ae, 0.27 lb MCP-p ae and 0.11lb dicamba ae per acre per application]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 7.2 **pounds lb** of product per 1,000 sq.ft. [2.4 lb 2,4-D ae, 0.54 lb MCP-p ae and 0.22 lb dicamba ae per acre].

WHEN TO APPLY:

Apply when broadleaf weeds are actively growing in spring or early fall.

WHAT TO EXPECT:

Weed stems and leaves will begin to twist and curl in a few days, but a month or more may be required for complete control. This product will not prevent weed seeds from producing new plants.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
4. Avoid applying when temperatures are consistently above 90°F.
5. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

BROADLEAF WEEDS			
Annual yellow sweetclover	Clover, white	Mustard	Spiny cocklebur
Aster	Cockle	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Cocklebur	Narrowleaf vetch	Spotted catsear
Bedstraw	Common mullein	Nettle	Spotted spurge
Beggarticks	Creeping jenny	Orange hawkweed	Spurweed
Betony, Florida	Cudweed	Oriental cocklebur	Stinging nettle
Bindweed, field	Curly dock	Oxalis	Strawberry, India mock
Bird vetch	Daisy, English	Parsley-piert	Tall nettle
Bitter wintercress	Daisy fleabane	Parsnip	Tall vervain
Bittercress, hairy	Daisy, oxeye	Pearlwort	Tansy ragwort
Bitterweed	Dandelion	Pennycress	Tansy mustard
Black-eyed Susan	Dichondra	Pepperweed	Tanweed
Black medic	Dogfennel	Pigweed	Thistle
Black mustard	Dollarweed	Pineywoods bedstraw	Trailing crownvetch
Blackseed plantain	False dandelion	Plains coreopsis	Tumble mustard
Blessed thistle	Falseflax	Plantain	Tumble pigweed
Bloodflower milkweed	False sunflower	Poison ivy	Velvetleaf
Blue lettuce	Fiddleneck	Poison oak	Venice mallow
Blue vervain	Florida pusley	Pokeweed	Virginia buttonweed
Bracted plantain	Frenchweed	Poorjoe	Virginia creeper
Bristly oxtongue	Galinsoga	Prairie sunflower	Virginia pepperweed
Broadleaf dock	Goldenrod	Prickly lettuce	Wavyleaf bullthistle
Broadleaf plantain	Ground ivy	Prickly sida	Western clematis
Broomweed	Gumweed	Prostrate knotweed	Western salsify
Buckhorn	Hairy fleabane	Prostrate pigweed	White mustard
Buckhorn plantain	Hawkweed	Prostrate spurge	Wild aster
Bulbous buttercup	Healall	Prostrate vervain	Wild buckwheat
Bull thistle	Heartleaf drymary	Puncturevine	Wild carrot
Bullnettle	Heath aster	Purslane, common	Wild four-o'clock
Burclover	Hedge bindweed	Ragweed	Wild garlic
Burdock	Hedge mustard	Red sorrel	Wild geranium
Burning nettle	Henbit	Redroot pigweed	Wild lettuce
Burweed	Hoary cress	Redstem filaree	Wild marigold
Buttercup	Hoary plantain	Rough cinquefoil	Wild mustard
Buttonweed	Hoary vervain	Rough fleabane	Wild onion
Canada thistle	Horsenettle	Russian pigweed	Wild parsnip
Carolina geranium	Jimsonweed	Russian thistle	Wild radish
Carpetweed	Knawel	Scarlet pimpernel	Wild rape
Catchweed bedstraw	Knotweed	Scotch thistle	Wild strawberry
Catnip	Kochia	Sheep sorrel	Wild sweet potato
Catsear	Lambsquarters	Shepherdspurse	Wild vetch
Chickweed, common	Lespedeza	Slender plantain	Woodsorrel
Chickweed, mouseear	Mallow	Smallflower galinsoga	Woolly croton
Chicory	Matchweed	Smooth dock	Woolly morningglory
Cinquefoil	Mexicanweed	Smooth pigweed	Woolly plantain
Clover, crimson	Milk vetch	Sorrel	Wormseed

BROADLEAF WEEDS			
Clover, hop	Morningglory	Sowthistle	Yarrow
Clover, red	Mouseear hawkweed	Spanishneedles	Yellow rocket
Clover, strawberry	Mugwort	Speedwell	Yellowflower pepperweed
Clover, sweet	Musk thistle	Spiny amaranth	and other broadleaf weeds

Fertilizer Labeling Requirements:

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.xxxxx.com>.

[Begin Optional Language]

[Note to Supplemental Registrants: Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, equipment settings of drop spreaders, rotary spreaders, and other granular applicators may be presented on the container labels.]

CALIBRATION INSTRUCTIONS:

To provide proper distribution, calibrate spreader before application. [Refer to table for appropriate bag sizes.]

- For 2.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 420 square feet and adjust to apply at the rate of 1 pound of product per 420 square feet.
- For 2.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 385 square feet and adjust to apply at the rate of 1 pound of product per 385 square feet.
- For 2.8 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 360 square feet and adjust to apply at the rate of 1 pound of product per 360 square feet.
- For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 333 square feet and adjust to apply at the rate of 1 pound of product per 333 square feet.
- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.

Spreader settings for 1.0 pound of product per 310 square feet	
Scotts EasyGreen® Rotary	26
Scotts Drop	5
Scotts SpeedyGreen® Rotary	3.5
GreenView Drop (Spred II)	4
Central Drop (#205)	5.5
Sears Drop (#19810)	4
Ortho Drop (#5000)	5
Precision Drop (Red Devil)	4

- For 3.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 300 square feet and adjust to apply at the rate of 1 pound of product per 300 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.

DISTRIBUTION AND SPREADER PATTERNS:

Always apply with a properly calibrated rotary or drop type fertilizer spreader to distribute the granules evenly. Never apply by hand.


Coverage - To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to XXX square feet. [Graphics Optional]

SINGLE PASS DISTRIBUTION:

First spread the borders of your lawn. Then spread the center by spreading in straight line passes. Spread at a uniform walking speed to insure even coverage [Graphics Optional]. To avoid gaps and streaks, first spread the border and fill in center with uniform, parallel passes.


Alternate Spreader Settings:


Spreader Settings



Apply in a single pass at a rate of 3.2 lbs. per 1,000 sq. ft. Start on the outside border and apply in a circular path working toward the center. Overlap the coverage slightly, and shut off spreader before making hard turns.


Broadcast Spreaders	Setting
Agri-Fab® Rotary	5
Cyclone® Rotary	4
Earthway® Rotary	14
Greenview® Jet-Spred® SS1	14
Lesco® Rotary (dial A-S)/(number dial)	H/16
Red Devil® Rotary (1-20)	5
Republic® Rotary	4
Scotts® EasyGreen®	27
Scotts® EdgeGuard®	5
Scotts® SpeedyGreen® 2000, 3000	5
Sears® Craftsman Rotary	5
Sta-Green® Rotary (1-10)	4
True-Temper® Rotary	4
Vigoro® Rotary (1-10)	4





Apply in a single pass at a rate of 3.2 lbs. per 1,000 sq. ft. Start on the outside border and apply in a rectangular path working back and forth across the length of the lawn. Overlap the coverage slightly, and shut off spreader before making hard turns.

Drop Spreaders	Setting
Agri-Fab® Drop	5
Greenview® Drop-Spred® XP1	18
Red Devil® Drop (1-5)	4 1/2
Republic® Drop	6 1/2
Scotts® Drop AccuGreen (2-18)	5
Scotts® Drop PF1 (2-18)	5
Sta-Green® Drop (1-5)	4 1/2



If your spreader model does not appear above, go to our website at www.preen.com, or call toll-free (800) 233-1067, 8 A.M. - 5 P.M. EDT, Monday - Friday, for additional model settings.

[Note: Additional spreader models and settings may be added or substituted by the supplemental registrant provided they deliver the labeled rate of product.]

[Note: Spreader settings may be varied by the supplemental registrant as different grades of fertilizer are used, provided they deliver the labeled rate of pesticide.]

[End Optional Language]

[Advisory For Fertilizers Containing Iron:

Keep this product off sidewalks, patios, driveways, etc. This fertilizer contains iron which will stain concrete and painted surfaces. If it comes in contact with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.]

24-4-8 GUARANTEED ANALYSIS	
Total Nitrogen (N)	24%
Water Insoluble Nitrogen	5%
Available Phosphate (P2O5)	4%
Soluble Potash (K2O)	8%

Referrals to other sequentially used products:

- For best results, apply (brand name) products.
- Apply using the 4-Step Program
- **4-Step Program:**

STEP 1: Menards Premium Crabgrass Preventer - Apply March - early May - Prevents crabgrass, foxtail, goosegrass, poa annua and other annual weed grasses. - Nutralene feeds your turfgrass for up to 2 months with long lasting deep greening.

STEP 2: Menards Premium Weed & Feed - Apply May - June - Kills dandelion, clover, plantain and over XXX other broadleaf weeds. - Nutralene feeds your turfgrass for up to 2 months with long lasting deep greening.

STEP 3: Menards Premium Lawn Food - Apply June - August - Nourishes your lawn throughout the summer months with extended feeding of Nitrogen. - Nutralene feeds your turfgrass for up to 2 months with long lasting deep greening.

STEP 4: Menards Premium Winterizer & Weed Control - Apply late August - mid October - Kills dandelion, clover, plantain and over XXX other broadleaf weeds. - Maximizes root building. - Nutralene feeds your turfgrass for up to 2 months with long lasting deep greening.

OR

STEP 4: Menards Premium Starter & Winterizer - Apply late August - November - Maximizes root building. - Nutralene feeds your turfgrass for up to 2 months with long lasting deep greening. - Use with new seeding and sodding of turfgrass.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or www.website.com for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

SUBLABEL 2: Optional Use Directions

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND RESTRICTIONS:

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

INSTRUCTIONS:

Gordon's Trimec® Weed & Feed 33½ controls a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a 24-4-8 fertilizer provides a supply of plant nutrients.


KEYS FOR SUCCESS	CALIBRATION
<ul style="list-style-type: none">Use when weeds are young and actively growing.	Measure your lawn. For example 20' X 50' equals 1,000 square feet.
<ul style="list-style-type: none">Mow lawn to normal height 1 to 2 days before application. Water thoroughly before application to sustain moisture until the next watering.	Make sure your spreader is dry. If spreader is not listed below, adjust to apply 1 pound of product per 280 square feet (28' X 10'). 18 pounds will cover 5,000 square feet.
<ul style="list-style-type: none">Apply preferably in the morning, when grass is moist from dew, or sprinkle lightly. Water will hold the particles on the foliage. Apply when air is calm. During application, close spreader openings when turning.	The spreader settings are only approximate. Drop Spreader Settings: Republic EZ Drop/ 4.75 Scotts 5.75 Sears, K-Gro 4.75 True Temper 3.75 Red Devil 8.5 Old or rusty spreaders will discharge products at different rates.
<ul style="list-style-type: none">Do not water treated area for 1 to 2 days after application. Then water thoroughly.	The spreader settings are only approximate. Rotary Spreader Settings Republic EZ Rotary 2.75 Scotts Easy Green 26 Scotts Speedy Green 5.5 Republic EZ Grow 5.25 Cyclone, True Temper 3.75 Red Devil 5 Sears, K-Gro 2.75 Ortho Whirlybird 2.5
Use Precautions: Do not use on desirable clovers, dichondra, carpetgrass, bentgrass or St. Augustinegrass. Do not apply to newly seeded lawns until after the third mowing.	
Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result.	
Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.	

If necessary, reapply in 30 days, if lawn is heavily overrun by weeds, or if adverse low moisture conditions prevail.

The maximum broadcast application rate is 3.6 pounds of product per 1,000 sq.ft. per application [1.2 lb 2,4-D ae, 0.27 lb MCP P-p ae and 0.11 lb dicamba ae per acre per application]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 7.2 pounds of product per 1,000 sq.ft. [2.4 lb 2,4-D ae, 0.54 lb MCP P-p ae and 0.22 lb dicamba ae per acre].


Alternate Spreader Settings:


Spreader Settings



Apply in a single pass at a rate of 3.2 lbs. per 1,000 sq. ft. Start on the outside border and apply in a circular path working toward the center. Overlap the coverage slightly, and shut off spreader before making hard turns.


Broadcast Spreaders	Setting
Agri-Fab® Rotary	5
Cyclone® Rotary	4
Earthway® Rotary	14
Greenview® Jet-Spred® SS1	14
Lesco® Rotary (dial A-S)(number dial)	H/16
Red Devil® Rotary (1-20)	5
Republic® Rotary	4
Scotts® EasyGreen®	27
Scotts® EdgeGuard®	5
Scotts® SpeedyGreen® 2000, 3000	5
Sears® Craftsman Rotary	5
Sta-Green® Rotary (1-10)	4
True Temper® Rotary	4
Vigoro® Rotary (1-10)	4





Apply in a single pass at a rate of 3.2 lbs. per 1,000 sq. ft. Start on the outside border and apply in a rectangular path working back and forth across the length of the lawn. Overlap the coverage slightly, and shut off spreader before making hard turns.

Drop Spreaders	Setting
Agri-Fab® Drop	5
Greenview® Drop-Spred® XP1	18
Red Devil® Drop (1-5)	4½
Republic® Drop	6½
Scotts® Drop AccuGreen (2-18)	5
Scotts® Drop PF1 (2-18)	5
Sta-Green® Drop (1-5)	4½



If your spreader model does not appear above, go to our website at www.preen.com, or call toll-free (800) 233-1067, 8 A.M. - 5 P.M. EDT, Monday - Friday, for additional model settings.

[Note: Additional spreader models and settings may be added or substituted by the supplemental registrant provided they deliver the labeled rate of product.]

[Note: Spreader settings may be varied by the supplemental registrant as different grades of fertilizer are used, provided they deliver the labeled rate of pesticide.]

24-4-8 GUARANTEED ANALYSIS	
Total Nitrogen (N)	24%
Water Insoluble Nitrogen	5%
Available Phosphate (P ₂ O ₅)	4%
Soluble Potash (K ₂ O)	8%

9 of 19
002217-00660.20170113.notif-proposed-highlighted

SUBLABEL 3: Optional Use Directions for the Scotts Company

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Do not use on dichondra, carpetgrass or St. Augustinegrass.
- Do not use this product where desirable clovers are present. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning.
- Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

INSTRUCTIONS:

Gordon's Trimec® Weed & Feed 33½ is a Trimec® complex, a combination of three proven weed killers - 2,4-D, Mecoprop-p and Dicamba. Together, they control a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a 24-4-8 fertilizer provides a supply of plant nutrients. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth.

Apply 2.4 to 3.6 pounds of product per 1,000 square feet of turfgrass. A second application may be made in 30 days.

The maximum broadcast application rate is 3.6 pounds of product per 1,000 sq.ft. per application [1.2 lb 2,4-D ae, 0.27 lb MCP-P ae and 0.11 lb dicamba ae per acre per application]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 7.2 pounds of product per 1,000 sq.ft. [2.4 lb 2,4-D ae, 0.54 lb MCP-P ae and 0.22 lb dicamba ae per acre].

HOW AND WHEN TO APPLY:

Apply when broadleaf weeds are actively growing in spring or early fall.

WHAT TO EXPECT:

Weed stems and leaves will begin to twist and curl in a few days, but a month or more may be required for complete control. This product will not prevent weed seeds from producing new plants.

- Mow lawn to normal height 1 to 2 days before application.
- Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see 4th bullet).
- Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants.

- Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.
- Avoid applying when temperatures are consistently above 90°F.

SPREADING TIPS:

To provide proper distribution, calibrate spreader before application.
For best results, use a Scotts® Spreader.

Spreader Settings		DIRECTIONS
Scotts AccuGreen® Drop Spreaders	X	Mow lawn first; apply 2 header strips; apply longest direction of lawn, overlapping wheel marks.
Scotts PrecisionGreen® Drop Spreader	X x/x	Mow lawn first; apply 2 header strips; apply longest direction of lawn, overlapping wheel marks.
Scotts SpeedyGreen® & EdgeGuard® Rotary Spreaders	X	Make each pass about 5 feet from previous pass to overlap spread width.
Scotts EasyGreen® Rotary	X	Make each pass about 30 inches from previous pass to overlap spread width.

[Optional Direction Illustrations May Be Presented]

Additional Spreader Settings:

- EZ Drop, Republic (# 530)..... X
- EZ Broadcast, Republic (# 540)..... X x/x
- Scotts Broadcast, Tractor Spreader..... X

(Non-Scotts spreader settings are approximate. You may need to adjust for proper coverage. Check your spreader on xxx sq.ft. and adjust to apply at the rate of 1 pound of product per xxx sq.ft.)

[Note to Supplemental Registrants: Refer to table for appropriate bag sizes.]

- For 2.4 lbs. of product per 1,000 sq.ft. of turfgrass: Check your spreader on 420 sq.ft. and adjust to apply at the rate of 1 lb. of product per 420 sq.ft.
- For 2.6 lbs. of product per 1,000 sq.ft. of turfgrass: Check your spreader on 385 sq.ft. and adjust to apply at the rate of 1 lb. of product per 385 sq.ft.
- For 2.8 lbs. of product per 1,000 sq.ft. of turfgrass: Check your spreader on 360 sq.ft. and adjust to apply at the rate of 1 lb. of product per 360 sq.ft.
- For 2.8 lbs. of product per 1,000 sq.ft. of turfgrass: Check your spreader on 360 sq.ft. and adjust to apply at the rate of 1 lb. of product per 360 sq.ft.
- For 3.0 lbs. of product per 1,000 sq.ft. of turfgrass: Check your spreader on 333 sq.ft. and adjust to apply at the rate of 1 lb. of product per 333 sq.ft.
- For 3.2 lbs. of product per 1,000 sq.ft. of turfgrass: Check your spreader on 310 sq.ft. and adjust to apply at the rate of 1 lb. of product per 310 sq.ft.
- For 3.4 lbs. of product per 1,000 sq.ft. of turfgrass: Check your spreader on 300 sq.ft. and adjust to apply at the rate of 1 lb. of product per 300 sq.ft.
- For 3.6 lbs. of product per 1,000 sq.ft. of turfgrass: Check your spreader on 275 sq.ft. and adjust to apply at the rate of 1 lb. of product per 275 sq.ft.

GUARANTEED ANALYSIS XX-X-X	
Total Nitrogen (N)	XX%
x.x% ammoniacal nitrogen	
x.x% urea nitrogen*	
Available Phosphate (P2O5)	X%
Soluble Potash (K2O)	X%
Sulfur (S)	X%
x.x% free sulfur (S)	
Derived from: polymer-coated sulfur-coated urea, urea, ammonium phosphate, potassium chloride.	
*A portion of the urea has been coated to provide x.x% coated slow-release nitrogen.	
This product and/or its manufacture is covered by one or more of the following U.S. patents and international counterparts thereof: US Patent Nos: 5,219,465; 5,300,135; 5,405,426; 5,466,274.	

APPENDIX

1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Weight: _____
- EPA Est. No. _____

21. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

- Container Packaging and Size: Approved retail container package sizes include: 12, 13, 14, 14.29, 15, 16, 17, 18, 24, 25, 26, 28, 34, 36, 37, 39, 42, 43, 48, and 49 pounds.
- Previously Approved Fertilizer Grades: Fertilizer grades with nutrient contents expressed as the percentages of N, P2O5, and K2O include the following:

19-6-8	23-3-7	26-3-6	29-3-4
20-0-4	24-0-4	26-4-4	30-3-3
22-3-3	24-4-8	26-4-6	30-4-4
22-3-6	25-3-3	28-0-3	32-3-3
22-3-17	25-4-8	29-0-4	32-4-8
22-4-11			

The fertilizer grade guaranteed analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.

23-3-7 GUARANTEED ANALYSIS	
Total Nitrogen (N)	23%
Water Insoluble Nitrogen	5%
Available Phosphate (P2O5)	3%
Soluble Potash (K2O)	7%

28-0-3 GUARANTEED ANALYSIS	
Total Nitrogen (N)	28%
— 28% urea nitrogen*	
Available Phosphate (P ₂ O ₅)	0%
Soluble Potash (K ₂ O)	3%
Sulfur (S) (Free)	4%
Derived from: polymer-coated sulfur-coated urea, urea and potassium chloride.	
*A portion of the urea has been coated to provide 9.6% coated slow-release nitrogen.	

22-4-11 GUARANTEED ANALYSIS	
Total Nitrogen (N)	22%
— 1.6% ammoniacal nitrogen	
— 20.4% urea nitrogen*	
Available Phosphate (P ₂ O ₅)	4%
Soluble Potash (K ₂ O)	11%
Sulfur (S) (Free)	2.2%
Derived from: sulfur-coated polymer-coated urea, urea, ammonium phosphate, potassium chloride.	
*A portion of the urea has been coated to provide 5.5% coated slow-release nitrogen.	

Herbicide Benefit Claims:

- Controls [Kills] [desired weed species from attached listing may be inserted here e.g. dandelion, chickweed, knotweed, plantain, henbit, spurge] and many other broadleaf weeds [as listed] [including those listed on the back panel]
- Small particle size for uniform coverage to control a wide range of weeds.
- Controls [Kills] broadleaf lawn weeds.
- Controls [Kills] [Number] common lawn weeds including dandelion and clover.
- An easy guide to timing your application is to look for dandelions on your lawn.
- This product can be applied in the fall when weeds are actively growing. Because plants are taking nutrients into their roots at this time of year, it's easier to control some weeds in the fall than in the spring.
- Knocks out weeds
- [Kills] [Controls] [Clears out] (desired weed species from attached list may be inserted here) and over (Number) other broadleaf weeds [after they spout and helps make your lawn its thickest, greenest ever].
- [Kills] [Controls] [Clears out] (desired weed species from attached list may be inserted here) and other major lawn weeds.
- All you need for a thicker, greener lawn without (desired weed species from attached list may be inserted here).
- [Contains] [With] [2,4-D] [For Difficult Weeds] [For Strong Weed Control]
- Weed stems and leaves will begin to twist and curl in a few days, but a month or more may be required for complete control.
- This product will not prevent weed seeds from producing new plants.
- Apply anytime [when broadleaf] weeds are actively growing.
- Controls fall lawn weeds and prepares your lawn from the stress of winter.
- Herbicide eliminates weeds, while fertilizer [plus natural microorganisms] give you a [thicker,] [greener] lawn.
- [Controls] [Kills] (desired weed species from attached list may be inserted here) [and [most] other [broadleaf] weeds.] [and many more.] [while greening your lawn.]
- One product that does it all, [kills] [controls] weeds and gives you a [healthier,] [greener] lawn
- Feeds, greens and thickens lawns while controlling weeds.
- Weed control penetrates down to the root to kill dandelion, clover and other broadleaf weeds.
- Clears out dandelions and other broadleaf weeds.

- Knocks out weeds.
- Kills Dandelion, Chickweed, Clover (Over 200 Weeds).
- Apply in Fall to Reduce Spring Weeds.
- A combination of 3 proven weed killers controls a wide range of [listed] lawn weeds

Fertilizer Benefit Claims:

- Long lasting, lush green lawn.
- Stimulates deep greening.
- Stimulates quick, deep greening.
- Stimulates early, vigorous root growth.
- Stimulates thick, lush grass blades.
- Thickens and greens lawns.
- One-step application for a lush green lawn.
- Fast deep greening balanced with long-term feeding by patented Polyon® coated fertilizer technology.
- Patented Polyon® coated fertilizer technology for timed release of fertilizer.
- Polyon® coating prevents burning.
- Scotts® slow-release polymer-encapsulated nitrogen fertilizer for uniform growth with extended feeding.
- Provides a regular turf feeding.
- Feeds up to 8 weeks.
- The fertilizer combination of nitrogen, phosphate, and potash gives your lawn strength above and below the surface. This product promotes root and rhizome development, helping the grass to produce more blades. The result is a thicker, greener lawn.
- Phosphorous-Free Formulation
- Scotts® Phosphorous-Free fertilizer provides the best formula for soils already containing a high natural presence of available phosphorous. A soil test can help determine if your soil will support normal turf growth without a phosphorous supplement.
 - Nitrogen for Color — Contributes to thicker, greener growth.
 - Phosphorous for strength — Stimulates root growth. Test your soil regularly to see if phosphorous should be applied.
 - Potassium for Protection — Protects turf against disease and drought damage.
- Scotts Phosphorous-Free products are specifically formulated with Scotts Poly-S® timed-release fertilizer. The unique Poly-S® coating provides your lawn with uniform growth and an extended feeding for up to two months.
- With the right weather conditions and correct application, your lawn will be greener and thin areas of turf will fill in, making the density of your lawn more uniform.
- We guarantee a thick, green lawn without the risk of surge growth
- A thick, green lawn
- [We guarantee a] thick(er), green(er) lawn.
- Poly-S® Technology.
- Specially formulated with Scotts Poly S® timed-release fertilizer [as used by landscape professionals.]
- Specially formulated with Scotts® Poly-S® timed-release fertilizer to provide uniform growth and extended feeding.
- Contains 40% of total guarantee of nitrogen as controlled release nitrogen for uniform growth and extended greening.
- Feeds up to 2 months.
- Contains Nutralene® Controlled Release Nitrogen.
- Nutralene is a controlled release homogenous proven technology. It releases nitrogen gradually through hydrolysis and microbial action in the soil for a sustained feeding. Nutralene is unlike other nitrogen sources that depend on coatings, particle size, and temperature for their nitrogen release.
- The ultimate timed-released fertilizer as used by professional golf courses.

- Contains both readily available nitrogen for quick green up and controlled release nitrogen for uniform growth and extended feeding.
- Feeds, greens and thickens lawns.
- Strengthens Roots for Winter Hardiness.
- [For Clorox/Naturize] Fertilize with (This Product -or- Registrant Name) [lawn products] for thick, green lawns
- [For Clorox/Naturize] Promotes a deep, green lawn [with stronger roots]
- [For Clorox/Naturize] Promotes a strong, deep root system in the fall for [a] quicker [green-up] [greening] in the spring.
- [For Clorox/Naturize] [Feeds your lawn] with time-released nitrogen for extended greening.
- [For Clorox/Naturize] Builds a healthy, green, and thick lawn.
- [For Clorox/Naturize] Apply anytime for a [greener,] [thicker] lawn.
- [For Clorox/Naturize] Makes your lawn healthier.
- [For Clorox/Naturize] (This Product -or- Registrant Name) [Lawn Food] [lawn products] add(s) [natural] [beneficial] [soil] microorganisms.
- [For Clorox/Naturize] The Result: [thicker] [,][heartier][,] [greener] grass than ordinary lawn fertilizers....Guaranteed!
- [For Clorox/Naturize] [(This Product -or- Registrant Name)] [lawn products] Add(s) [Natural] [Beneficial] [Soil] Microorganisms to Feed the Soil as Well as Your Grass.
- [For Clorox/Naturize] Better Results than Ordinary Fertilizer Guaranteed!
- [For Clorox/Naturize] Ultra Greening Power!
- [For Clorox/Naturize] The patented formula in (This Product -or- Registrant Name) [Lawn Food] [Lawn Food with Weed Control] [lawn products] [actually] blends [natural] [,] [beneficial] microorganisms – the same found in [fertile soil] [compost] – with [traditional] lawn food nutrients.
- [For Clorox/Naturize] These microorganisms allow your lawn to absorb more nutrients.
- [For Clorox/Naturize] The result is a [thicker] [,] [healthier] [,] [more robust] lawn.
- [For Clorox/Naturize] (This Product -or- Registrant Name) [lawn products] provide(s) both the nutrients and the microbials that a [healthy] lawn needs.
- [For Clorox/Naturize] Builds [stronger][,] [deeper] [,][bigger] roots
- [For Clorox/Naturize] [Feeds] [Restores] the soil [as well as the plant].
- [For Clorox/Naturize] Made in the U.S.A.
- [For Clorox/Naturize] Image Statements (possible text statements beneath any approved image)
 - Fed [lawn] [grass]
 - Unfed [lawn] [grass]
 - [Traditional] [ordinary] fertilizer
 - (This Product -or- Registrant Name) [lawn]
 - [With] [Contains][Without] [No][Beneficial] [Natural] [Soil] Microorganisms

General Marketing Claims:

- XX pounds covers up to XX square feet.
- Ortho product quality you've come to trust.
- Scientific blend of small particles promotes fast deep greening and balanced long term feeding.
- Outdoor use only
- Satisfaction guaranteed* or your purchase price refunded with proof-of-purchase and original cash register receipt.
- This product is designed to be used in the spring, when weeds are actively growing.
- Apply anytime weeds are actively growing.
- (Number) pounds treats and feeds (Number) square feet of turfgrass.
- For best results, [always] use a Scotts® spreader.
- Visit our website www.xxxxx.com.
- Scotts No Quibble Guarantee(TM). If for any reason you are not satisfied with results after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.
- Answers to FAQ's on the internet at www.xxxxx.com

- Produce superior results . . . every time.
- Treats and feeds _____ sq.ft.
- Proudly produced in the U.S.A.
- Can we help you? Call toll-free: 1-XXX-XXX-XXXX or visit our website at www.XXXXXXX.com.
- Apply in late fall.
- Feeds & treats 5,000 sq.ft.
- Apply once in Spring when weeds are growing
- To spot treat individual weeds, use Ortho Weed-B-Gon Weed Killer®.
- Guarantees offered by supplemental registrants may include the following:
 - CAN'T FAIL GUARANTEE. If you are not completely satisfied with the results after using this product, (Company Name) will refund your purchase price in full. Simply send proof of purchase and register receipt to (Company Name and Address).

Referrals to other products used sequentially:

- These products are the quality choice in complete lawn care. You determine what your lawn needs, and you'll find a product to help you. Fertilizers, lawn starters, crabgrass prevention, weed and insect control - everything for a lush green lawn and year-round protection. Each one is easy to apply and guaranteed to produce quick, long-lasting results.
- This product makes it easier to have a lush, green lawn. Our four (4) simple applications eliminate the guesswork and give you professional lawn care results. Each stage combines just the right amount of fertilizer with the right amount of control and prevention. Just match each stage to the correct season and you'll prevent crabgrass, control weeds and insects, and prepare your lawn for the winter. It's what your lawn needs for strong growth.
 - Stage One (February to April) is a pre-emergent -- Scotts Phosphorous-Free fertilizer with Crabgrass Preventer
 - Stage Two is a broadleaf postemergent – Scotts Phosphorous-Free fertilizer with Weed Control
 - Stage Three is an insecticide.
 - Stage Four is a fall broadleaf postemergent.
- For best results, apply (brand name) products.
- Apply using the 4-Step Program
- **4-Step Program:**
 - STEP 1:** Menards Premium Crabgrass Preventer - Apply March - early May - Prevents crabgrass, foxtail, goosegrass, poa annua and other annual weed grasses. - Nutralene feeds your turfgrass for up to 2 months with long lasting deep greening.
 - STEP 2:** Menards Premium Weed & Feed - Apply May - June - Kills dandelion, clover, plantain and over XXX other broadleaf weeds. - Nutralene feeds your turfgrass for up to 2 months with long lasting deep greening.
 - STEP 3:** Menards Premium Lawn Food - Apply June - August - Nourishes your lawn throughout the summer months with extended feeding of Nitrogen. - Nutralene feeds your turfgrass for up to 2 months with long lasting deep greening.
 - STEP 4:** Menards Premium Winterizer & Weed Control - Apply late August - mid October - Kills dandelion, clover, plantain and over XXX other broadleaf weeds. - Maximizes root building. - Nutralene feeds your turfgrass for up to 2 months with long lasting deep greening.
- OR**
- STEP 4:** Menards Premium Starter & Winterizer - Apply late August - November - Maximizes root building. - Nutralene feeds your turfgrass for up to 2 months with long lasting deep greening. - Use with new seeding and sodding of turfgrass.

2. State mandated fertilizer labeling requirements:

[Note to Supplemental Registrants: Please select the statements relevant to the areas of the country in which your specific product is registered and sold; if any of these statements are relevant to your labeling, please place in the Directions for Use under the heading "3.1 State Restrictions"]

GUARANTEED ANALYSIS XX-X-X	
Total Nitrogen (N)	XX%
x.x% ammoniacal nitrogen	
x.x% urea nitrogen*	
Available Phosphate (P ₂ O ₅)	X%
Soluble Potash (K ₂ O)	X%
Sulfur (S)	X%
x.x% free sulfur (S)	
Derived from: polymer-coated sulfur-coated urea, urea, ammonium phosphate, potassium chloride.	
*A portion of the urea has been coated to provide x.x% coated slow-release nitrogen.	
This product and/or its manufacture is covered by one or more of the following U.S. patents and international counterparts thereof: US Patent Numbers: xxxxxx	

[Note: Fertilizer grades with nutrient contents expressed as the percentages of N, P₂O₅ and K₂O may include the following: 23-3-7 or x-x-x. The fertilizer grade analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.]

- California, Oregon, Washington: “[When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials.]Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>”
- Florida: [Note to reviewer: maximum number of applications in this section is dually based upon maximum ai load as well as fertilizer analysis effect on maximum nitrogen rates in different regions in Florida]:
 - In North Florida: Do not exceed X applications per year on bermudagrass. Do not exceed X applications per year on centipedegrass. Do not exceed X applications per year on St. Augustine grass. Do not exceed X applications per year on zoysiagrass.
 - In Central Florida: Do not exceed X applications per year on bermudagrass. Do not exceed X applications per year on centipedegrass. Do not exceed X applications per year on St. Augustine grass. Do not exceed X applications per year on zoysiagrass.
 - In South Florida: Do not exceed X applications per year on bermudagrass. Do not exceed X applications per year on centipedegrass. Do not exceed X applications per year on St. Augustine grass. Do not exceed X applications per year on zoysiagrass.
- Florida: “Apply only to actively growing turf.”
- Florida: “Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices. Check with your county or city government to determine if there are local regulations for fertilizer use.
- Maryland: “This fertilizer contains phosphorus and may not be used on turf in the state of Maryland except when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory identified under § 8-803.7 of the Agriculture Article, Annotated Code of Maryland, performed no more than 3 years before the application; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State; or 3) Reestablishing or repairing a turf area.”
- New Jersey: “In the state of New Jersey, this product may be used only when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary pursuant to a soil test conducted by a laboratory identified pursuant to subsection b. of section 6 of P.L.2010, c.112 (C.58:10A-66) and performed no more than three years before the application, and pursuant to the associated annual fertilizer recommendation issued by the New Jersey Agricultural Experiment Station at Rutgers, the State University; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is in accordance with the standards and requirements established under the “Soil Erosion and Sediment Control Act,” P.L.1975, c.251 (C.4:24-39 et seq.)

and the rules and regulations adopted pursuant thereto; 3) Reestablishing or repairing a turf area; or 4) Delivering liquid or granular fertilizer under the soil surface, directly to the feeder roots.”

- New Hampshire: In the state of New Hampshire, this product may only be used on 1) Establishing new lawns, for repairing a lawn, for seeding, or for use when a soil test indicates a phosphorus deficiency.
- Virginia: “In the state of Virginia, this product may be used only on 1) Newly established turf areas from sod or seed during the first growing season; 2) Turf areas being repaired or renovated, and; 3) Turf areas where soil tests performed within the past three years indicate nutrient deficiency.”
- [Virginia, Florida, Maryland, others] DO NOT APPLY NEAR WATER, STORM DRAINS, OR DRAINAGE DITCHES. DO NOT APPLY IF HEAVY RAIN IS EXPECTED. APPLY THIS PRODUCT ONLY TO YOUR LAWN/GARDEN. [AND SWEEP ANY PRODUCT THAT LANDS ON THE DRIVEWAY, SIDEWALK, OR STREET, BACK INTO YOUR LAWN/GARDEN.] [Note to Supplemental Registrant: The bracketed part of this statement is not required for liquid products.]
- [For products containing Iron:] Keep this product off sidewalks, patios, driveways, etc. This fertilizer contains iron which will stain concrete and painted surfaces. If it comes in contact with concrete surface, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.
- [Note to Supplemental Distributor: In Kentucky if the product supplies less than 1lb of Nitrogen per application, include the following:] A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing XX pounds of nitrogen per 1,000 square feet and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.”

[Note to Supplemental Registrants: Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, equipment settings of drop spreaders, rotary spreaders, and other granular applicators may be presented on the container labels.]

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.	Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.
12	2.4	5,000	26	2.4	10,800
	2.6	4,600		2.6	10,000
	2.8	4,300		2.8	9,300
	3.0	4,000		3.0	8,700
	3.2	3,800		3.2	8,100
	3.4	-----		3.4	7,600
	3.6	-----		3.6	7,200
13	2.4	5,400	28	2.4	11,700
	2.6	5,000		2.6	10,800
	2.8	4,600		2.8	10,000
	3.0	4,300		3.0	9,300
	3.2	4,100		3.2	8,800
	3.4	3,800		3.4	8,200
	3.6	3,600		3.6	7,800
14	2.4	5,800	36	2.4	15,000
	2.6	5,400		2.6	13,800
	2.8	5,000		2.8	12,900
	3.0	4,700		3.0	12,000
	3.2	4,400		3.2	11,300
	3.4	4,100		3.4	10,500
	3.6	3,900		3.6	10,000

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.	Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.
15	2.4	6,300	37	2.4	15,400
	2.6	5,800		2.6	14,200
	2.8	5,400		2.8	13,200
	3.0	5,000		3.0	12,300
	3.2	4,700		3.2	11,600
	3.4	4,400		3.4	10,900
	3.6	4,200		3.6	10,300
16	2.4	6,700	39	2.4	16,300
	2.6	6,200		2.6	15,000
	2.8	5,700		2.8	13,900
	3.0	5,300		3.0	13,000
	3.2	5,000		3.2	12,200
	3.4	4,700		3.4	11,500
	3.6	4,400		3.6	10,800
17	2.4	7,100	42	2.4	17,500
	2.6	6,500		2.6	16,200
	2.8	6,100		2.8	15,000
	3.0	5,700		3.0	14,000
	3.2	5,300		3.2	13,100
	3.4	5,000		3.4	12,400
	3.6	4,700		3.6	11,700
18	2.4	7,500	43	2.4	17,900
	2.6	6,900		2.6	16,500
	2.8	6,400		2.8	15,400
	3.0	6,000		3.0	14,300
	3.2	5,600		3.2	13,400
	3.4	5,300		3.4	12,600
	3.6	5,000		3.6	11,900
24	2.4	10,000	48	2.4	20,000
	2.6	9,200		2.6	18,500
	2.8	8,600		2.8	17,100
	3.0	8,000		3.0	16,000
	3.2	7,500		3.2	15,000
	3.4	7,100		3.4	14,100
	3.6	6,700		3.6	13,300
25	2.4	10,400	49	2.4	20,400
	2.6	9,600		2.6	18,800
	2.8	8,900		2.8	17,500
	3.0	8,300		3.0	16,300
	3.2	7,800		3.2	15,300
	3.4	7,400		3.4	14,400
	3.6	6,900		3.6	13,600

DOCUMENT CONTROL INFORMATION

1. Unique Label Identifier: 002217-00660.20170113.notif-proposed-highlighted

2. Reason for Issue: ad claims, state mandated fertilizer language, formatting, minor corrections, delete obsolete sublabels

Internal Usage Notes
Associated Container Label: Supplemental Distributors