

# TRIMEC®

## WEED & FEED

### FOR SOUTHERN LAWNS

ACCEPTED  
20-10-5  
EPA  
88-17-661A



**Eighteen Pounds Covers up to 5,000 Square Feet**

**KILLS:** Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and many other broadleaf weeds in St. Augustine, Centipede, Bahia, Bermuda, Zoysia, Bluegrass, Rye, Fescue, and Bentgrass lawns; some of which are listed on the back panel.

**ACTIVE INGREDIENTS:**

- \*Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy)propionic acid 0.641%
  - \*\*Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid 0.144%
  - \*\*\*Dimethylamine Salt of Dicamba 3,6-dichloro-o-anisic acid 0.124%
- INERT INGREDIENTS** 99.093%
- TOTAL** 100.000%

**THIS PRODUCT CONTAINS:**

- 1) 609 lbs. or 0.530% 2-(2-methyl-4-chlorophenoxy)propionic acid per ton or 0.832 lb. per acre
  - 2) 3062 lbs. or 0.153% 2,4-Dichlorophenoxyacetic acid per ton or 0.240 lb. per acre
  - 3) 304 lbs. or 0.065% 3,6-dichloro-o-anisic acid per ton or 0.102 lb. per acre
- \*Isomer Specific by AOAC Method 6 D01-5

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**STATEMENT OF PRACTICAL TREATMENT**

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention.

See back panel for additional precautionary statements.

TRIMEC is a registered trademark of PBI-GORDON CORPORATION  
U.S. Patent No. 3,341,186

**NET WEIGHT 18 POUNDS**



KANSAS CITY, KANSAS 66118



**READ THE ENTIRE LABEL FIRST**  
**OBSERVE ALL CAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans & Domestic Animals**  
CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. Wash thoroughly after handling.

**Environmental Hazards**  
Do not apply directly to water. Do not apply when weather conditions favor drift from target area.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs. To prevent cross contamination, do not store near other fertilizers or near herbicides, insecticides, fungicides or seeds. **PRODUCT DISPOSAL:** To dispose of bags which contain unused product, seal, wrap bag in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse empty bag. Fold empty bag and discard in trash.

**USE PRECAUTIONS**

Do not use this product where desirable plants are present or where the material may drift onto lawns, vegetable gardens, ornamentals, shrubbery. Do not apply in areas of high traffic. Do not use on lawns. Apply when the air is calm. Do not apply to newly seeded lawns. During application, take care to avoid contact with children. Do not contaminate food or feedstuffs.

**INSTRUCTIONS**

1. This Weed & Feed for Southern Lawns is a TRIMEC product and contains active ingredients: 2-(2-methyl-4-chlorophenoxy)propionic acid, 2,4-Dichlorophenoxyacetic acid, and Dicamba. It is a weed and feed product. It is designed to kill weeds and feed your lawn. It is designed to kill weeds in St. Augustine, Centipede, Bahia, Bermuda, Zoysia, Bluegrass, Rye, Fescue, and Bentgrass lawns; some of which are listed on the back panel. It is designed to kill weeds in St. Augustine, Centipede, Bahia, Bermuda, Zoysia, Bluegrass, Rye, Fescue, and Bentgrass lawns; some of which are listed on the back panel. It is designed to kill weeds in St. Augustine, Centipede, Bahia, Bermuda, Zoysia, Bluegrass, Rye, Fescue, and Bentgrass lawns; some of which are listed on the back panel.
2. Water lawns thoroughly at least 1 day before application. Do not water lawns for 2 days after application.
3. Apply when appropriate wind direction will prevent drift onto desirable plants. Do not overapply. Do not apply to newly seeded lawns or to lawns that have been recently fertilized. Do not apply to lawns that have been recently mowed. Do not apply to lawns that have been recently watered. Do not apply to lawns that have been recently treated with other lawn care products.
4. Do not walk on lawns for 2 days after application. Avoid contact with lawns for 2 days after application.

**Be patient: TRIMEC works slowly and gently and is friendly to your lawn.**

**SPREADER SETTINGS**

The appropriate spreader settings are approximate. They vary with the type of spreader used and the type of lawn. For more information, see the back panel of this product.

**FERTILIZER GUARANTEE**

This fertilizer contains 18% Nitrogen and 18% Phosphorus. It is a complete fertilizer. It is designed to feed your lawn. It is designed to feed your lawn. It is designed to feed your lawn.

WEEDS			
BEDSTRAW	DANDELION	LESPEDEZA	POISON IVY
BLACK MEDIC	DOCK	MALLOW	POISON OAK
BUCKHORN	GROUND IVY	MORNING GLORY	PURSLANE
BURDOCK	HEAL ALL	OTALIS	RAGWEED
CHICORY	HENBIT	PEPPERGRASS	SHEEP SORREL
CHICKWEED	KNOTWEED	PIGWEEED	SHEPHERD'S PURSE
CLOVER	LAMBSQUARTER	PLANTAINS	SPEEDWELL
			SPURGE
			THISTLE
			WILD CARROT
			WILD GARLIC
			WILD LETTUCE
			WILD ONION
			YARROW

**NOTICE:** Do not walk on lawns for 2 days after application. Avoid contact with lawns for 2 days after application. Do not walk on lawns for 2 days after application. Avoid contact with lawns for 2 days after application.