

TRIMEC®

WEED & FEED

FOR SOUTHERN LAWNS

20-10-5



16 Pounds Covers up to 5,000 Square Feet

KILLS: Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and many other broadleaf weeds in St. Augustine, Centipede, Bahia, Bermuda, Zoysia and Bentgrass lawns; some of which are listed on the back panel.

ACTIVE INGREDIENTS:

- 1. Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid 0.641%
- 2. Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid 0.184%
- 3. Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid) 0.078%

INERT INGREDIENTS

99.097%

TOTAL 100.000%

THIS PRODUCT CONTAINS

- 10.609 lbs. or 0.530% 2-(2-methyl-4-chlorophenoxy) propionic acid per ton
- 3.062 lbs. or 0.153% 2,4-Dichlorophenoxyacetic acid per ton
- 1.314 lbs. or 0.065% 3,6-dichloro-o-anisic acid per ton

Isomer Specific by AOAC Method

JUL 22 1961

KEEP OUT OF REACH OF CHILDREN
CAUTION

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water. For eyes, flush with water for 15 minutes and get medical attention. See back panel for additional precautionary statements.

TRIMEC is a registered trademark of PBI-GORDON CORPORATION

U.S. Patent No. 3,284,188

NET WEIGHT 16 POUNDS



KANSAS CITY, KANSAS 66118



READ THE ENTIRE LABEL FIRST
OBSERVE ALL CAUTIONS AND FOLLOW DIRECTIONS CAREFULLY

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Do not breathe dust. Do not get in contact with skin, eyes or clothing. Wash thoroughly after handling.

Environmental Hazards

Do not apply directly to water. Do not apply when weather conditions favor drift to target area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE

Store in original container and keep away from children and pets. Do not use in a storage area inaccessible to children and domestic animals. Do not store in a room with food, feed, or bedding. To prevent cross contamination, do not store near other fertilizers or herbicides. Do not store in a room with fungicides or seeds. **PRODUCT DISPOSAL:** To dispose of this material, seal the unused product securely in a bag in several layers of newspaper and place in trash. **CONTAINER DISPOSAL:** Do not reuse empty bag. Fold empty bag and discard in trash.

USE PRECAUTIONS

Do not use this product where desirable plants are present. Do not use on lawns with flowers, fruiting flowers, vegetables, ornamental shrubs, etc. Do not use on trees, shrubs, vines, etc. Do not use on lawns or shrubs. Apply when the lawn is calm. Do not apply to lawns or shrubs during a rain or heavy dew. Do not use spreader openings when turning. Do not use on roads, driveways, sidewalks, etc.

INSTRUCTIONS

Gordon's Weed & Feed For Southern Lawns is a TRIMEC complex, a combination of three proven weed killers - 2,4-D, Mecoprop and Dicamba. Together they control a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge, etc. This complex is found in home lawns. In addition, a 20-10-5 fertilizer is also present. A supply of plant food is provided for a normal season's growth. Post-treatment may be necessary if a lawn is heavily overgrown by weeds. In case of adverse lawn moisture conditions prevail and weeds are in a state of poor growth. A second application may be made approximately 30 days later. Do not use in drought. Do not overapply. In the conditions of adequate moisture, the normal rate of 20-10-5 is sufficient. Do not use in drought conditions. Do not use on St. Augustine, Centipede, Bermuda, Zoysia, or other warm season grasses. Do not use on St. Augustine, Centipede, Bermuda, Zoysia, or other warm season grasses. Do not use on St. Augustine, Centipede, Bermuda, Zoysia, or other warm season grasses.

1. Mow lawn to normal height (1 1/2 to 2 inches) before application.
2. Water lawn thoroughly at least 3 days before application. Water lawn thoroughly after treatment. Use step 4.
3. Apply when weeds are young and active. Apply preferentially to the young weeds. Do not apply to grass if grass is in a state of poor growth. Do not apply to lawns with flowers, fruiting flowers, or vegetables. Do not apply to lawns or shrubs during a rain or heavy dew.
4. Do not water lawn until weeds are dead. Do not apply to lawns with flowers, fruiting flowers, or vegetables.

Be patient. TRIMEC works slowly and gently and is friendly to your lawn.

SPREADER SETTINGS

The application settings are given in terms of pounds of product per 1,000 square feet. For broadcast application, the rate is 1.314 lbs. per 1,000 sq. ft.

FERTILIZER GUARANTEE

Total Nitrogen... Water Soluble Nitrogen... Available Phosphorus... Soluble Phosphate...

WEEDS			
BEDSTRAW	DANDELION	LESPEDEZA	POISON IVY
BLACK MEDIC	DOCK	MALLOW	POISON OAK
BUCKHORN	GROUND IVY	MORNING GLORY	PURSLANE
BURDOCK	HEAL ALL	OXALIS	RAGWEED
CHICORY	HENBIT	PEPPERGRASS	SHEEP SORREL
CHICKWEED	KNOTWEED	PIGWEED	SHEPHERD'S PURSE
CLOVER	LAMBSQUARTER	PLANTAINS	SPEEDWELL
			SPURGE
			THISTLE
			WILD CARROT
			WILD GARLIC
			WILD LETTUCE
			WILD ONION
			YARROW

NOTICE: For best results, this product should be applied to lawns during the growing season. Do not apply to lawns with flowers, fruiting flowers, or vegetables. Do not apply to lawns or shrubs during a rain or heavy dew. Do not use on St. Augustine, Centipede, Bermuda, Zoysia, or other warm season grasses. Do not use on St. Augustine, Centipede, Bermuda, Zoysia, or other warm season grasses.

ESTABLISHED 1951

1982, PBI-GORDON CORP.

†TRIMEC®

WEED & FEED

FOR SOUTHERN LAWNS

20-10-5



Eighteen Pounds Covers up to 5,000 Square Feet

KILLS: Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and many other broadleaf weeds in St. Augustine, Centipede, Bahia, Bermuda, Zoysia and Bentgrass lawns; some of which are listed on the back panel

ACTIVE INGREDIENTS

- ***Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid 0.641%
- **Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid 0.184%
- ***Dimethylamine Salt of Dicamba(3,6-dichloro-o-anisic acid) 0.078%

INERT INGREDIENTS

TOTAL 100.000%

THIS PRODUCT CONTAINS:

- 10.609 lbs. or 0.530% 2-(2-methyl-4-chlorophenoxy) propionic acid per ton
- 3.062 lbs. or 0.153% 2,4-Dichlorophenoxyacetic acid per ton
- 1.104 lbs. or 0.065% 3,6-dichloro-o-anisic acid per ton
- †††Isomer Specific by AOAC Method

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water. For eyes, flush with water for 15 minutes and get medical attention.

See back panel for additional precautionary statements

TRIMEC® is a registered trademark of PBI GORDON CORPORATION

U.S. Patent No. 3,284,186

NET WEIGHT 18 POUNDS



READ THE ENTIRE LABEL FIRST
OBSERVE ALL CAUTIONS AND FOLLOW DIRECTIONS CAREFULLY

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Do not inhale dust. Eye irritant. Wear contact with skin. Avoid getting on clothing. Wash thoroughly after handling.

Environmental Hazards

Do not apply directly to water. Do not apply when weather conditions favor drift to non-target area.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or livestock. To prevent cross contamination, do not store near other fertilizers, herbicides, insecticides, fungicides, or seeds. **PRODUCT DISPOSAL:** This use of bags which contain unused product. Securely wrap bag in several layers of newspaper, place in trash. **CONTAINER DISPOSAL:** Do not reuse empty bag. Fold empty bag and discard in trash.

USE PRECAUTIONS

Do not use this product where desirable covers are present. Do not use on lawns with annual weeds, flowers, vegetables or ornamental shrubs. Do not use on lawns of Bentgrass, Zoysia, St. Augustine, Centipede, or shrubs. Apply when the air is calm to avoid blowing dust. Do not use on lawns of Bentgrass, Zoysia, St. Augustine, Centipede, or shrubs when the air is calm to avoid blowing dust. Do not use on lawns of Bentgrass, Zoysia, St. Augustine, Centipede, or shrubs when the air is calm to avoid blowing dust.

INSTRUCTIONS

Gordon's Weed & Feed For Southern Lawns is a TRIMEC complex combination of three proven weed killers - 2,4-D, MCP, and a 3,6-Dichloro-o-anisic acid. It kills actively growing broadleaf weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge, and many other broadleaf weeds in addition to a 20-10-5 fertilizer. It is a pre-emergent herbicide for lawns. Apply to lawns in a normal season (e.g., with the retreatment) may be necessary if a lawn is heavily infested by weeds (0.1 lb) if adverse low moisture conditions prevail and weeds are in a state of rapid growth. A second application may be made in approximately 30 days. Do not use under the following conditions: (1) inadequate moisture; (2) high temperature and high humidity; (3) drought; (4) major or temporary discoloration of St. Augustine under these conditions. Where the application of St. Augustine will choke out most weeds and a first-rate treatment would not be necessary.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not in a state of application, sprinkle gently with water. Do not water immediately to prevent dust drift to non-target plants.
4. Do not wash from lawns for 1 to 2 days after application. Allow some additional watering if needed to be made.

Be patient. TRIMEC works slowly and gently and is friendly to your lawn.

SPREADER SETTINGS

Spreader	1/2
Lawn Fertilizer	1/2
BAKERS	1/2
Spans	1/2

The above spreader settings are approximate. The actual rate of application should be determined by the rate of flow per 275 sq. ft.

FERTILIZER GUARANTEE

Total Nitrogen (not less than)	20%
Water-soluble Nitrogen (not less than)	10%
Available Phosphorus (acid insoluble)	10%
Available Potash (not less than)	5%

WEEDS

BEDSTRAW	DANDELION	LESPEDEZA	POISON IVY	SPURGE
BLACK MEDIC	DOCK	MALLOW	POISON OAK	THISTLE
BUCKHORN	GROUND IVY	MORNING GLORY	PURSLANE	WILD CARROT
BURDOCK	HEAL ALL	OXALIS	RAGWEED	WILD GARLIC
CHICORY	HENBIT	PEPPERGRASS	SHEEP SORREL	WILD LETTUCE
CHICKWEED	KNOTWEED	PIGWEEED	SHEPHERD'S PURSE	WILD ONION
CLOVER	LAMBSQUARTER	PLANTAINS	SPEEDWELL	YARROW

NOTICE: Seller warrants that this product is a true and correct representation of the actual composition of this product as weathered. It is not a guarantee of performance. The actual performance will vary. Seller warrants that this product is a true and correct representation of the actual composition of this product as weathered.

EEA REG. No. 2217-881

© 1982 PBI-GORDON CORP.

1982, PBI-GORDON CORP.