

# TRIMEC®

## WEED & FEED

### FOR SOUTHERN LAWNS

20-10-5



Fifteen Pounds Covers up to 5,000 Square Feet

**KILLS:** Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and many other broadleaf weeds in St. Augustine, Centipede, Bahia, Bermuda, Zoysia and Bentgrass lawns, some of which are listed on the back panel.

**ACTIVE INGREDIENTS**

Methylamine Salt of 2,2-dimethyl-4-chlorophenoxypropionic acid 0.641%  
 Methylamine Salt of 2,4-Dichlorophenoxypropionic acid 0.164%  
 Methylamine Salt of Cloanthra 3,6-dimethylpicolinic acid 0.078%  
 99.997%  
**TOTAL 100.000%**

**INERT INGREDIENTS**

**THIS PRODUCT CONTAINS**

- ✓ 10.641% Methylamine Salt of 2,2-dimethyl-4-chlorophenoxypropionic acid
- ✓ 0.164% Methylamine Salt of 2,4-Dichlorophenoxypropionic acid
- ✓ 0.078% Methylamine Salt of Cloanthra 3,6-dimethylpicolinic acid
- ✓ 99.997% Inert Ingredients

352-101937  
 2217-661

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**STATEMENT OF PRACTICAL TREATMENT**

Apply to the lawn with a broadcast spreader. Follow directions on the back panel. Wash hands thoroughly after use.

U.S. Patent No. 3,284,196

**NET WEIGHT 15 POUNDS**

**Gordon Corporation**  
 KANSAS CITY, KANSAS 66110



**READ THE ENTIRE LABEL FIRST. OBSERVE ALL CAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans & Domestic Animals**  
**CAUTION:** Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. Wash thoroughly after handling.

**Environmental Hazards**  
 Do not apply directly to water. Do not apply when weather conditions favor drift from target area.

**DIRECTIONS FOR USE**  
 It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**STORAGE** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs. To prevent cross contamination, do not store near other fertilizers or near herbicides, insecticides, fungicides, or seeds. **PRODUCT DISPOSAL:** To dispose of bags which contain unused product, securely wrap bag in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse empty bag. Fold empty bag and discard in trash.

**USE PRECAUTIONS**  
 Do not use this product where desirable clovers are present or allow the material to wind drift onto flowers, vegetables or ornamental shrubbery. Do not apply in areas underlaid by roots of desirable trees or shrubs. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning. Do not contaminate domestic or irrigation waters.

**INSTRUCTIONS**  
 Gordon's Weed & Feed For Southern Lawns is a TRIMEC complex, a combination of three proven weed killers - 2,4-D, Mecoprop and Dicamba. Together they control a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a 20-10-5 fertilizer provides a generous supply of plant food needed for a normal season's growth. Possible retreatment may be necessary if a lawn is heavily overrun by weeds. If adverse low moisture conditions prevail and weeds are in a state of poor growth, a second application may be made in approximately 30 days. Do not use on dithionda. **Do not overapply.** Under conditions of adequate moisture, high temperature and high humidity, over application may cause temporary discoloration on St. Augustine. Under these conditions, however, the vigorous growth of St. Augustine will choke out most weeds and herbicide treatment would not be necessary.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent drift from target plants.
4. Do not wash from weed leaves for 1 to 2 days after application. At the same time, thorough watering should be made.

Be patient! TRIMEC works slowly and gently and is friendly to your lawn.

**SPREADER SETTINGS**

Scout	6
Lawn Beauty	5
Excelsior	7
Scots	4

The above spreader settings are approximate. Check your spreader's 330 sq. ft. and adjust to apply at the rate of 1.5 lbs per 330 sq. ft.

**FERTILIZER GUARANTEE**

Total Nitrogen, not less than	20%
Available Phosphate, not less than	5%
Available Potash, not less than	10%
Water Soluble Phosphate, not less than	5%

WEEDS			
BEDSTRAW	DANDELION	LESPEDEZA	POISON IVY
BLACK MEDIC	DOCK	MALLOW	POISON OAK
BUCKHORN	GROUND IVY	MORNING GLORY	PURLANE
BURDOCK	HEAL ALL	OXALIS	RAGWEED
CHICORY	HENBIT	PEPPERGRASS	SHEEP SORREL
CHICKWEED	KNOTWEED	PIGWEED	SHEPHERD'S PURSE
CLOVER	LAMBSQUARTER	PLANTAINS	SPEEDWELL
			SPURGE
			THISTLE
			WILD CARROT
			WILD GARLIC
			WILD LETTUCE
			WILD ONION
			YARROW

**NOTICE:** Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

EPA Reg. No. 2-17-661  
 EPA Est. No. 2217-RS-1  
 © 1982, PBI-GORDON  
 560 1283