TRIMEC[®] WEFD & FEED FOR SOUTHERN LAWNS 20-10-5

PM03 2217-661

1074

(FIFTEEN, SIXTEEN, OR EIGHTEEN) POUNDS COVERS UP TO 5,000 SQUARE FEET

Controls Dandelium, Chickweed, Knotweed, Plantain, Henbit, Spurge and many other broadleaf weeds in St. Augustinegrass, Centipedegrass, Bahiagrass, Bermudagrass, Zoysiagrass, Bluegrass, Ryegrass, Fescue and Bentgrass lawns; some of which are listed on the back panel.


ACTIVE INGREDIENTS: Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy)
propionic acid
propionic acid
acid
<pre>***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic</pre>
acid
INERT INGREDIENTS
TOTAL 100.000%
THIS PRODUCT CONTAINS:
10.609 lbs. or 0.530% 2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton. 3.62 lbs. or 0.153% 2,4-Dichlorophenoxyacetic acid equivalent per ton. 1.304 lbs. or 0.055% 3.6-dichlorophenoxyacetic acid equivalent per top.
3.062 lbs. or 0.153% 2,4-Dichlorophenoxyacetic acid equivalent per ton.
THE 1304 He or 0.0558 3.6 diablars a spisis paid aquivalant partos

1.304 lbs. or 0.065% 3,6-dichloro-o-anisic acid equivalent per ton. \sim


Isomer Spefic by AOAC Methods.

(

TRIMEC^(R) is a registered trademark of PBI/GORDON CORPORATION



See back panel for additional Precautionary Statements.



1

A WELL-MAINTAINED LAWN BENEFITS OUR ENVIRONMENT.

- Apply when weeds are young and actively growing.
- Provides vital nutrients plus time-released nitrogen which encourages grass to fill in bare spots weeds leave behind.
- Contains sulfur which helps maintain dark green color an.. encourages plant and root growth.

GRASS, CYCLING AND LAWN BENEFITS

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients will go back into your lawn.

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

- A 50 foot x 50 foot lawn produces enough oxygen for a family of four.

- Controls erosion
- Reduces glare
- Traps dust, smoke particles and other pollutants
- Purifies water

)

- Reduces noise
- Improves soil condition
- Modifies temperature

GRASSCYCLING GRASSCYCLING The Start of Starts

2014

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. Wear long pants, long-sleeved shirt and shoes when applying this product. Wash thoroughly after handling.

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water, For eyes, flush with water for 15 minutes and eget medical attention.

ENVIRONMENTAL HAZARDS: This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by disposing of equipment washwaters. Do not apply when weather conditions favor drift from target area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. \sim

Do not reenter or permit others to reenter treated areas until sprays have dried or dusts have settled.

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or foodctuffs. To prevent cross-contamination, do not store near other fartilizers, or near herbicides insecticides, fungicides, or seeds.

PRODUCT DISPOSAL: Securely wrap bag in several layers of newspaper and discard in trash. ----CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash. _---



BEST AVAILABLE CON

- USE PRECAUTIONS -

Do not use this product where desirable clovers are present or allow the material to "wind drift" onto flowers, vegetables or ornamental shrubbery. Do not apply in areas underlaid by roots of desirable trees or shrubs. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning. Do not contaminate domestic or irrigation waters.

- INSTRUCTIONS -

ð


GORDON'S WEED & FEED FOR SOUTHERN LAWNS is a TRIMEC^(FI) complex, a combination of three proven weed killers – 2,4-D, Mecoprop and Dicamba. Together they control a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantains, henbit, spurge and many others commonly found in home lawns. In addition, a 20-10-5 fertilizer provides a generous supply of plant food needed for a normal season's growth. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days. Do not use on dichondra. Do not overapply. Undar conditions of adequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on St. Augustinegrass.

- 1. Mow lawn to normal height 1 to 2 days before application.
- 2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
- 3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto non-target plants.
- 4. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

SPREADER SETTINGS

Ì

The above spreader settings are approximate. (15 pound bag: Check your spreader on 330 square feet and adjust to apply at the rate of 1 pound per 330 square feet)(16 pound bag: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound per 310 square feet)(18 pound bag: Check you: spreader on 275 square feet and adjust to apply at the rate of 1 pound per 275 square feet)



WEEDS

Bedstraw Black Medic Buckhorn Chicory Chickweed Clover Dandelion Dock Ground ivy Heal-All Henbit Knotweed

j

)

Lambsquarter Lespedeza Mallow Morningglory Oxalis Peppergrass Pigweed Plantains Poison lvy Poison Oak Purslane Ragweed Sheep Sorrel Shepherdspurse Speedwell Spurge Thistle Wild Carrot Wild Garlic Wild Lettuce Wild Onion Yarrow

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WAR-RANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warrings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.