GORDON'S TRIMEC® WEED & FEED FOR SOUTHERN LAWNS 20-10-5

(Fifteen, Sixteen, Eighteen or Twenty) Pounds Covers Up To 5,000 Square Feet.

Controls Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and many other broadleaf weeds in St. Augustinegrass, Centipedegrass, Bahiagrass, Bermudagrass, Zoysiagrass, Bluegrass, Ryegrass, Fescue and Bentgrass lawns.

ACTIVE INGREDIENTS:

*Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy)propionic acid	0.641%
**Dimethylamine Salt of 2,4-dichlorophenoxyacetic acid	0.184%
***Dimethylamine Salt of Dicamba: 3,6-dichloro-o-anisic acid	0.078%
INERT INGREDIENTS	99.097%
TOTAL	100 000%

THIS PRODUCT CONTAINS:

- *10.609 lbs. or 0.530% 2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton.
- **3.062 lbs. or 0.153% 2,4-dichlorophenoxyacetic acid equivalent per ton.


Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional Precautionary Statements.



^{***1.304} lbs. or 0.065% 3,6-dichloro-o-anisic acid equivalent per ton.

A WELL-MAINTAINED LAWN BENEFITS OUR ENVIRONMENT.

- Apply when weeds are young and actively growing.
- Provides vital nutrients plus time-released nitrogen which encourages grass to fill in bare spots weeds leave behind.
- Contains sulfur which helps maintain dark green color and encourages plant and root growth.

GRASS CYCLING AND LAWN BENEFITS

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients will go back into your lawn.

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

- Controls erosion
- Reduces glare
- Traps dust, smoke particles and other pollutants
- · Reduces noise
- Improves soil condition
- · Modifies temperature





READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

/ First Aid

In case of contact, wash skin with plenty of soap and water. For eyes, flush with water for 15 minutes and get medical attention.

ENVIRONMENTAL HAZARDS:

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater, ,Do not apply when weather conditions favor drift from target area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do, not enter treatment area until spray has dried or dust has settled.

Note to Supplemental Registrants: Some fertilizer grades are hygroscopic, or certain formulations absorb water during storage that decrease the flowability during application. Phosphate and potash are relatively nonhygroscopic. However, fertilizer mixtures of urea and/or ammonium nitrate absorb water at lower humidities during the summer than during the winter. Caking may be prevented by maintaining low moisture contents in the fertilizers at all times. For these fertilizer formulations Options B or C may be used in place of Option A as the Storage statement in the following Storage and Disposal section.

STORAGE & DISPOSAL

OPTION A - For use with nonhygroscopic fertilizers:

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.

OPTION B - For use with hygroscopic fertilizers:

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs. To prevent caking, this product should be stored in a cool, dry place.

OPTION C - For use with hygroscopic fertilizers:

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs. Under conditions of high humidity, this product should be stored in a cool, dry place.

PESTICIDE DISPOSAL: Securely wrap bag in several layers of newspaper and discard in trash. CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

USE PRECAUTIONS:

Do not use on dichondra. Do not overapply. Under conditions of adequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on St. Augustinegrass.

Do not use this product where desirable clovers are present or allow the material to wind drift onto flowers, vegetables or ornamental shrubbery. Do not apply in areas underlaid by roots of desirable trees or shrubs. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning.

Note to Supplemental Registrants: Application schedules will vary according to climatic conditions. The following application map and estimated application schedule may be presented on the container labeling of supplemental registrations.

WHEN TO APPLY:



Northern Zone	Transition Zone	Southern Znaș
April through	Mid-March	January throagh
Mid-June	through May	April

This application map estimates the schedule of Artifizer applications and should be used as a guide-ine poly. Abnormal climatic conditions can affect your schedule.

Note to Supplemental Registrants: The following referrals to other pesticides or nonpesticides used sequentially may be presented on the container labeling of supplemental registrations.

SEQUENTIAL APPLICATIONS: Maintaining a healthy and lush green lawn involves several steps. The most important factor is an annual fertilizer application program.

ANNUAL FERTILIZER PROGRAM:

Step 1. Apply Late Winter - Early Spring.

(Brand Name) Crabgrass Preventer + Lawn Fertilizer prevents crabgrass poa annua, and other grass weeds while providing a deep green lawn with a controlled-release fertilizer.

Step 2. Apply Mid - Late Spring.

(Brand Name) Weed and Feed... kills over (number) weeds, roots and all, and provides a deep green law) with a controlled-release fertilizer.

Step 3. Apply Late Spring - Summer.

(Brand Name) Lawn Fertilizer... provides extended greening throughout the summer months with controlled-release fertilizer.

Step 4. Apply Fall.

(Brand Name) Winterizer Lawn Food (or Fail/Winterizer Weed & Feed). Is specially formulated to prepare your lawn for winter and controls weeds in one easy step. Encourages earlier spring greening.

INSTRUCTIONS:

Gordon's Trimec® Weed & Feed For Southern Lawns is a Trimec® complex, a combination of three proven weed killers — 2,4-D, Mecoprop and Dicamba. Together, they control a wide range of lawn weeds including dandelion, chickweed, knotweed, plantains, henbit, spurge and many others commonly found in home lawns. In addition, a 20-10-5 fertilizer provides a balanced supply of nutrients. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days.

The maximum application rate to turf is 0.25 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

- 1. Mow lawn to normal height 1 to 2 days before application.
- 2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
- Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants.
- 4. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

Note to Supplemental Registrants: Equipment settings may vary and may be presented on the container labeling of supplemental registrations. Illustrations or pictograms of spreader patterns may also be displayed.

Spreader Settings	
Scott	6
Lawn Beauty	5
Jackson	71/2
Sears	4

Page 4 of 6

The above spreader settings are approximate. (15 pound bag: Calibrate your spreader on 330 square feet and adjust to apply at the rate of 1 pound of product per 330 square feet) (16 pound bag: Calibrate your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet) (18 pound bag: Calibrate your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet) (20 pound bag: Calibrate your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet).

20-10-5 GUARANTEED ANALYSIS	
Total Nitrogen (N)	20%
Water Insoluble Nitrogen	5%
Available Phosphorus (P ₂ O ₅)	10%
Soluble Potash (K ₂ O)	5%

	- WEEDS -	
Bedstraw	Lambsquarters	Ragweed
Black Medic	Lespedeza	Sheep Sorrel
Buckhom	Mallow	Shepherdspurse
Chicory	Momingglory	Speedwell
Chickweed	Oxalis	Spurge
Clover	Peppergrass	Thistle
Dandelion	Pigweed	Wild Carrot
Dock	Plantains	Wild Garlic
Ground Ivy	Poison Ivy	Wild Lettuce
Healall	Poison Oak	Wild Onion
Henbit	Purslane	Yarrow
Knotweed		

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or yearbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

Page 5 of 6

APPENDIX

I. Approved fertilizer grades with nutrient contents expressed as the percentages of N, P₂O₅, and K₂O include the following:

10-6-4	22-3-3	26-3-3	28-3-3	•
15-5-7	22-3-6	26-3-5	30-3-3	
16-7-7	25-3-3	27-3-3	:	
20-10-5	25-4-6	28-2-5	_	

The fertilizer grade guaranteed analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.

- II. Guarantees offered by supplemental registrants may include the following:
 - ♦ Money-Back Guarantee:
 - (Brand Name) fertilizers are manufactured to the highest of quality standards. If you are not completely satisfied, (Company Name) will replace your product or refund your money. Call at (telephone number).
- III. Advertising claims that supplemental registrants may present on the retail container labeling.
 - (15, 16, 18, or 20) Pounds covers up to 5,000 square feet.
 - ◆ Contains Trimec[®] herbicide.
 - Controls over (number) broadleaf weeds, roots and all, including dandelion, chickweed, plantain, knotweed, henbit and spurge.
 - Can be used on bahiagrass, centipedegrass, bermudagrass, zoysiagrass, bluegrass, ryegrass, fescue and bentgrass lawns.
 - Extended feeding formula.
- IV. Note to supplemental registrants: Equipment settings may vary and may be presented on the container labeling of supplemental registrations. Illustrations or pictograms of spreader patterns may also be displayed.
 - ♦ SPREADER SETTINGS: Rotary spreaders are easier to use and reduce streaking that's often caused by drop spreaders. While drop spreaders work well, they require more care to avoid gaps and streaks which appear as light green stripes about a week after fertilizer application.
 - Always apply with a properly calibrated rotary or drop type fertilizer spreader. Never apply by hand.

ROTARY SPREADER SETTINGS	
Republic EZ Rotary	2
Scotts Easy Green	25
Quaker Rotary	31/2
Ortho Rotary (6000)	2
Ev N Spread	13
True Temper	31/2
Precision Rotary	5

DROP SPREADER SETTINGS	
Ortho Drop	5
Republic EZ Drop	5
Scotts Drop (PF3)	61/4
Scotts Even Green	61/4