

2217-661

9/29/2010

1/11



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

James Kuntsman
PBI/Gordon Corporation
P.O. Box 014090
Kansas City, Missouri 64101

SEP 29 2010

Subject: EPA Reg. 2217-661 / Gordon's Trimec Weed & Feed For Southern Lawns Notification

Dear Mr. Kuntsman:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 8-13-10 for the product EPA Reg. 2217-661 / Gordon's Trimec Weed & Feed For Southern Lawns. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions please call Erik Kraft at 703-308-9358 or email at Kraft.Erik@epa.gov or Kathryn Montague at 703-305-1243 or Montague.Kathryn@epa.gov.

Sincerely,

Kathryn Montague
Team Leader
Herbicide Branch, Team 23
Registration Division (7504P)

2
11



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 2217-661	2. EPA Product Manager Kathryn V. Montague	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Gordon's Trimec Weed & Feed For Southern Lawns	PM# Product Manager—Team 23	
5. Name and Address of Applicant (Include ZIP Code) PBI/Gordon Corporation Post Office Box 014090 Kansas City, Missouri 64101 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Labeling notification per Pesticide Registration Notice (PRN) 98-10.

We propose to add instructions for use with **drop or broadcast (rotary) spreaders**. These instructions are consistent with use directions previously approved. The methods of application, the application or dosage rate, and the application schedules have been maintained without change. Please refer to page 4 of the labeling.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or Confidential Statement of Formula of this product. I understand that it is a violation of 18 U.S.C Sec. 1001 to willfully make any false statement to the EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Section 12 and 14 of FIFRA.

e-mail to jkunstman@pbigordon.com FAX: 816-421-2731

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input checked="" type="checkbox"/> Plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per Container	<input type="checkbox"/> Glass	<input checked="" type="checkbox"/> Paper
				<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input checked="" type="checkbox"/> Container		4. Size(s) Retail Container 15, 16, 18 or 20 pound bags		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product			<input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Other _____		
			<input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)			
Name James L. Kunstman, Ph.D.	Title Director of Regulatory Services	Telephone No. (Include Area Code) 816-460-6292	
<p align="center">Certification</p> <p><i>I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.</i></p>			6. Date Application Received (Stamped)
2. Signature 	3. Title Director of Regulatory Services		
4. Typed Name James L. Kunstman, Ph.D.	5. Date August 13, 2010		


Certification with Respect to Label Integrity

version: 9/11/02

I certify that the information (including, but not limited to, text, tables, and graphics) contained in the electronic file identified below by file name and submitted with this certification is the same information as that on the paper copies of these documents included with this submission.

PROPOSED LABEL		
EPA Registration #	Date Submitted to EPA	Electronic file name
2217-661	8/13/2010	002217-00661.20100813. notif-proposed-highlighted 002217-00661.20100813. notif-proposed-clean

I certify that the statements that I have made on this form are true, accurate, and complete. I acknowledge that any knowingly false or misleading statements may be punishable by fine or imprisonment or both under applicable law.



Signature

08/13/2010

Date

James L. Kunstman, Ph.D.

Name (typed)

Director, Regulatory Services

Title

5/1

GORDON'S TRIMEC® WEED & FEED FOR SOUTHERN LAWNS

EPA Reg. No. 2217-661

ACTIVE INGREDIENT:

Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid	0.32%
Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	0.18%
Dimethylamine salt of dicamba: 3,6-dichloro-o-anisic acid	0.08%

INERT INGREDIENTS:	99.42%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- 5.40 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton or 0.27%
- 3.00 lb 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.15%
- 1.40 lb 3,6-dichloro-o-anisic acid equivalent per ton or 0.07%

Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

NOTIFICATION

SEP 29 2010

KEEP OUT OF REACH OF CHILDREN

CAUTION

- Apply when weeds are young and actively growing.
- Provides vital nutrients plus time-released nitrogen which encourages grass to fill in bare spots weeds leave behind.
- Contains sulfur which helps maintain dark green color and encourages plant and root growth.

GRASS CYCLING AND LAWN BENEFITS

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients will go back into your lawn.

Besides increasing property values, did you know a properly-maintained lawn can provide the following benefits to your home and environment:

- Helps control erosion
- Reduces glare
- Traps dust, smoke particles and other pollutants
- Reduces noise
- Helps improve soil condition
- Modifies temperature



STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

G pbi/gordon
corporation
 An Employee-Owned Company
 1217 West 12th Street
 Kansas City, Missouri 64101

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

First Aid	
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or on clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call toll-free, 1-877-800-5556 for emergency medical treatment information.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate. This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

[Environmental Hazards (alternate per PRN 2008-1)

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.]

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Do not use on dichondra. Do not overapply. Under conditions of adequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on St. Augustinegrass.
- Do not use this product where desirable clovers are present. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning.
- Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result.

- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

[Note to Supplemental Registrants: Application schedules will vary according to climatic conditions. The following application map and estimated application schedule may be presented on the container labeling of supplemental registrations.]

WHEN TO APPLY:



Northern Zone	Transition Zone	Southern Zone
April through Mid-June	Mid-March through May	January through April

This application map estimates the schedule of fertilizer applications and should be used as a guideline only. Abnormal weather conditions can affect your schedule.

[Note to Supplemental Registrants: The following referrals to other pesticides or nonpesticides used sequentially may be presented on the container labeling of supplemental registrations.]

SEQUENTIAL APPLICATIONS: Maintaining a healthy and lush green lawn involves several steps. The most important factor is an annual fertilizer application program.

ANNUAL FERTILIZER PROGRAM:

Step 1. Apply Late Winter - Early Spring.

(Brand Name) Crabgrass Preventer + Lawn Fertilizer... prevents crabgrass, poa annua, and other grass weeds while providing a deep green lawn with a controlled-release fertilizer.

Step 2. Apply Mid - Late Spring.

(Brand Name) Weed and Feed... kills over (number) weeds, roots and all, and provides a deep green lawn with a controlled-release fertilizer.

Step 3. Apply Late Spring - Summer.

(Brand Name) Lawn Fertilizer... provides extended greening throughout the summer months with controlled-release fertilizer.

Step 4. Apply Fall.

(Brand Name) Winterizer Lawn Food (or Fall/Winterizer Weed & Feed)... prepares your lawn for winter and controls weeds in one easy step. Encourages earlier spring greening.

INSTRUCTIONS:

Gordon's Trimec® Weed & Feed For Southern Lawns is a Trimec® complex, a combination of three proven weed killers — 2,4-D, Mecoprop-p and Dicamba. Together, they control a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a 20-10-5 fertilizer provides a supply of nutrients. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in 30 days.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).

- 3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
- 4. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

[Note to Supplemental Registrants: Equipment settings may vary and may be presented on the container labeling of supplemental registrations. Illustrations or pictograms of spreader patterns may also be displayed.]

Spreader Settings	
Scotts	6
Lawn Beauty	5
Jackson	7.5
Sears	4

Application Equipment and Instructions

This product can be applied with a drop or broadcast (rotary) spreader. For best results, apply this product evenly and uniformly over the turfgrass (lawn) area avoiding streaking, skips or overlaps.

Before each application, determine the correct amount (lbs.) of product to place in the spreader hopper for the area to be treated. For example, if the label states to apply 3.6 lbs. per 1000 sq. ft. and your area is 5000 sq. ft., then place 18 lbs. of product in the hopper and set the spreader setting at 4 (per table below) if using an Accugreen drop spreader.

A more uniform application can be made by spreading one-half of the recommended amount of this product over the turfgrass area and then applying the remaining one-half of this product at a right angle to the previous application direction.


NOTE: The spreader settings listed below are approximate. The speed at which you walk and the working condition of the spreader will affect the rate of application.

[These settings can only be used as a guide; they are not absolute. The speed at which the applicator walks, spreader condition and the texture of the turf will alter rate of application.]

The above spreader settings are approximate.

- 15 pound bag:** Calibrate your spreader on 330 square feet and adjust to apply at the rate of 1 pound of product per 330 square feet.
- 16 pound bag:** Calibrate your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
- 18 pound bag:** Calibrate your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- 20 pound bag:** Calibrate your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- 20 pound bag:** Calibrate your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

[Spread product over area going at right angles (see diagram).]



The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application (0.26 lb 2,4-D ae, 0.47 lb MCP-p ae and 0.12 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. (0.52 lb 2,4-D ae, 0.94 lb MCP-p ae and 0.24 lb dicamba ae per acre).

20-10-5 GUARANTEED ANALYSIS	
Total Nitrogen (N)	20%
Water Insoluble Nitrogen	5%
Available Phosphorus (P2O5)	10%
Soluble Potash (K2O)	5%

Labeling Requirements:

- Information regarding the contents and levels of metals in this product is available on the Internet as <http://www.aapfco.org/metals.htm>.

BROADLEAF WEEDS			
Annual yellow sweetclover	Cockle	Mustard	Spiny amaranth
Aster	Cocklebur	Narrowleaf plantain	Spiny cocklebur
Austrian fieldcress	Common mullein	Narrowleaf vetch	Spiny sowthistle
Bedstraw	Creeping jenny	Nettle	Spotted catsear
Beggaricks	Cudweed	Orange hawkweed	Spurweed
Betony, Florida	Curly dock	Oriental cocklebur	Stinging nettle
Bindweed, field	Daisy, English	Oxalis	Strawberry, India mock
Bird vetch	Daisy fleabane	Parsley-piert	Tall nettle
Bitter wintercress	Daisy, oxeye	Parsnip	Tall vervain
Bittercress, hairy	Dandelion	Pearlwort	Tansy ragwort
Bitterweed	Dichondra	Pennycress	Tansy mustard
Black-eyed Susan	Dogbane	Pennywort	Tanweed
Black medic	Dogfennel	Peppergrass	Thistle
Black mustard	Dollarweed	Pepperweed	Trailing crownvetch
Blackseed plantain	False dandelion	Pigweed	Tumble mustard
Blessed thistle	Falseflax	Pigweeds bedstraw	Tumble pigweed
Bloodflower milkweed	False sunflower	Plains coreopsis (tickseed)	Velvetleaf
Blue lettuce	Fiddleneck	Plantain	Venice mallow
Blue vervain	Florida pusley	Poison ivy	Virginia buttonweed
Bracted plantain	Frenchweed	Poison oak	Virginia creeper
Brassbuttons	Galinsoga	Pokeweed	Virginia pepperweed
Bristly oxtongue	Goathead	Poorjoe	Water pennywort
Broadleaf dock	Goldenrod	Prairie sunflower	Wavyleaf bullthistle
Broadleaf plantain	Ground ivy	Prickly lettuce	Western clematis
Broomweed	Gumweed	Prickly sida	Western salsify
Buckhorn	Hairy fleabane	Prostrate knotweed	White mustard
Buckhorn plantain	Hawkweed	Prostrate pigweed	Wild aster
Bulbous buttercup	Healall	Prostrate spurge	Wild buckwheat
Bull thistle	Heartleaf drymary	Prostrate vervain	Wild carrot
Bullnettle	Heathaster	Puncturevine	Wild four-o'clock
Burclover	Hedge bindweed	Purslane, common	Wild garlic
Burdock	Hedge mustard	Ragweed	Wild geranium
Burning nettle	Hemp	Red sorrel	Wild lettuce
Burweed	Henbit	Redroot pigweed	Wild marigold
Buttercup	Hoary cress	Redstem filaree	Wild mustard
Buttonweed	Hoary plantain	Rough cinquefoil	Wild onion
Canada thistle	Hoary vervain	Rough fleabane	Wild parsnip
Carolina geranium	Horsenettle	Russian pigweed	Wild radish
Carpeweed	Jimsonweed	Russian thistle	Wild rape
Catchweed bedstraw	Knawel	Scarlet pimpernel	Wild strawberry
Catnip	Knotweed	Scotch thistle	Wild sweet potato
Catsear	Kochia	Sheep sorrel	Wild vetch

BROADLEAF WEEDS			
Chickweed, common	Lambsquarters	Shepherdspurse	Woodsorrel
Chickweed, mouseear	Lespedeza	Slender plantain	Woolly croton
Chicory	Mallow	Smallflower galinsoga	Woolly morningglory
Cinquefoil	Matchweed	Smooth dock	Woolly plantain
Clover, crimson	Mexicanweed	Smooth pigweed	Wormseed
Clover, hop	Milk vetch	Sorrel	Yarrow
Clover, red	Momingglory	Sowthistle	Yellow rocket
Clover, strawberry	Mouseear hawkweed	Spanishneedles	Yellowflower pepperweed
Clover, sweet	Mugwort	Speedwell	and other broadleaf weeds
Clover, white	Musk thistle	Spotted spurge	

[Note to Supplemental Registrants: Some fertilizer grades are hygroscopic, or certain formulations absorb water during storage that decrease the flowability during application. Phosphate and potash are relatively nonhygroscopic. However, fertilizer mixtures of urea and/or ammonium nitrate absorb water at lower humidities during the summer than during the winter. Caking may be prevented by maintaining low moisture contents in the fertilizers at all times. For these fertilizer formulations Options B or C may be used in place of Option A as the Storage statement in the following Storage and Disposal section.]

STORAGE AND DISPOSAL
 Do not contaminate water, food, or feed by storage and disposal.

OPTION A - For use with nonhygroscopic fertilizers:
PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

OPTION B - For use with hygroscopic fertilizers:
PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. To prevent caking, this product should be stored in a cool, dry place.

OPTION C - For use with hygroscopic fertilizers:
PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Under conditions of high humidity, this product should be stored in a cool, dry place.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partially filled:** Call your local solid waste agency or toll free number (1-800-xxx-xxxx) for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER
 FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

APPENDIX

1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Weight: _____
- EPA Est. No. _____

2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

- (15, 16, 18, or 20) Pounds covers up to 5,000 square feet.
- Contains Trimec® herbicide.
- Controls over (number) broadleaf weeds, roots and all, including dandelion, chickweed, plantain, knotweed, henbit and spurge.
- Controls dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf weeds in St. Augustinegrass, centipedegrass, bahiagrass, bermudagrass, zoysiagrass, bluegrass, perennial ryegrass, fescue spp. and bentgrass lawns.
- Can be used on bahiagrass, centipedegrass, bermudagrass, zoysiagrass, bluegrass, ryegrass, fescue and bentgrass lawns.
- Extended feeding formula.

3. Note to supplemental registrants: Equipment settings may vary and may be presented on the container labeling of supplemental registrations. Illustrations or pictograms of spreader patterns may also be displayed.

- **SPREADER SETTINGS:** Rotary spreaders are easier to use and reduce streaking that's often caused by drop spreaders. While drop spreaders work well, they require more care to avoid gaps and streaks which appear as light green stripes about a week after fertilizer application.
- Always apply with a properly calibrated rotary or drop type fertilizer spreader. Never apply by hand.

ROTARY SPREADER SETTINGS		DROP SPREADER SETTINGS	
Republic EZ Rotary	2	Ortho Drop	5
Scotts Easy Green	25	Republic EZ Drop	5
Quaker Rotary	3.5	Scotts Drop (PF3)	6.25
Ortho Rotary (6000)	2	Scotts Even Green	6.25
Ev N Spread	13		
True Temper	3.5		
Precision Rotary	5		

4. Guarantees offered by supplemental registrants may include the following:

- Money-Back Guarantee:
- (Brand Name) fertilizers are manufactured to the highest of quality standards. If you are not completely satisfied, (Company Name) will replace your product or refund your money. Call at (telephone number).

5. Approved fertilizer grades with nutrient contents expressed as the percentages of N, P2O5, and K2O include the following:

10-6-4	20-10-5	25-4-6	28-2-5
15-5-7	22-3-3	26-3-3	28-3-3
15-5-10	22-3-6	26-3-5	30-3-3
16-7-7	25-3-3	27-3-3	

The fertilizer grade guaranteed analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.

DOCUMENT CONTROL INFORMATION

1. **Unique Label Identifier:** 002217-00661.20100813.notif-proposed-highlighted.doc
2. **Reason for Issue:** restatement of spreader information, PRN 2007-4