

2217-668

5-20-2005

1/11



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

MAY 20 2005

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Mr. Craig Martens
Manager of Regulatory Services
PBI/Gordon Corporation
Post Office Box 014090
Kansas City, Missouri 64101

Dear Mr. Martens:

Subject: Gordon's Amine 600 2,4-D Weed Killer
Registration No. 2217-668
Application and Letter Dated May 5, 2005, Request
to Amend Registration To Add Maximum Use Rates for
Forestry and Non-Crop Sites To Label, Based on
A Memorandum of Understanding Agreed to between
the Industry Task Force II on 2,4-D Research Data on
October 14, 2004

The labeling referred to above has been reviewed and found acceptable as an amendment to the registration of "Amine 400 2,4-Weed Killer" under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, provided that you:

1. Revise the footnotes for Table 6 by deleting the word "general" as a descriptor to noncropland sites, and in last statement change the name "non-crop sites" to read "noncropland sites".
2. Do not release this product for shipment under any other labeling after October 1, 2005.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA, section 6(e). Your release for shipment of this product under this labeling constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records. Please submit one copy of the final printed label before you release the product for shipment.

Sincerely yours,

Joanne I. Miller

Joanne I. Miller
Product Manager 23
Herbicide Branch
Registration Division (7505C)

Enclosure

2/11

GORDON'S AMINE 600 2,4-D WEED KILLER

ACCEPTED
with **COMMENTS**
in EPA Letter Dated:

MAY 20 2005
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

2217-668

ACTIVE INGREDIENT:

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid 66.57%

INERT INGREDIENTS: 33.43%

TOTAL 100.00%

THIS PRODUCT CONTAINS:

5.6 lbs. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 55.30%.

Isomer Specific by AOAC Methods.

KEEP OUT OF REACH OF CHILDREN

DANGER - PELIGRO

Si Usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle. (If you do not understand the label, find some one to explain it to you in detail.)

See additional Precautionary Statements and First Aid elsewhere on this label.

NET CONTENTS (1, 2.5, 30, or 55) U.S. GALLONS

802/ APxxxxxx
EPA REG. NO. 2217-668
EPA EST. NO. 2217-KS-1
MANUFACTURED BY:



Telephone: 1-800-821-7925

3/11



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

DANGER: Corrosive. Causes irreversible eye damage. Causes skin irritation. Do not get in eyes, on skin, or on clothing. Harmful or fatal if swallowed.

Personal Protective Equipment (PPE):

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, and shoes plus socks, protective eyewear, and chemical resistant apron when cleaning equipment, mixing, or loading.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

Engineering control statements:

Containers over 1 gallon and less than 5 gallons: Mixers and loaders who do not use a mechanical system (probe and pump) to transfer the contents of this container must wear coveralls or a chemical resistant apron in addition to the other required PPE.

Containers of 5 gallons or more: A mechanical system (probe and pump) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal. If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40CFR 170.240 (d)(4)], the handler PPE requirements may be reduced or modified as specified in the WPS.

When handlers use enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

Users should:

- ◆ Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- ◆ Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- ◆ Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

4/11

First Aid

If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.</p> <p>NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.</p>	

ENVIRONMENTAL HAZARDS:

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water except as specified on this label. Do not contaminate water when disposing of equipment washwater. Do not apply when weather conditions favor drift from target area. Do not contaminate domestic or irrigation waters.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170.

This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

For early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear: coveralls, waterproof gloves, shoes plus socks, and protective eyewear.

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Gordon's Amine 600 2,4-D Weed Killer may be stored in an unheated building. However, product exposed to subfreezing temperatures should be warmed to at least 45°F and mixed thoroughly before using.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: For Metal Containers - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures. For Plastic Containers - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities by burning. If burned stay out of smoke.

USE PRECAUTIONS:

Do not overdose! Avoid spray drift to cotton, soybeans, tomatoes, tobacco, grapes, fruit trees, flowers, garden crops, ornamental plants, shrubs, trees and other hormone herbicide-sensitive desirable plants. Do not apply near these plants since small quantities of wind-drifted herbicide may cause severe injury. Do not apply when wind speed is sufficient to cause drift. Do not apply when temperature air inversion exists. An air inversion may be detected by creating a smoke column and observing a layering effect. Do not apply when temperature exceeds 90°F and humidity is high.

To aid in avoiding spray drift, use coarse sprays and low pressure. Do not use nozzles which produce fine spray droplets under high pressure. The use of thickening agents or anti-drift additives and drift reducing equipment is of value in preventing spray drift.

Care should be taken not to make applications where runoff could carry the chemical to nontarget areas. Local spray conditions will vary and check local recommendations.

Do not apply this product through any type of irrigation system.

WEEDS CONTROLLED

Annual and Biennial Weeds:		
Beggarticks	Jewelweed	Puncturevine
Bitterweed	Jimsonweed	Radish (wild)
Broomweed	Kochia	Ragweed
Bull Thistle	Knotweed	Russian thistle
Burdock	Lambsquarters	Shepherdspurse
Carpetweed	Lettuce (wild)	Smartweed
Cinquefoil	Mallow	Sneezeweed
Cockle	Marshelder	Sowthistle (common)
Cocklebur	Marijuana	Spanishneedles
Coffeeweed	Morningglory (annual)	Sunflower
Croton	Musk thistle	Tumbleweed
Devil's claw	Mustard	Velvetleaf
Fleabane (daisy)	Parsnip	Vetch
Flixweed	Pennycress	Wild carrot
Frenchweed	Peppergrass	Wild parsnip
Galinsoga	Pigweed	Witchweed
Goatsbeard	Prickly lettuce	Wormwood

6/11

Goosefoot	Primrose	Yellow starthistle
Perennial Weeds:		
Artichoke	Goldenrod	Rushes
Aster	Ground ivy	Southern wild rose
Austrian fieldcress	Gumweed	Sowthistle
Bindweed	Healall	Stinging nettles
Black eyed Susan	Hoary cress	Strawberry (wild)
Blue lettuce	Horsetail	Tall buttercup
Canada thistle	Ironweed	Tanweed
Catnip	Locoweed	Toadflax
Chicory	Nettles	Vervains
Clover (many types)	Orange hawkweed	Wild garlic
Dandelion	Plantain	Wild onion
Docks	Poverty weed	Wild sweet potato
Dogbane	Yellow rocket	

SPRAY PREPARATION:

Fill the spray tank with one-half the required amount of water, then add the recommended amount of product and continue filling the spray tank. Keep agitator running when filling the spray tank and during spray operations. Use enough spray volume for uniform coverage by ground or air application. In any case, use the same amount of 2,4-D recommended per acre. For crop uses, DO NOT MIX WITH OIL, SURFACTANT OR OTHER ADJUVANTS UNLESS SPECIFICALLY RECOMMENDED ON THIS LABEL AS THIS MAY REDUCE SELECTIVITY TO CROPS RESULTING IN CROP DAMAGE.

WHEN TO USE:

Best results are obtained when weeds are young and actively growing. Applications of lower recommended rates to susceptible annual weeds usually will be satisfactory, but for perennial weeds and other conditions, such as in very dry areas where weed kill is difficult, use the higher recommended rates. When used as a selective spray on crops, the stage of growth of the crop must be considered. Some woody plants and weeds are hard to kill and repeat applications may be necessary.

7/11

Table 1. Recommendations For Use In Corn And Sorghum By Air Or Ground Applications.		
Crop	Amount Per Acre	Directions
Corn (Preemergence)	1 ¹ / ₃ to 2 ² / ₃ Pints	Spray after planting before corn emerges, preferably 2 to 5 days after planting.
Corn (Emergence)	² / ₃ Pint	Apply just as corn plants are breaking ground.
Corn (Postemergence)	¹ / ₃ to ² / ₃ Pint	Spray when weeds are small and crop is 4 to 12 inches tall. Use drop nozzles when corn is more than 8 inches tall or beyond the 5 leaf stage to keep the spray off the corn foliage.
	¹ / ₂ to ² / ₃ Pint	Application of ¹ / ₂ to ² / ₃ pt. per acre may be needed to control resistant weeds but crop injury is more likely.
Sweet Corn	¹ / ₃ to ² / ₃ Pint	Spray before sweet corn is 6 inches in height.
Corn (Preharvest)	² / ₃ to 1 ¹ / ₃ Pints	Apply after the hard dough or denting stage by air or ground equipment to suppress perennial weeds, and control tall weeds such as bindweed, cocklebur, dogbane, jimsonweed, ragweed, sunflower, velvetleaf and vines that interfere with harvesting. Do not forage or feed corn fodder for 7 days following application.
Grain Sorghum (Milo) Postemergence	¹ / ₃ to ² / ₃ Pint	Spray when weeds are small and crop is 6 to 8 inches tall. Use drop nozzles when sorghum is more than 8 inches tall or during boot, flowering or early dough stage.

Footnotes For Table 1: Do not use atrazine, oil or other adjuvants. Do not apply from the tasseling to dough stage in corn. Crop injury is more likely to occur if corn or sorghum is growing rapidly under high temperature and high moisture conditions. In such cases use the lower rate. To reduce stalk breakage from temporary brittleness caused by 2,4-D, delay cultivation for 8 to 10 days after treatment. Hybrids vary in response to 2,4-D and some are easily injured. Spray only hybrids known to be tolerant to 2,4-D. Contact seed company and Extension Service weed specialists for this information.

FOR USE IN REDUCED OR NO-TILLAGE IN SOYBEANS (Preplant Only)

GENERAL INFORMATION:

Gordon's Amine 600 2,4-D Weed Killer is a phenoxy-type herbicide that provides postemergence control of many susceptible annual and perennial broadleaf weeds. Gordon's Amine 600 2,4-D Weed Killer may be applied prior to planting soybeans to provide foliar burndown control of susceptible annual and perennial broadleaf weeds and certain broadleaf cover crops such as those listed on this label. Gordon's Amine 600 2,4-D Weed Killer should only be applied preplant to soybeans in situations such as reduced tillage production systems, where emerged weeds are present. Apply only according to the application instructions given below.

MIXING INSTRUCTIONS:

Mix Gordon's Amine 600 2,4-D Weed Killer only with water, unless otherwise directed on this label. Compatible crop oil concentrates, agricultural surfactants and fluid fertilizers approved for use on growing crops may increase the herbicidal effectiveness of 2,4-D on certain weeds and may be added to the spray tank. Read and follow all directions and precautions on this label and on all labels of adjuvants or fertilizers mixed with this product.

APPLICATION PROCEDURES:

Apply using air or ground equipment in sufficient gallonage to obtain adequate coverage of weeds. Use 2 or more gallons of water per acre in aerial equipment and 10 or more gallons of water per acre in ground equipment.

8/11

APPLICATION TIMING AND USE RATES FOR AMINE SALTS

Maximum Amount Of Amine 600 to Apply/Acre	Maximum Rate (Pounds 2,4-D to Apply/Acre)	When to Apply (Days Prior to Planting Soybeans)
2/3 Pint	0.5	Not Less Than 15 Days
2/3 Quart	1.0	Not Less Than 30 Days

WEEDS CONTROLLED		
alfalfa*	garlic, wild*	ragweed, common
bindweed*	horseweed or marestalk	ragweed, giant
bullnettle	Ironweed	shepherdspurse
bittercress, small flowered	lambsquarters, common	smartweed, Pennsylvania
buttercup, small flowered	lettuce, prickly	sowthistle, annual
Carolina geranium	morningglory, annual	speedwell
cinquefoil, common and rough	Mousetail	thistle, Canada*
clover, red*	mustard, wild	thistle, bull
cocklebur, common	onion, wild*	velvetleaf
dandelion	pennycress, field	vetch, hairy*
dock, curly*	Plantain	Virginia copperleaf
evening primrose, cutleaf	purslane, common	

*These species are only partially controlled.

In general, weeds should be small, actively growing and free of stress caused by extremes in climatic conditions, diseases, or insect damage at the time of treatment. The response of individual weed species to Gordon's Amine 600 2,4-D Weed Killer is variable. Consult your local county or state Agricultural Extension Service or crop consultant for advice.

APPLICATION RESTRICTIONS AND PRECAUTIONS FOR SOYBEANS (PREPLANT)

IMPORTANT NOTICE: Unacceptable injury to soybeans planted in fields previously treated with Gordon's Amine 600 2,4-D Weed Killer may occur. Whether or not soybean injury occurs and the extent of the injury will depend on weather (temperature and rainfall) from herbicide application until soybean emergence and agronomic factors such as the amount of weed vegetation and previous crop residue present. Injury is more likely under cool rainy conditions and where there is less weed vegetation and crop residue present.

Apply a maximum of one application per growing season regardless of the treatment rate.

Do not apply Gordon's Amine 600 2,4-D Weed Killer when weather conditions such as temperature air inversions or wind favor drift from treated areas to susceptible plants.

Livestock Grazing Restriction: Do not feed hay, forage, or fodder. Restrict livestock from grazing treated fields.

In fields previously treated with 2,4-D, plant soybean seed as deep as practical or at least 1.5 to 2.0 inches deep. Adjust the press wheel of the planter, if necessary, to ensure that planted seed is completely covered.

9/11

Table 2. Recommendations For Use In Small Grains And Sugarcane With Aerial Or Ground Application.

Crop	Amount Per Acre	Directions
Wheat, Oats, Barley, Rye (fall seeded)	2/3 to 1 Pint	Apply in the spring after full tiller but before boot stage. Do not treat in fall. Do not apply from boot to milk stage or in seedling stage. Do not forage or graze treated grain fields within 2 weeks after treatment with 2,4-D. Do not feed treated straw to livestock.
	2/3 to 1 1/3 Pints	For special situations such as wild onion or wild garlic. These rates may injure the crop.
Wheat, Barley, Rye (spring-seeded)	2/3 to 1 Pint (1 1/2 to 2 Pints for special situations)	Apply after full tiller (usually 4 to 8 inches) but before boot stage. Do not apply from boot to milk stage. Do not forage or graze treated grain fields within 2 weeks after treatment with 2,4D. Do not feed treated straw to livestock.
Oats (spring-seeded)	1/3 to 2/3 Pints (1 to 1 1/3 Pints for special situations)	Apply at the full tiller stage but before the early boot stage. Do not apply from boot to milk stage. Do not forage or graze treated grain field within 2 weeks after treatment with 2,4-D. Do not feed treated straw to livestock. Oats are less tolerant to 2,4-D than wheat or barley.
Wheat, Oats, Barley, Rye (Preharvest)	2/3 to 1 1/3 Pints (1 1/3 to 2 Pints for special situations)	Apply when grains are in the hard dough stage to control large weeds that may interfere with harvest. Best results are obtained when weeds are growing vigorously. Do not feed treated straw to livestock.
Oats for grain Southern U.S. (fall seeded)	1/2 to 1 Pints	Apply after full tillering but before the early boot stage. Do not spray during or immediately following cold weather. Do not forage or graze treated fields within 2 weeks after treatment with 2,4-D. Do not feed treated straw to livestock.
Grain stubble for wild garlic	1 1/3 to 2 Quarts	Apply following harvest. Do not forage for 7 days following application. Do not plant to any crop until 3 months after treatment or until chemical has disappeared from soil.
Rice	2/3 to 2 Pints	Apply when the first elongating internode increases to 1/2 inch long. Do not apply when internode exceeds 1/2 inch long. Do not apply nitrogen during 7 to 21 days before application of 2,4-D. Follow State Extension Service recommendations.
Sugarcane	1 1/3 to 2 2/3 Pints	Spring, once or twice before dose-in fall after harvest or planting.

Footnotes for Table 2:
 The use of the higher rates in special situations to handle difficult weed problems is more likely to cause crop damage. Oats are less tolerant to 2,4-D than wheat or barley and more susceptible to injury. Do not feed treated straw to livestock. When using high rates in special situations consult your state weed specialist for recommendations to fit local conditions.

Table 3. Recommendations For Use In Rangeland And Grass Pastures

Weeds & Crops	Amount Per Acre	Directions
Weeds & Brush	2/3 to 1 1/3 Quarts	Apply when weeds are small and actively growing. Spray perennials when in the bud stage. The lower rate can be used for susceptible species.
Newly sprigged Coastal Bermudagrass	2/3 to 1 1/3 Quarts	Spray Pre-emergence and/or postemergence.
Musk Thistle	1 Quart	In spring in rosette stage before stalk lengthens and plants bolted.
	1 1/3 Quarts	In spring after bolting or in late fall or winter.

Footnotes for Table 3:
 The maximum application rate to pasture and rangeland is 2 pounds 2,4-D acid equivalent per acre per application per site. Legumes such as clover or alfalfa may be damaged. Do not graze dairy cattle in treated areas for 7 days after application. Remove meat animals from freshly treated areas for 3 days before slaughter. Do not cut treated grass for hay within 30 days after application. Do not apply to newly seeded areas or after heading begins. Do not apply to grass when in boot to milk stage. Do not apply to bentgrass or susceptible grass pastures. If weeds are young and growing vigorously, 2/3 quart per acre will provide control of some species. Deep-rooted perennial weeds may require repeated applications in the same year or in subsequent years

Table 4. Recommendations For Use In Grass Seed Crops.

Crops	Amount Per Acre	Directions
Grass seed crops	2/3 to 1 Pint	Use sufficient water to give uniform application by ground or air. Apply to established stands in spring from the tiller to early boot stage. Can be applied in spring or fall.

Footnotes for Table 4:
 Do not spray from early boot stage to milk stage. New spring seedlings can be treated with the lower rate after the grasses have at least five leaves. Perennial weed regrowth can be treated in the fall. Do not use on bentgrass unless injury can be tolerated. For best results apply when soil moisture is adequate.

Table 5. Recommendations For Use In Fallow Lands.

Site	Amount Per Acre	Directions
Fallow land	2/3 to 1 1/3 Quarts	For established annual broadleaf weeds.
	2/3 to 2 Quarts	For established perennial species such as Canada thistle or field bindweed.

Footnotes for Table 5:
 Apply to actively growing weeds. Do not plant any crop for 3 months or until chemical has disappeared from the soil. Consult your state weed specialist for information on local conditions.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

11/11

Table 6. Recommendations For Use In Noncropland Areas		
Site	Amount Per Acre	Directions
Fencerows, lawns, airfields, roadsides, vacant lots, drainage ditches.	1 to 2 Quarts	Use the amount of water needed for uniform application. Usually 1½ quarts provides good weed control under average conditions when weeds are growing.
<p>Footnotes for Table 6:</p> <p>Treat when weeds are young and growing well. Do not use on golf greens nor on dichondra or other broadleaf herbaceous groundcovers. Do not use on creeping grasses such as bentgrass and St. Augustinegrass except for spot treatment. Do not use on newly seeded turf until grass is well established. Reseeding of treated areas should be delayed following treatment. With spring application, reseed in the fall; with fall application, reseed in the spring. Legumes are usually damaged or killed. Do not treat areas where legumes are desired. Deep-rooted perennial weeds may require repeated treatments in the same season or in the subsequent years.</p> <p>The maximum application rate to general non-crop sites is 2 pounds of 2,4-D acid equivalent per acre per application per site. The maximum non-crop application for tree, brush and woody plant control on rights-of-way is 4 pounds of 2,4-D acid equivalent per acre per application per site. The maximum seasonal application rate to non-crop sites is 4 pounds 2,4-D acid equivalent per acre per application per site.</p>		

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.