



READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals:

DANGER: Corrosive causes eye damage and skin irritation. Do not get in eyes, on skin or clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed.

Environmental Hazards

Do not apply directly to water. Do not apply when weather conditions favor drift away from target area. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate domestic or irrigation waters.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

— KEEP FROM FREEZING —

Prohibitions: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Do not store near other pesticides or seeds.

Pesticide Disposal: Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under Subtitle C of the Resource Conservation and Recovery Act.

Container Disposal: Triple rinse (or equivalent) and offer for recycling or dispose of in a sanitary landfill, or by other approved State and local procedures.

**GORDON'S
PHENOMETMC 2+1**

FOR SALE TO, USE AND STORAGE BY
PROFESSIONAL TURF MAINTENANCE PERSONNEL
ONLY.

For the control of broadleaf weeds in bluegrass,
fescue, Bermudagrass and bentgrass turf.

A 2:1 formula... CMPP plus 2,4-D in a mixed amine solution.

— KEEP FROM FREEZING —

ACTIVE INGREDIENTS:

*Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	16.16%
*Diethanolamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	12.20%
**Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid	8.04%
**Diethanolamine Salt of 2,4-Dichlorophenoxyacetic acid	6.04%
INERT INGREDIENTS	57.56%
TOTAL	100.00%

This Product Contains:

- *2 lbs. 2-(2-methyl-4-chlorophenoxy) propionic acid per gallon or 21.53%
- *1 lb. 2,4-Dichlorophenoxyacetic acid per gallon or 10.77%
- **Isomer Specific by AOAC Method 6 D01-5

KEEP OUT OF REACH OF CHILDREN

DANGER

Statement of Practical Treatment

- If In Eyes:** In case of eye contact immediately flush eyes with plenty of water for 15 minutes. Call a physician at once.
- If On Skin:** Wash promptly with soap and water. Rinse thoroughly. If irritation develops, get medical attention.
- If Swallowed:** Drink one or two glasses of water. Induce vomiting by touching back of throat with finger. Call a physician at once. Do not induce vomiting or give anything by mouth to an unconscious person.
- If Inhaled:** Remove victim to fresh air and apply respiration if indicated.

See additional precautionary information elsewhere on this label.

NET CONTENTS ONE U.S. GALLON

BEST DOCUMENT AVAILABLE



USE PRECAUTIONS:

- Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near non-target plants. Do not use on warm season grasses such as St. Augustine, Centipede, Dichondra or on lawns and turf with desirable clovers are present. Note: Coarse sprays are less likely to wind-drift. Do not spray when air temperature exceeds 85 degrees F. Seed can be safely sown 3 to 4 weeks after application. Do not contaminate domestic or irrigation waters. After using this product, clean sprayer with soap or detergent and water and rinse thoroughly before applying other pesticides. Failure to observe the above precautions may result in injury.

GENERAL INFORMATION AND INSTRUCTIONS:

PHENOMETMC 2+1 is a mixed amine selective herbicide for the control of broadleaf weeds in bluegrass, fescue, bermudagrass, and bentgrass turf. Do not use this product on closely-mowed bentgrass or bermudagrass as on golf greens or tees. Weeds are best controlled when they are young and actively growing.

Weeds Controlled:

Susceptible — Dandelion, clover, common chickweed, mouse ear chickweed, plantain, pigweed, ragweed and lambsquarter.
Moderately Susceptible — Knotweed, black medic, English daisy, dock, purslane, oxalis, mallow.

Application Rate:

For established turf, apply 1-1/4 oz. to 1-1/2 oz. for 1,000 sq. ft. (3-1/3 pints to 4 pints per acre) in 1 to 2 gallons of water per 1,000 sq. ft. (30 to 60 gallons per acre). Use higher rates when hard-to-kill weeds are present. Maximum control of weeds will be obtained from spring or early fall application when weeds are actively growing. Avoid applying during periods when turf is under stress from drought or extreme heat. Do not water for 24 hours after application. Do not apply to newly seeded turf until after the second or third mowing.

NOTE: Use application equipment that is capable of applying a uniform droplet over the entire area to be covered.

WARRANTY:

The manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled use when applied according to label directions. Because of widely varying use conditions, it is impossible to eliminate all risks even when label directions are followed. Therefore, the manufacturer makes no other implied or express warranty nor is any agent of the manufacturer allowed to do so. Upon purchase of this product, the buyer agrees to accept a refund of the purchase price of the product as full discharge of the manufacturer's liability.

EPA REG. NO. 2217-671
EPA EST. NO. 2217-KS-1
837/582



ACCEPTED
NOV 30 1982
EPA REG. NO. 2217-671