

# USE PRECAUTIONS:

Use only on the plants listed on the label. Do not apply to transplant or cutting beds or to newly set transplants, and do not seed or transplant into treated areas. Do not use on extremely light sandy soils. Do not use on plants, any part of which is used for animal consumption. This product, when applied to the soil as directed, will give selective control of annual grassy and broadleaved weeds and certain perennial weeds such as:

Annual Bluegrass	Foxtail	Plantain
Bluegrass	Gisela	Purslane
Bull Thistle	Groundsel	Ragweed
Camphorweed	Henbit	Redroot Pigweed
Carpetweed	Horsetail	Rosarypea
Chickweed	Jerusalem	Russian Thistle
Citron Melon	Goosefoot	Shepherdspurse
Coffeeweed	Knotweed	Smartweed
Crabgrass	Lambquarters	Spanishneedles
Cudweed	Milkweed Vine	Spurge
Dandelion	(Morone)	Teaweed
Evening Primrose	(Morone)	Texas Panicum
Fiddleneck	Natalgrass	(Harradgrass)
Florida Purslane	Pepperglass	Wild Mustard
(Pusley)	Pineapple Weed	Yellow Woodsorrel

## WOODY ORNAMENTALS

This product is to be applied only to established Woody Plants - Those which have been planted for at least four months. May be used for weed control on the following plants.

Amur Corktree	Filbert	Mountain Ash
Arbor Vitae	Flowering Almond	Nanadina
Ash	Flowering Quince	Oak
Azalea*	Forsythia	Osmanthus
Barberry	Gardenia	(Holly-Olive)
Beauty Bush	Goldenraintree	Pachistima
Birch	Hackberry	Pecan
Boxelder	Heather	Photinia
Boxwood	Holly*	Pittosporum
Camellia	Honeysuckle	Poplar
Caragana	Ivy (Hedera)	Privet
Cleyera Japonica	Holly	Pyracantha
Cotoneaster	Juniper	Rhododendron
Cottonwood	Kinnikinnick	Rockrose
Crabapple	Laurel	Rose
Deutzia	Leucothoe	Russian Olive
Dogwood	Lilac	Spiraea
Eastern Red Cedar	Linden	Square-carpel
Elm	Locust	Wetgalla
Elwood Cypress	Magnolia	Willow
English Walnut	Maple	Yew (Taxus)
Eunonymus	Mock Orange	

1. Kurume, Mollis, Hardy Hybrid types and hardy native species only.
2. Except I. Crenata, I. Rotunda, I. Vomitoria

## WARRANTY:

The manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled uses when applied according to label directions. Because of widely varying use conditions, it is impossible to eliminate all risks even when label directions are followed. Therefore, the manufacturer makes no other implied or express warranty nor is any agent of the manufacturer allowed to do so. Upon purchase of this product, the buyer assumes all risks associated with use of this product. In the event of damage resulting from a breach of warranty the buyer agrees to accept a refund of the purchase price of the product as full discharge of the manufacturer's liability.

# ACME® barrier 2G landscaping herbicide

56-89

Around Trees

Edge  
Along Fences

Landscaped Areas

For  
Weed Free  
Rose Gardens

## CONTROLS WEEDS & GRASSES in ROSES, EVERGREENS and OTHER SHRUBS

### ACTIVE INGREDIENT:

*Dichlobenil	2.0%
INERT INGREDIENTS	98.0%

\* 2,6-dichlorobenzonitrile.

TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN  
CAUTION

SEE OTHER PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET WEIGHT 1.25 POUNDS

STOP. READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Do not breathe dust. Do not allow contact with eyes or skin.

### Statement of Practical Treatment

If Swallowed: Induce vomiting by drinking water and touching back of throat with finger.

If Inhaled: Remove to fresh air.

If In Eyes: Rinse with water for 15 minutes.

If On Skin: Wash with soap and water.

### Environmental Hazards

Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. "Cover, collect, or incorporate granules spilled on the soil surface. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate water by cleaning of equipment or disposal of wastes."

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**STORAGE:** Store in original container in a locked, cool storage area. Keep container cap tightly closed. Do not store near seeds, bulbs, or in greenhouses. Do not contaminate water, food or feed by storage or disposal. **PRODUCT DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Do not reuse container. Rinse thoroughly before discarding in trash.

## BEFORE APPLICATION:

- Remove existing weeds.
- Thoroughly work and smooth the area to be treated.
- Carefully measure area to be treated and determine the number of square feet.

NOTE: If area to be treated is large, it should be divided into smaller sections, and each treated individually.

## APPLICATION AND RATE:

Shake granules uniformly on the soil at the rate of ¼ ounce (3 level tablespoons) for each 10 square feet to be treated. The 20 ounce shaker package treats 270 square feet, the 5 pound bag treats 1,080 square feet. As a guide for proper application and uniformity of distribution of granules, consult the following diagram.

- Rates higher than those indicated may cause injury.
- More accurate and uniform application can be obtained by applying one-half of the required material over the entire treatment area and repeating the application with the remaining material.
- This product should not be used on sloping areas, as run-off may result in injury to lawn or other plants in adjacent areas.

## AFTER APPLICATION:

- Unless the soil is wet, sprinkle lightly to moisten the soil ¼ to ½ inch deep. Do not flood, avoid puddling and washing.
- Avoid disturbing soil in the treated area as long as weed control is being maintained.

503/983  
EPA REG. NO. 2217-678-33955  
EPA EST. NO. 2217-KS-1

DISTRIBUTED BY ACME DIVISION  
**G pbi / Gordon**  
**CORPORATION**  
KANSAS CITY, KANSAS 64118



2217-678

06/22/1987

1/3

# FRUIT AND NUTS IN ORCHARDS AND NURSERIES

This product may be used on the following crops: do not allow livestock to graze treated orchards:

- A. Almond, Apple, Blueberry, Cherry, Grape, Mango, Nectarine, Peach, Pear, Prune Plum, Filbert, Pecan (Do not use on pecan seedlings within 6 months of planting), and English Walnut. May be used on bearing, non-bearing and nursery stock. Do not apply until 4 weeks after transplanting. Do not make application within one month of harvest on nuts or stone fruits. Apply 200 to 300 pounds per acre for annual weed control based on area actually treated. For control of perennial weeds apply 300 pounds per acre. Follow General Directions for Use.
- B. Blackberry and Raspberry: For control of annual weeds follow General Directions for Use, except use only on established plantings and do not apply more than 200 pounds per acre. Do not apply during new shoot emergence.
- C. Citrus (bearing and non-bearing oranges, grapefruit, tangerines, lemons, and limes) in groves: Apply 150 to 300 pounds per acre and incorporate immediately to a depth of 1 to 2 inches by use of power driven rotary hoe. Do not apply less than one year after transplanting, do not apply before water ring is removed.

On citrus nurseries this product may be used for control of annual weeds in the following citrus root stocks: Cleopatra mandarin, rough lemon, sour orange, and sweet orange. Apply to weed-free soil at 200 to 300 pounds per acre. Make first application in early spring after removal of banks, and second in early summer, not less than 90 days after the first application. Apply broadcast and irrigate lightly with overhead system (1/4 to 1/2 inch).

Do not apply during 6 months following possible freeze damage on any citrus.

## RATE CONVERSION CHART

BARRIER 2G LBS. PER ACRE =====	BARRIER 2G LBS. PER 1.000 SQFT =====
150 .....	7.00
200 .....	9.32
250 .....	11.00
300 .....	14.00

**ACCEPTED**  
**JUN 22 1987**  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 3217-678

2000

0000

2000

0000