



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY**

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals):

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes or skin with plenty of water. If irritation develops get medical attention. Avoid contamination of food or feedstuffs.

Environmental Hazards: Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate domestic or irrigation waters.

DIRECTIONS FOR USE:

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children. Do not store near food or feedstuffs. To prevent cross contamination, do not store near other fertilizers, insecticides, fungicides or seeds.

PRODUCT DISPOSAL: Securely wrap original container with unused product in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Rinse thoroughly before discarding in trash.

OTHER USE PRECAUTIONS

Do not allow spray to drift to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result. Do not spray where grass is emerging from dormancy. Do not apply to newly seeded lawn grasses until well established. Do not spray when the temperature exceeds 85°F. Seed can be safely sown 3-4 weeks after application. Do not use on carpetgrass, dichondra, St. Augustine, Centipede, closely mowed bentgrass, nor on lawns or turf where desirable clovers are present. Do not spray roots of trees and ornamentals. Do not exceed specified dosages for any area, be particularly careful within dripline of tree and other ornamental species.

TRIMEC is a registered trademark of PBI Gordon Corporation. U.S. Patent No. 3,284,186.
© 1982 PBI Gordon Corporation.

ACME®
liquid
weed & feed
15-0-0

78-16

READY TO USE

TRIMEC®
COMPLEX
PROFESSIONAL QUALITY

Controls Dandelion, Chickweed, Knotweed, Henbit, Plantains, Clover, Spurge and many other broadleaf lawn weeds; some of which are shown on the outside box.

ONE QUART COVERS UP TO 5,000 SQ. FT.

ACTIVE INGREDIENTS

**Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid	5.670%
**Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy)propionic acid	2.736%
**Dimethylamine Salt of Dicamba (3,6-dichloro-anisic acid)	0.613%
INERT INGREDIENTS	90.981%
TOTAL	100.000%

This Product Contains:
 *0.44 lb. 2,4-Dichlorophenoxyacetic acid per gallon or 4.711%
 *0.211 lb. 2-(2-methyl-4-chlorophenoxy)propionic acid per gallon or 2.261%
 *0.348 lb. Dicamba (3,6-dichloro-anisic acid) per gallon or 3.614%

KEEP OUT OF REACH OF CHILDREN

CAUTION

Statement of Practical Treatment
 In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention.
 See Side Panels for additional precautionary statements

NET CONTENTS ONE QUART (NET WEIGHT: 2 LBS 5 OZ.)

ACCEPTED

SEP 19 1986

FBI Gordon Corp.

GUARANTEED ANALYSIS

Total Nitrogen (N), Not Less Than 15.000%
 15% Urea Nitrogen

**No Spreading — No Complicated Mixing
 Just attach to a garden hose & spray**

WHERE TO USE:

LIQUID WEED & FEED may be used on Bahia, Bermuda, Bluegrass, Fescue, Rye or Zoysia lawns, vacant lots, playgrounds and other similar areas where broad leaf weeds are present. This product is especially effective on all hard to kill weeds such as Poison Ivy, Poison Oak, Sheep Sorrel and Spurge. In addition, **LIQUID WEED & FEED** controls most other broadleaf weeds including sensitive weeds such as Chickweed, Clover, Dandelion, Knotweed and Plantain.

WHEN TO USE:

1. Spray at any time during the growing season when weeds are actively growing.
2. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. Wait 3 weeks before putting new grass seed or stolons down in the bare spots.
3. For weed control, spray is applied during drought or just before rain.
4. For best results, mow 1 to 2 days before treating.
5. Do not water lawn within 24 hours after treatment.

**** HOW TO USE:**

SHAKE WELL BEFORE USING


1. Connect spray nozzle to Garden hose, remove safety cap from container, attach tube to applicator and in soft tube section. Attach container to end of nozzle.
2. Turn on water, place finger over hole on right side of nozzle and proper ratio of **LIQUID WEED & FEED** will automatically be dispersed. Spray directly over area requiring application evenly and uniformly. Check handy measuring guide to see how much of this liquid has been used.

803 RB5
 EPA Reg. No. 2217-681-33955
 EPA Est. 9521-C-4-2




MEASURING GUIDE

1,000 sq. ft. coverage
2,000 sq. ft. coverage
3,000 sq. ft. coverage
4,000 sq. ft. coverage
5,000 sq. ft. coverage when applied



1/3

One-half gallon will cover up to 10,000 square feet.



HOSE END SPRAYER INCLUDED!
COVERS UP TO 5,000 SQ. FT.
KILLS WEEDS & FEEDS GRASS
ONE-STEP APPLICATION

weed & feed
liquid



ACME



READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals):

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. In case of contact immediately flush eyes or skin with plenty of water. If irritation develops get medical attention. Avoid contamination of food or feeds with. Environmental Hazards: Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate domestic or irrigation waters.

DIRECTIONS FOR USE:

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children. Do not store near food or feeds/feeds. To prevent cross-contamination do not store near other fertilizers, insecticides, fungicides, or seeds.

PRODUCT DISPOSAL: Securely wrap original container with unused product in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Rinse thoroughly before discarding in trash.

OTHER USE PRECAUTIONS:

Do not allow spray to drift to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result. Do not spray where grass is emerging from dormancy. Do not apply to newly seeded lawn grasses until well established. Do not spray when the temperature exceeds 85°F. Seed can be safely sown 3-4 weeks after application. Do not contaminate domestic or irrigation waters. Do not use on carpetgrass, dichondra, St. Augustine, Bermuda, closely-mowed bentgrass, nor on lawns or turf where desirable clovers are present. Do not spray roots of trees and ornamentals. Do not exceed specified dosages for any area, be particularly careful within dripline of tree and other ornamental spaces.

Limited Warranty: Manufacturer warrants that the chemical ingredients contained in the ingredients statement shown on the label and that this product is used for the intended use when applied according to label directions. Because of safety carrying use control laws, it is impossible to estimate all uses from which label directions are followed.

Exclusion of Other Warranties and Remedies: Except where such exclusions and limitations are specifically provided by applicable law, THE FOREGOING IS THE ONLY WARRANTY OR REMEDY APPLICABLE TO THIS PRODUCT AND IS GIVEN EXPRESSLY AND IN LIEU OF ALL OTHER WARRANTIES EXPRESS OR IMPLIED OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, AND ALL SUCH WARRANTIES WHICH EXCEED OR DIFFER FROM SAID LIMITED WARRANTY ARE DISCLAIMED BY MANUFACTURER and their manufacturers companies with and without warranty. BUYER SHALL HAVE NO REMEDY AGAINST MANUFACTURER FOR ANY TYPE OF DAMAGE OR LOSS, AND IN NO EVENT SHALL MANUFACTURER BE LIABLE FOR CONSEQUENTIAL DAMAGE OR LOSS.

TRACOR is a Registered Trademark of PPG-INDUSTRIES CORPORATION
U.S. Patent No. 3,276,888

- NO SPREADING
 - NO COMPLICATED MIXING
- JUST ATTACH TO A GARDEN HOSE AND SPRAY!!**

WHERE TO USE:

LIQUID WEED & FEED may be used on Bahia Bermuda, Bluegrass, Fescue, Rye or Zoysia lawns, vacant lots, playgrounds and other similar areas where broadleaf weeds are present. This product is especially effective on all hard-to-kill weeds such as Poison Ivy, Poison Oak, Sheep Sorrel and Spurge. In addition LIQUID WEED & FEED controls most other broadleaf weeds, including sensitive weeds such as Chickweed, Clover, Dandelion, Knotweed and Plantain.

WHEN TO USE:

1. Spray at any time during the growing season when weeds are actively growing.
2. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. Wait 3 weeks before putting new grass seed or stolons down in the bare spots.
3. Poor weed control may result if spray is applied during drought or just before rain.
4. For best results, mow 1 to 2 days before treating.
5. Do not water lawn within 24 hours after treatment.

HOW TO USE

JUST ATTACH TO A GARDEN HOSE & SPRAY




INSTRUCTIONS

1. Connect spray nozzle (see diagram) to garden hose.
2. Shake contents of bottle well.
3. Remove safety cap.
4. Attach tube to nozzle applicator.
5. Attach bottle to open end of spray nozzle. Turn bottle counter-clockwise until threads mesh. Then turn the bottle clockwise until slightly closed.
6. Turn on water, place finger over hole on right side of nozzle and apply rate of LIQUID WEED & FEED until automaticity is observed. Spray directly over area requiring application evenly and uniformly. (Check handy measuring guide to see how much of the liquid has been used.)



WEEDS CONTROLLED


Bedstraw	Ground Ivy	Prosperrilla	Spurge
Black Medic	Hand-Ail	Ragweed	Spurge
Buckhorn	Horsetail	Plantain	Thistle
Burdock	Knotweed	Poison Ivy	Weed Carrot
Chickweed	Lambquarters	Poison Oak	Weed Garlic
Clover	Lynxseed	Purplestem	Weed Larkspur
Dandelion	Mallow	Ragweed	Weed Onion
Dock	Morning Glory	Sheep Sorrel	Yarrow
	Oxalis	Shepherd's Purse	



COVERS UP TO 5,000 SQ. FT.



COVERS UP TO 5,000 SQ. FT.



COVERS UP TO 5,000 SQ. FT.




Complete with
**HOSE-END
SPRAYER**

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid	5.87%
Dimethylamine Salt of 2,4,4'-trichlorophenoxyacetic acid	2.73%
Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)	0.19%
INERT INGREDIENTS:	90.87%
TOTAL:	100.00%

This Product Contains:
 * 9.44 lb. 2,4-Dichlorophenoxyacetic acid per gallon or 4.71%
 * 9.21 lb. 2,4,4'-trichlorophenoxyacetic acid per gallon or 4.28%
 * 0.30 lb. Dicamba (3,6-dichloro-o-anisic acid) per gallon or 0.14%

KEEP OUT OF REACH OF CHILDREN
CAUTION
 Statement of Symbolic Treatment
 In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention.
 See Side Panels for additional precautionary statements.



PPG-INDUSTRIES CORPORATION
 P.O. BOX 1000
 PITTSBURGH, PA. 15201