



~~***~~ OPTION A OR B TO BE USED DEPENDING ON THE SUBREGISTRATION

OPTION (A)

" A lawn will typically utilize 1 to 4 pounds of nitrogen per 1,000 square feet each growing season. When applied as directed this product supplies 0.069 pounds of nitrogen per 1,000 square feet with each application. Use this product in conjunction with an additional fertilization program to provide at least 1 pound of nitrogen per 1,000 square feet each growing season. Consult your local agricultural extension agent for the proper amount of nitrogen to be applied in your area."

OPTION (B)

"Trimec Liquid Weed & Feed should be supplemented by Liquid Lawn Fertilizer every 3 weeks from March through September (or April through October depending upon the growing season in your area). Please consult your local extension office for questions regarding frequency of this product in your area."



MEASURING GUIDE

1,000 sq. ft. coverage

3,000 sq. ft. coverage

5,000 sq. ft. coverage when empty

STOP. READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals)

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes or skin with plenty of water. If irritation develops get medical attention. Avoid contamination of food or feed stuffs.

Environmental Hazards: Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate domestic or irrigation waters.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children. Do not store near food or feedstuffs. To prevent cross contamination, do not store near other fertilizers, insecticides, fungicides, or seeds.

PRODUCT DISPOSAL: Securely wrap original container with unused product in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Rinse thoroughly before discarding in trash.

OTHER USE PRECAUTIONS:

Do not allow spray to drift to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result. Do not spray where grass is emerging from dormancy. Do not apply to newly seeded lawn grasses until well established. Do not spray when the temperature exceeds 85°F. Seed can be safely sown 3-4 weeks after application. Do not use on carpetgrass, dichondra, St. Augustine, Centipede, citrusly mowed bentgrass, nor on lawns or turf where desirable clovers are present. Do not spray roots of trees and ornamentals. Do not spray over bird droppings. In any area, be particularly careful within dripline of tree and other ornamental species.

© 1982, PBI-GORDON CORP.

**TRIMEC®
LIQUID WEED & FEED**

15-0-0

NET CONTENTS ONE QUART
(NET WEIGHT: 2 LBS 5 OZ)

Contains TRIMEC® Turf Herbicide

-- KEEP FROM FREEZING --

1 QUART COVERS UP TO 5,000 SQ. FT.

KILLS: Dandelion, Chickweed, Knotweed, Henbit, Plantains, Clover, Spurge and many other broadleaf lawn weeds, some of which are shown on the outside of box.

The fast, easy way to weed control and fertilize your lawn.

No Spreading — No Complicated Mixing
Just attach to a garden hose & spray

ACTIVE INGREDIENTS:

***Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid	5.51%
**Dimethylamine Salt of 2,4-dimethyl-4-chlorophenoxy propionic acid	2.19%
***Dimethylamine Salt of Dicamba (3,6-dichloro-p-toleric acid)	0.87%
INERT INGREDIENTS	90.97%
This Product Contains	TOTAL 100.00%

10-44-0: 2,4-Dichlorophenoxyacetic acid per gallon of 4.11%
10-211-0: 2,4-dimethyl-4-chlorophenoxy propionic acid per gallon of 2.26%
10-048-0: Dicamba (3,6-dichloro-p-toleric acid) per gallon of 0.54%

KEEP OUT OF REACH OF CHILDREN

-- CAUTION --

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water, for eyes, flush with water for 15 minutes and get medical attention.

See back panel for additional precautionary statements.

TRIMEC® is a registered trademark of PBI-Gordon Corporation, U.S. Patent No. 3,284,186

Mfg. by **G** **pbi/gordon**
CORPORATION
KANAS CITY, KANSAS 66108

Total Nitrogen (N) N
Nitrogen from Ur



EPA REG. NO.
EPA EST. NO.

TRIMEC® LIQUID WEED & FEED Fertilize Rye or Zoysia lawns, where broad leaf weeds are present. Hard to kill weeds such as Poa. In addition TRIMEC® LIQUID WEED & FEED kills Knotweed and Plantain.

WHEN TO USE:

1. Spray at any time during actively growing.
2. On new lawns, wait 4 inches before making before putting new grass.
3. For best weed control mow or just before rain.
4. For best results, mow.
5. Do not water lawn with



HOW TO USE:

SHAKE WELL BEFORE USE

- * Connect spray nozzle to attach hose to applicator nozzle.
- ** Turn on water, place line at end of TRIMEC® LIQUID WEED & FEED. Spray directly over area. Check handy measuring used.

NOTICE: See warnings that the following uses of use, such as water application equipment will vary.



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY**

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals)

CAUTION Harmful if swallowed. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes or skin with plenty of water. If irritation develops, get medical attention. Avoid contamination of food or feedstuffs.

Environmental Hazards: Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate domestic or irrigation waters.

DIRECTIONS FOR USE:

This is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children. Do not store near food or feedstuffs. To prevent cross contamination, do not store near other fertilizers, insecticides, fungicides, or seeds.

PRODUCT DISPOSAL: Securely wrap original container with unused product in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Rinse thoroughly before discarding in trash.

OTHER USE PRECAUTIONS:

Do not allow spray to drift to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result. Do not spray where grass is emerging from dormancy. Do not apply to newly seeded lawn grasses until well established. Do not spray when the temperature exceeds 85 F. Seed can be safely sown 3-4 weeks after application. Do not use on carpetgrass, dichondra, St. Augustine. Do not use on closely mowed turfgrass, nor on lawns or turf where desirable clovers are present. Do not spray roots of trees and shrubs. Excessive application of sprays may cause injury to particularly sensitive plants, shrubs and other nearby plants, trees.

TRIMEC is a registered trademark of G. P. B. Gordon Corporation. U.S. Patent No. 3,284,196.



Controls Dandelion, Chickweed, Knotweed, Henbit, Plantains, Clover, Spurge and many other broadleaf lawn weeds; some of which are shown on the outside box. ONE QUART COVERS UP TO 5,000 SQ. FT.

ACTIVE INGREDIENTS
 *Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid 5.670%
 **Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy)propionic acid 2.736%
 ***Dimethylamine Salt of Dicamba (3,6-dichloroaniline acid) 6.619%
INERT INGREDIENTS 30.955%
TOTAL 100.000%
 This Product Contains:
 10.44 lb. 2,4-Dichlorophenoxy, etc. and per gallon or 4.711%
 110.211 lb. 2-(2-methyl-4-chlorophenoxy)propionic acid per gallon or 2.261%
 110.048 lb. Dicamba (3,6-dichloroaniline acid) per gallon or 0.514%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**
 Statement of Practical Treatment
 In case of contact, wash skin with plenty of soap and water. For eyes, flush with water for 15 minutes and get medical attention.
 See Side Panels for additional precautionary statements.

NET CONTENTS ONE QUART (NET WEIGHT: 2 LBS 5 OZ.)

BEST AVAILABLE COPY

GUARANTEED ANALYSIS

Total Nitrogen (N) Not Less Than 15.00%
 15% Total Nitrogen

No Spreading — No Complicated Mixing
 Just attach to a garden hose & spray

WHERE TO USE


LIQUID WEED & FEED may be used on Bahia, Bermuda, Bluegrass, Fescue, Ryegrass, and other cool season grasses. It is also effective on broadleaf weeds and grasses. It is applied as a pre-emergent or a post-emergent. It is applied as a pre-emergent on lawns, and as a post-emergent on lawns, golf courses, and other areas where broadleaf weeds are present. It is applied as a pre-emergent on lawns, and as a post-emergent on lawns, golf courses, and other areas where broadleaf weeds are present. It is applied as a pre-emergent on lawns, and as a post-emergent on lawns, golf courses, and other areas where broadleaf weeds are present.

WHEN TO USE

1. Spray at any time during the growing season when weeds are beginning to grow.
2. On new lawns, wait until the grass has reached a height of 2 inches before making any application. After treatment, wait 3 weeks before putting new grass seed in spots down in the bare spots.
3. For weed control, re-spray as needed for maintenance.
4. For best results, mow 1 to 2 days before treating.
5. Do not water lawn within 24 hours after treatment.

HOW TO USE

SHAKE WELL BEFORE USING
 Connect spray nozzle to garden hose, remove safety cap from container, attach to the applicator and invert the section. Attach container to end of nozzle. Turn on water, place trigger over bare or spotted area of lawn, and spray. LIQUID WEED & FEED will kill weeds and grasses. Spray directly over the weeds and grasses. Do not spray on desirable plants. Do not spray on lawns or turf where desirable clovers are present. Do not spray roots of trees and shrubs. Excessive application of sprays may cause injury to particularly sensitive plants, shrubs and other nearby plants, trees.



5,000 sq. ft. coverage

MEASURING GUIDE

100 sq. ft. coverage

1000 sq. ft. coverage


1000 sq. ft. coverage

1000 sq. ft. coverage

MU

3. Remove both parts of separated lock ring.
4. Turn arrow on activator switch to OFF position.
5. Push activator switch securely down against top of sprayer. This will pierce the safety seal inside the dispenser.
6. To begin spraying, turn on water and then move activator switch to the ON position.
7. To stop dispensing the product, turn off water.

✓




(Interstate Nozzle Directions)

★ HOW TO USE:

SHAKE WELL BEFORE USING.

1. Attach spray nozzle to garden hose. Turn nozzle clockwise to firmly attach it to hose.
2. Remove cap from bottle (directions on cap) and screw bottle into spray nozzle.
3. Turn the on/off switch on top of the spray nozzle counter clockwise to the "OFF" position. Then turn on the water at the faucet.
4. To begin spraying, turn the sprayer nozzle switch clockwise to the "ON" position. The water automatically mixes with the product. Walk at a steady pace spraying in a side to side arc, slightly overlapping treated areas. Refer to graduated scale on side of container to guide coverage.
5. To stop spraying, turn the sprayer nozzle switch counter clockwise to the "OFF" position and turn off water at the faucet.




**

HOW TO USE:

This half-gallon package is intended as a refill for the quart hose-end sprayer applicator/container of Acme Liquid Weed & Feed 15-0-0. However, this product may be used in any hose-end sprayer in which herbicides have been used.

Add Acme Liquid Weed & Feed 15-0-0 to the hose-end sprayer jar. Do not add water to the jar. Shake jar well. One-half gallon will fill your quart hose-end sprayer jar twice. One quart covers 5,000 square feet. One-half gallon will cover up to 10,000 square feet.

✓



HOSE END SPRAYER INCLUDED!
COVERS UP TO 5,000 SQ. FT.
KILLS WEEDS & FEEDS GRASS
ONE-STEP APPLICATION

ACME[®] Liquid Weed & Feed



READ THE ENTIRE LABEL FIRST
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals):
CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes or skin with plenty of water. If irritation develops, get medical attention. Avoid contamination of food or feedstuffs.
Environmental Hazards: Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate water by leaking of equipment or disposal of wastes. Do not contaminate domestic or irrigation waters.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children. Do not store near food or feedstuffs. To prevent cross-contamination, do not store near other fertilizers, insecticides, fungicides or seeds.

PRODUCT DISPOSAL: Securely wrap original container with unused product in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Rinse thoroughly before discarding in trash.


OTHER USE PRECAUTIONS:

Do not allow spray to drift to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result. Do not spray where grass is emerging from dormancy. Do not apply to newly seeded lawn grasses until well established. Do not spray when the temperature exceeds 85°F. Seed can be safely sown 3-4 weeks after application. Do not contaminate domestic or irrigation waters. Do not use on carpetgrass, dichondra, St. Augustine, Centipede, closely-mowed bermgrass, nor on lawns or turf where desirable clovers are present. Do not spray roots of trees and ornamentals. Do not exceed specified dosages for any area, be particularly careful within drooping of tree and other ornamental species.

United Technology Manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the intended use when applied according to label directions. Because of water varying use conditions, it is impossible to guarantee all results over other label structures are followed.

Exclusion of Other Warranties and Remedies. Except where such warranties and remedies are specifically provided by separate law, THE FOREGOING IS THE ONLY WARRANTY OR REMEDY APPLICABLE TO THIS PRODUCT AND IS GIVEN EXPRESSLY AND IN LIEU OF ALL OTHER WARRANTIES EXPRESS OR IMPLIED, OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, AND ALL SUCH WARRANTIES WHICH EXCEED OR DIFFER FROM SAID LIMITED WARRANTY ARE DISCLAIMED BY MANUFACTURER, and upon Manufacturer's compliance with said limited warranty, BUYER SHALL HAVE NO REMEDY AGAINST MANUFACTURER FOR ANY TYPE OF DAMAGE OR LOSS and IN NO EVENT SHALL MANUFACTURER BE LIABLE FOR CONSEQUENTIAL DAMAGE OR LOSS.

Trade-C is a Registered Trademark of PBI GORDON CORPORATION
U.S. Patent No. 3,264,488



COVERS UP TO 5,000 SQ. FT.

BEST AVAILABLE COPY

- NO SPREADING
 - NO COMPLICATED MIXING
- JUST ATTACH TO A GARDEN HOSE AND SPRAY!!**

WHERE TO USE:

LIQUID WEED & FEED may be used on Bahia, Bermuda, Bluegrass, Fescue, Rye or Zoysia lawns, vacant lots, playgrounds and other similar areas where broadleaf weeds are present. This product is especially effective on all hard to kill weeds such as Poison Ivy, Poison Oak, Sheep Sorrel and Spurge. In addition, LIQUID WEED & FEED controls most other broadleaf weeds, including sensitive weeds such as Chickweed, Clover, Dandelion, Knotweed and Plantain.

WHEN TO USE:

1. Spray at any time during the growing season when weeds are actively growing.
2. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. Wait 3 weeks before putting new grass seed or sods down in the bare spots.
3. Poor weed control may result if spray is applied during drought or just before rain.
4. For best results, mow 1 to 2 days before treating.
5. Do not water lawn within 24 hours after treatment.

HOW TO USE

JUST ATTACH TO A GARDEN HOSE & SPRAY



INSTRUCTIONS

1. Connect spray nozzle (see diagram) to garden hose.
2. Shake contents of bottle well.
3. Remove safety cap.
4. Attach tube to nozzle applicator.
5. Attach bottle to open end of spray nozzle. Turn bottle counter-clockwise until threads mesh. Then turn the bottle clockwise until tightly closed.
6. Turn on water, place finger over hole on right side of nozzle and proper rate of LIQUID WEED & FEED will automatically be dispersed. Spray directly over area requiring application evenly and uniformly. (Check handy measuring guide to see how much of this liquid has been used.)



WEEDS CONTROLLED

Betulae	Ground Ivy	Prostratae	Sarcocolla
Black Alder	Heart-As	Spurred	Scourge
Buckhorn	Hemlock	Plantain	Thrush
Burdock	Ironweed	Shower Ivy	Wild Control
Chicory	Lambquarters	Trifolium	Wild Garlic
C. reticulata	Lupinus	Purple	Wild Onion
Clover	Mallow	Sheep Sorrel	Yarrow
Dandelion	Morning Glory	Shepherd's Purse	
Oxalis	Oxalis		



COVERS UP TO 5,000 SQ. FT.




Complete with
**HOSE-END
SPRAYER**



ACTIVE INGREDIENTS:
 *Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid 5.670%
 **Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid 2.736%
 ***Dimethylamine Salt of Dicamba (3,6-dichloro-p-aminic acid) 0.619%
INERT INGREDIENTS: 90.975%
TOTAL 100.000%

This Product Contains:
 * 0.44 lb. 2,4-Dichlorophenoxyacetic acid per gallon of 4.71%
 ** 0.27 lb. 2-(2-methyl-4-chlorophenoxy) propionic acid per gallon of 2.73%
 *** 0.06 lb. Dicamba (3,6-dichloro-p-aminic acid) per gallon of 0.62%

KEEP OUT OF REACH OF CHILDREN!
CAUTION
 Statement of Practical Treatment
 In case of contact, wash skin with plenty of clean water; for eyes, flush with water for 15 minutes and get medical attention.
 See Side Panel for full list of occupational statements.



COVERS UP TO 5,000 SQ. FT.