

2217-681

12/12/2013

1/13



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
Washington, D.C. 20460

OFFICE OF  
CHEMICAL SAFETY AND  
POLLUTION PREVENTION

DEC 12 2013

James L. Kunstman, Ph.D.  
PBI/Gordon Corporation  
Post Office Box 014090  
Kansas City, Missouri 64101

**Subject:** Amendment adding a sub-label, advertising claims and updating the Storage and Disposal and Environmental Hazard Statement.  
**Product name:** Trimec® Liquid Weed and Feed  
**EPA Reg. No:** 2217-681  
**Application Dated:** May 2, 2013


Dear Mr. Kunstman,

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

A stamped copy of your label is enclosed for your records. This label supersedes all previously accepted labels. You must submit one (1) copy of the final printed label before you release the product for shipment. Products shipped after eighteen (18) months from the date of this letter must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA §6(e). Your release for shipment of the product constitutes acceptance of these conditions.

If you have any questions, please contact Grant Rowland at 703-347-0254 or at Rowland.Grant@epa.gov.

Sincerely,

  
Kathryn Montague  
Product Manager 23  
Herbicide Branch  
Registration Division (7505P)


2/13

# TRIMEC® LIQUID WEED & FEED

## EPA Reg. No. 2217-681

Two sublabels represent the entire or master label of TRIMEC® LIQUID WEED & FEED. Please refer to the appropriate section of the labeling as shown as follows:

<b>Section</b>
Label language common to all sublabels
Sublabel 1. Draft labeling previously approved on November 4, 2008
Sublabel 2. Proposed instructions for small trailer sprayers
Appendix
Document Control Information



3/13

**Label Language Common To All Sublabels**

**TRIMEC® LIQUID WEED & FEED**

**EPA Reg. No. 2217-681**

**Alternate Brand Names:**

**ACTIVE INGREDIENT:**

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid .....	5.67%
Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid .....	1.37%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid) .....	0.63%

<b>INERT INGREDIENTS:</b> .....	<b>92.33%</b>
<b>TOTAL</b> .....	<b>100.00%</b>


**THIS PRODUCT CONTAINS:**

- 0.44 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 4.71%
- 0.11 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 1.13%
- 0.05 lb Dicamba (3,6-dichloro-o-anisic acid equivalent per gallon or 0.52%

Isomer Specific by AOAC Methods.

Note: If the design, format, or small size of the container labels makes it impractical to present the entire ingredient statement on the front panel, then add a referral statement and present this substatement on the back panel: "See back panel for substatement of ingredient statement."

GUARANTEED ANALYSIS	
Total Nitrogen (N), Not less than .....	15%
15% Urea Nitrogen - Derived from Urea	



**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals**

**CAUTION:** Harmful if swallowed. Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

<b>First Aid</b>	
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call toll-free, 1-877-800-5556 for emergency medical information.</p>	

**Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

**[Environmental Hazards (alternate per PRN 2008-1)**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.]

5/13

## **DIRECTIONS FOR USE**

### **STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Keep from freezing. Store in original container in a locked storage area inaccessible to children.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or [www.website.com](http://www.website.com) for disposal instructions. Never place unused product down any indoor or outdoor drain.

### **LIMITED WARRANTY AND DISCLAIMER**

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

6/13

## SUBLABEL 1

**Draft labeling previously approved on November 4, 2008**

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### **PRECAUTIONS AND RESTRICTIONS:**

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Do not apply as a fine mist because of potential for injury to desirable plants. Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.
- Do not spray when grass is emerging from dormancy. Do not apply to newly seeded lawn grasses until well established. Do not spray when air temperatures exceed 85°F. Seed can be sown 3 to 4 weeks after application.
- Do not use on carpetgrass, dichondra, St. Augustinegrass, centipedegrass, nor on lawns or turf where desirable clovers are present.
- Do not exceed specified dosages for any area.

#### **State fertilizer requirements:**

##### **OPTION (A)**

A lawn will typically utilize 1 to 4 pounds of nitrogen per 1,000 square feet each growing season. When applied as directed this product supplies 0.069 pounds of nitrogen per 1,000 square feet with each application. Use this product in conjunction with an additional fertilization program to provide at least 1 pound of nitrogen per 1,000 square feet each growing season. Consult your local Agricultural Extension agent for the proper amount of nitrogen to be applied in your area.

##### **OPTION (B)**

Trimec® Liquid Weed & Feed should be supplemented by Liquid Lawn Fertilizer every 3 weeks from March through September (or April through October depending upon the growing season in your area) Please consult your local extension office for questions concerning a regular fertilizer program.

#### **WHERE TO USE:**

Trimec® Liquid Weed & Feed may be used on bahiagrass, bermudagrass, bluegrass, fescue, perennial ryegrass or zoysiagrass lawns, vacant lots, playgrounds and other similar areas where broadleaf weeds are present. This product is especially effective on hard-to-kill weeds such as poison ivy, poison oak, sheep sorrel and spurge. In addition Trimec® Liquid Weed & Feed controls listed broadleaf weeds, including sensitive weeds such as chickweed, clover, dandelion, knotweed and plantain.

#### **WHEN TO USE:**

1. Spray any time during the growing season when weeds are actively growing.
2. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. Wait 3 weeks after application before seeding or planting stolons.
3. Poor weed control may result if spray is applied during drought or just before rain.

- 4. For best results, mow 1 to 2 days before treating.
- 5. Do not water lawn within 24 hours after treatment.

**Limitations on broadcast applications to residential turfgrass:**

The maximum application rate is 32 fl.oz. [1 quart] of product per 5,000 sq.ft. per application [0.96 lb 2,4-D ae, 0.24 lb MCP-P ae and 0.11 lb dicamba ae per acre per application]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 64.0 fl.oz. [2 quarts] of product per 5,000 sq.ft. [1.92 lb 2,4-D ae, 0.48 lb MCP-P ae and 0.22 lb dicamba ae per acre].

**HOW TO USE:** Shake well before using.

- One quart covers 5,000 square feet.
- One-half gallon covers up to 10,000 square feet.

**OPTIONS (C) THRU (K) - ONE TO BE USED DEPENDING ON TYPE OF READY-SPRAY® HOSE-END SPRAYER**

**OPTION (C)**  
Connect spray nozzle to garden hose; remove safety cap from container; attach tube to applicator and insert tube section. Attach container to end of nozzle. Turn on water; place finger over hole on right side of nozzle and proper ratio of Trimec® Liquid Weed & Feed will automatically be dispersed. Spray directly over area requiring application evenly and uniformly.

**OPTION (D)**  
This half-gallon package is intended as a refill. One-half gallon will fill your quart hose end sprayer twice. One quart covers 5,000 square feet. One-half gallon will cover up to 10,000 square feet.

**OPTION (E)**

1. Attach spray nozzle to garden hose. Turn nozzle clockwise to firmly attach it to hose.
2. Remove cap from bottle (directions on cap) and screw bottle into spray nozzle.
3. Turn the on/off switch on top of the spray nozzle counter clockwise to the "OFF" position. Then turn on the water at the faucet.
4. To begin spraying, turn the sprayer nozzle switch clockwise to the "ON" position. The water automatically mixes with the product. Walk at a steady pace spraying in a side to side arc, slightly overlapping treated areas. Refer to graduated scale on side of container to guide coverage.
5. To stop spraying, turn the sprayer nozzle switch counter clockwise to the "OFF" position and turn off water at the faucet.

**OPTION (F)**

1. Attach garden hose to connector on dispenser
2. Insert screwdriver into slot in orange lock ring and twist, snapping lock ring in two.
3. Remove both parts of separated lock ring.
4. Turn arrow on activator switch to OFF position.
5. Push activator switch securely down against top of sprayer. This will pierce the safety seal inside the dispenser.
6. To begin spraying, turn on water and then move activator switch to the ON position.
7. To stop dispensing the product, turn off water.

**OPTION (G)**

1. Connect spray nozzle to garden hose. Be sure that valve is in "OFF" position.
2. Remove bottle cap and screw bottle onto nozzle. Turn on water.
3. Turn valve to "ON" position to start flow of water. Use valve to shut off flow of water at any time.

**OPTIONS (H) THROUGH K - DEPENDING ON PACKAGING USED.**

**OPTION (H)**

Check handy measuring guide to determine remaining area to be covered.

**OPTION (I)**

Check handy measuring guide to see how much of this liquid has been used.

**OPTION (J) to be used on label for one-half gallon [2 quart] container.**

**HOW TO USE:**

1. SHAKE WELL BEFORE USING.
2. Unscrew the cap from this container and carefully remove the foil seal.
3. Fill the hose end sprayer jar with the amount of product specified in the chart below. **DO NOT** dilute this product with water when filling the sprayer jar.
4. Attach the sprayer to garden hose.
5. Turn water on at full volume.
6. Hold the sprayer at waist level with the nozzle pointed away from the face and body.
7. Walk back and forth over lawn while spraying (as you would when mowing) covering the lawn thoroughly. Overlap slightly for uniform coverage.

**OPTION (K) to be used on label for 1 quart container.**

**HOW TO USE:**

1. SHAKE WELL BEFORE USING.
2. Connect spray nozzle to garden hose.
3. Remove white cap from bottle and screw bottle into the spray nozzle.
4. Turn the black "on-off" lever to the "off" position by turning to the right. Turn water on full volume.
5. Hold the sprayer at waist level with the nozzle pointed away from the face and body.
6. To spray, turn the black "on-off" lever to the center "on" position.
7. Walk back and forth over lawn while spraying (as you would when mowing) covering the lawn thoroughly. Overlap slightly for uniform coverage.

**BROADLEAF WEEDS CONTROLLED**

Bedstraw	Ground ivy	Peppergrass	Speedwell
Black medic	Healall	Pigweed	Spurge
Buckhorn	Henbit	Plantain	Thistle
Burdock	Knotweed	Poison ivy	Wild carrot
Chicory	Lambsquarters	Poison oak	Wild garlic
Chickweed	Lespedeza	Purslane	Wild lettuce
Clover	Mallow	Ragweed	Wild onion
Dandelion	Morningglory	Sheep sorrel	Yarrow
Dock	Oxalis	Shepherdspurse	



## SUBLABEL 2

### Proposed instructions for small trailer sprayers

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### 1. Before You Use. Read & follow these directions when using.

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Use on established lawns. Will not harm lawn grasses when used as directed (Do not use on carpetgrass, centipedegrass, St. Augustinegrass, dichondra, or desirable clovers)
- Do not apply as a fine mist because of potential for injury to desirable plants. Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse sprayer over lawn area only.

#### 2. How Much To Apply

**Bahiagrass, Bermudagrass, Fescues, Kentucky Bluegrass, Perennial Ryegrass, and Zoysiagrass:**

Entire Lawn Treatments:

- Pump Style Sprayer: Mix 6.4 fl.oz. in [0.5 to] 1 gallon of water to treat 1,000 sq.ft.
- Hose-End Sprayer: Apply 6.4 fl.oz. per 1,000 sq.ft.
- [12 volt &] Tow-Behind Sprayers: Mix 6.4 fl.oz. in [0.5 to] 1 gallon of water to treat 1,000 sq.ft.

[Optional tabular presentation]

Table 1: Entire Lawn Treatment of Bahiagrass, Bermudagrass, Fescues, Kentucky Bluegrass, Perennial Ryegrass, and Zoysiagrass		
Tank Size	Concentrate to add to tank	Coverage Area
1 gal	6.4 fl.oz.	1,000 sq.ft.
5 gal	1.0 qt	5,000 sq.ft.
15 gal	3.0 qt	15,000 sq.ft.
25 gal	5.0 qt	25,000 sq.ft.

#### 3. How To Apply

##### 3.1 [12 Volt &] Tow-Behind Boom Sprayers [sprayer icon]

Ideal for treating large country and suburban lawns

1. To calibrate your application equipment consult your equipment owner's manual for speed and pressure settings required to deliver [0.5 to] 1 gallon of spray solution to treat 1,000 sq.ft.. For example, at 4 mph a sprayer might deliver 1/2 gallon per 1,000 sq.ft. at 40 psi.
2. To apply evenly, drive at a constant speed that delivers the prescribed amount of spray required for the area to be treated.

3. Mix the appropriate amount of product and water in the sprayer using the following tables as a guide.

<b>Table 2: [12 Volt &amp;] Pull Behind Sprayers: Amount of Trimec Liquid Weed &amp; Feed [product] required to prepare 15 to 25 gallons of spray solution for broadcast treatments on Bahiagrass, Bermudagrass, Fescues, Kentucky Bluegrass, Perennial Ryegrass, and Zoysiagrass:</b>			
<b>Tank Size</b>	<b>Spray volume</b>	<b>Coverage Area</b>	<b>Concentrate to add to tank</b>
15 gal	0.5 gal/1000 sq.ft.	30,000 sq.ft. (0.68 acre)	1.5 gal
	1 gal/1000 sq.ft.	15,000 sq.ft. (0.34 acre)	0.75 gal
25 gal	0.5 gal/1000 sq.ft.	50,000 sq.ft. (1.15 acre)	2.5 gal
	1 gal/1000 sq.ft.	25,000 sq.ft. (0.57 acre)	1.25 gal

### 3.2 Pump Style Sprayers [sprayer icon]

Ideal for spraying individual weeds

1. Add the appropriate amount of water and concentrate to the sprayer tank.
2. Close sprayer, shake well, and pump handle to pressurize.
3. Adjust nozzle to deliver a coarse spray pattern.
4. To broadcast over large areas: apply evenly to treated area.
5. Re-pressurize the sprayer as needed to maintain a good spray pattern.

### 3.3 Dial Style Hose-End Sprayers [sprayer icon]

Ideal for treating medium-sized lawns

1. Fill sprayer with sufficient concentrate to cover the total area to be treated. **DO NOT ADD WATER.**
2. Close sprayer and set dial to 1 oz.
3. Connect sprayer to hose
4. Extend hose to furthest point from faucet. Walking back towards the faucet, begin spraying, avoiding contact with the treated area.
5. Pour any unused product back into this container.

### 3.4 Concentrate Plus Water Hose-End Sprayers [sprayer icon]

Ideal for treating medium-sized lawns

1. Fill sprayer with sufficient concentrate to cover the total area to be treated. Pour product into sprayer jar to the "Fluid Oz." level desired (See sprayer jar for markings)
2. Add the appropriate amount of water to the "Gallons" level indicated on the side of the sprayer jar. Close sprayer, connect to garden hose, shake well.
3. Extend hose to furthest point from faucet. Walking back towards the faucet, begin spraying, avoiding contact with the treated area.

## 4. For Best Results

- Apply when weeds are young and actively growing.
- Apply 1 to 3 days after mowing.
- If the lawn is dry, water 24 hours before using this product.
- Do not water lawn within 24 hours after treatment.
- For new lawns, apply only when grass has reached a height of at least 2 inches.

- Bare spots may be seeded 3 to 4 weeks after application.
- Spray when air temperatures are between 60° and 90°F with good soil moisture.
- Avoid applications during the spring transition of the warm season grasses and during periods of extremely high temperatures.

### 5. Limitations

#### Broadcast applications to residential turfgrass:

The maximum application rate is 32 fl.oz. [1 quart] of product per 5,000 sq.ft. per application [0.96 lb 2,4-D ae, 0.24 lb MCPP-p ae and 0.11 lb dicamba ae per acre per application]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 64.0 fl.oz. [2 quarts] of product per 5,000 sq.ft. [1.92 lb 2,4-D ae, 0.48 lb MCPP-p ae and 0.22 lb dicamba ae per acre].

6. Weeds Killed			
Bedstraw	Ground ivy	Peppergrass	Speedwell
Black medic	Healall	Pigweed	Spurge
Buckhorn	Henbit	Plantain	Thistle
Burdock	Knotweed	Poison ivy	Wild carrot
Chicory	Lambsquarters	Poison oak	Wild garlic
Chickweed	Lespedeza	Purslane	Wild lettuce
Clover	Mallow	Ragweed	Wild onion
Dandelion	Morningglory	Sheep sorrel	Yarrow
Dock	Oxalis	Shepherdspurse	

[Back Panel] [Restatement of information in sections 1 and 2]

Trimec® Liquid Weed & Feed kills broadleaf weeds and greens lawns in one convenient step. Made for use with all 12 volt sprayers and concentrated to treat up to 5,000 square feet. It's the ideal choice for country and suburban lawns!

**Getting Started**

**Kills:**

Dandelion, Clover, Thistle and over [XXX] other weeds listed.

**Where To Use:**

On established lawns without harming lawn grasses when used as directed (Do not use on carpetgrass, centipedegrass, St. Augustinegrass, dichondra, or desirable clovers).

**Amount To Use:**

**Bahiagrass, Bermudagrass, Fescues, Kentucky Bluegrass, Perennial Ryegrass, and Zoysiagrass:**

Entire Lawn Treatments:

- Pump Style Sprayer: Mix 6.4 fl.oz. in [0.5 to] 1 gallon of water to treat 1,000 sq.ft.
- Hose-End Sprayer: Apply 6.4 fl.oz. per 1,000 sq.ft.
- [12 volt &] Tow-Behind Sprayers: Mix 6.4 fl.oz. in [0.5 to] 1 gallon of water to treat 1,000 sq.ft.

Mixing Instructions For a Typical Broadcast Application			
Amount of Water in Sprayer	15 Gallons	25 Gallons	50 Gallons
Amount of Product to add to sprayer	3.0 qt (0.75 gal)	5.0 qt (1.25 gal)	10.0 qt (2.5 gal)
Area to Treat	15,000 sq.ft.	25,000 sq.ft.	50,000 sq.ft.

**Coverage:**

This [size] bottle treats XXXX sq.ft. of Bahiagrass, Bermudagrass, Fescues, Kentucky Bluegrass, Perennial Ryegrass, and Zoysiagrass.

**Telephone Icon**

**Computer Icon**

For questions or comments call toll-free 1-800-821-7925 or visit our website at [www.pbigordon.com](http://www.pbigordon.com)

## **APPENDIX**

### **1. Statements which may appear on different label components depending on packaging configuration.**

- See next panel for additional Precautionary Statements and First Aid
- Net Contents: \_\_\_\_\_
- EPA Est. No. \_\_\_\_\_

### **2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:**

- Keep From Freezing
- 1 Quart Covers Up To 5,000 Square Feet or (2 Quarts Covers Up To 10,000 Square Feet.)
- Controls dandelion, chickweed, knotweed, henbit, plantain, clover, spurge and listed broadleaf lawn weeds.
- The fast, easy way to weed control and fertilize your lawn.
- No Spreading - No Complicated Mixing
- TRIMEC® is a registered trademark of PBI/Gordon Corporation.
- A lawn will typically utilize 1 to 4 pounds of nitrogen per 1,000 square feet each growing season. When applied as directed this product supplies 0.069 pounds of nitrogen per 1,000 square feet with each application. Use this product in conjunction with an additional fertilization program to provide at least 1 pound of nitrogen per 1,000 square feet each growing season. Consult your local Agricultural Extension agent for the proper amount of nitrogen to be applied in your area.
- [Product name] should be supplemented by Liquid Lawn Fertilizer every 3 weeks from March through September (or April through October depending upon the growing season in your area). Please consult your local extension office for questions concerning a regular fertilizer program.
- [Just one step] Kills [Listed] Broadleaf Weeds & Quickly Greens Lawns [in One Timesaving Step!]
- All in One Timesaving Step!
- Pre-Mixed Formula Eliminates [Listed] Broadleaf Weeds While Greening Lawns
- One-Step Weed & Feed for Lawns
- [Perfect] For Use [in Lawns] with ATV or Tow-behind sprayers
- Made for 12 volt ATV & Tow-Behind Sprayers
- Won't Harm Lawns\* . \*when used as directed
- Kills Dandelion, Clover, henbit and over XXX [listed] weeds listed [in Country and Suburban Lawns]
- Quickly Greens Lawns
- Provides Quick Green-Up
- Rainfast in 8 hours
- [For 2.5 gallon container] Makes up to 50 gallons [of spray]
- [For 2.5 gallon container] Treats up to 50,000 square feet [1.15 acres]
- Makes up to XX gallons [of spray]
- Treats up to XXX square feet [xxx acres]

## **DOCUMENT CONTROL INFORMATION**

1. Unique Label Identifier: 002217-00681.20130502.amend-proposed-clean.doc

2. Reason for Issue: new sublabel, ad claims, storage and disposal