



**STOP: READ THE ENTIRE LABEL FIRST.  
OBSERVE ALL PRECAUTIONS AND  
FOLLOW DIRECTIONS CAREFULLY.**

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans & Domestic Animals**

**CAUTION:** Harmful if swallowed. Do not breathe dust.  
Do not allow contact with eyes or skin.

**Statement of Practical Treatment**

**IF SWALLOWED:** Induce vomiting by drinking water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

**IF INHALED:** Remove victim to fresh air.

**IF IN EYES:** Flush with plenty of water. Get medical attention if irritation persists.

**IF ON SKIN:** Wash with plenty soap and water.

**Environmental Hazards**

Do not contaminate water by cleaning of equipment or disposal of wastes.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE & DISPOSAL**

**STORAGE:** Store in original bag in a locked storage area inaccessible to children or pets. Storage area should be cool and dry. Keep bag tightly closed. Reseal bag by rolling down top to product level and taping or clipping. Do not store near seeds, bulbs, or in greenhouses.

**PESTICIDE DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**BEST AVAILABLE COPY**

IMPORTANT! Always wash hands, face, and arms with soap and water before smoking, eating, drinking, or toileting.

Keep all unprotected persons, children, livestock, and pets away from treated area or where there is a danger of drift.

Do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away. See Statement of Practical Treatment.

[(STATEMENT PLACED ON LABEL) "Refer to product labeling for use restrictions to protect ENDANGERED SPECIES."]


"ENDANGERED SPECIES RESTRICTIONS"

Before using this product in forests in a county listed below, you must contact the Endangered Species Specialist in the Regional/Field Office of the U.S. Fish and Wildlife Service (FWS) indicated below. You must provide FWS with your name and phone number, the product you intend to use, and the specific location in which you intend to use it. The U.S. Fish and Wildlife Service will inform you whether your proposed use is in the range of endangered species. Use of this product in the range of endangered species, as defined for you by FWS is prohibited.

Contact FWS Field Offices at the following numbers:

- ALABAMA (Jackson, Mississippi, 601-965-4900)  
Cherokee, De Kalb, Etowah, Jackson, Marshall and St. Clair
- ARKANSAS (JACKSON, MISSISSIPPI, 601-965-4900)  
Clay, Jackson, Lawrence and Woodruff
- CALIFORNIA (Sacramento, California, 916-978-4613)  
Lake
- CONNECTICUT (Concord, New Hampshire, 603-225-1411)  
Litchfield
- DELAWARE (Annapolis, Maryland, 301-269-5448)  
New Castle
- FLORIDA (Jacksonville, Florida, 904-791-2580)  
Clay, Gadsden, Gulf, Jackson, Liberty, Marion, Orange and Polk
- GEORGIA (Jacksonville, Florida, 904-791-2580)  
Baker, Brantley, Burke, Decatur, Dooly, Floyd, Gordon, Habersham, Lee, Rabun, Stephens, Sumter, Towns, Walker, Wayne, and Wheeler
- ILLINOIS (Bloomington, Indiana, 309-793-5800)  
Randolph
- IOWA (St. Paul, Minnesota, 612-725-7131)  
Allamakee, Clayton, Delaware, Dubuque, Fayette and Jackson
- MAINE (Concord, New Hampshire, 603-225-1411)  
Aroostook, Cumberland, Kennebec and York

- ✓ MARYLAND (Annapolis. Maryland. 301-269-5448)  
Queen Anns
- ✓ MASSACHUSETTS (Concord. New Hampshire. 603-225-1411)  
Essex and Hampshire
- ✓ MICHIGAN (E. Lansing. Michigan. 517-337-6650)  
Berrien
- ✓ MINNESOTA (Twin Cities. Minnesota 612-725-3276)  
Goodhue and Rice
- ✓ MISSISSIPPI (Jackson. Mississippi. 601-965-4900)  
Sharkey and Sunflower
- ✓ MISSOURI (Columbia. Missouri. 314-875-5374)  
Christian. Dade. Greene and Ripley
- ✓ NEW HAMPSHIRE (Concord. New Hampshire. 603-225-1411)  
Belknap. Carroll. Merrimack. Rockingham and Strafford
- ✓ NEW JERSEY (State College. Pennsylvania. 814-234-4090)  
Sussex
- ✓ NEW YORK (Cortland. New York. 607-753-9334)  
Delaware. Sullivan and Ulster
- ✓ NORTH CAROLINA (Asheville. North Carolina. 704-259-0321)  
Avery. Bladen. Burke. Haywood. Henderson. Macon. Mitchell and  
Scotland
- ✓ OHIO (Columbus. Ohio. 614-231-3416)  
Portage and Summit
- ✓ PENNSYLVANIA (State College. Pennsylvania. 814-234-4090)  
Centre and Venango
- ✓ RHODE ISLAND (Concord. New Hampshire. 603-225-1411)  
Providence
- ✓ SOUTH CAROLINA (Asheville. North Carolina. 704-259-0321)  
Bamberg. Barnwell. Berkeley. Clarendon. Colleton. Lee. Oconee.  
Orangeburg. Poincoter and Richland
- TENNESSEE (Asheville. North Carolina. 704-259-0321)  
Carter. Hamilton and Walker
- ✓ UTAH (Salt Lake City. Utah. 801-524-4430)  
Cache. Emery. Sevier and Wayne
- ✓ VIRGINIA (Annapolis. Maryland. 301-269-5448)  
Caroline. James City. Prince William and Smyth
- ✓ WISCONSIN (Green Bay. Wisconsin. 414-465-2682)  
Grant. Monroe. Richland. Sauk and Vernon



**GENERAL DIRECTIONS  
PERENNIAL WEED CONTROL**

This product is recommended for control of certain perennial weeds such as ARTEMISIA, CANADA THISTLE, CURLY DOCK, FALSEDANDELION (CATEAR), FESCUE, LEAFY SPURGE, ORCHARDGRASS, QUACKGRASS, RUSSIAN KNAPWEED, TIMOTHY, WILD ARTICHOKE, WILD ASTER, WILD CARROT, AND YELLOW ROCKET IN ESTABLISHED WOODY ORNAMENTALS, AND FRUIT and for control in non-crop areas such as roadways, around buildings and fences, railroad and utility rights-of-way and installations.

**SOIL SURFACE TREATMENT:** Apply as a soil surface treatment in late fall from November 15 to February 15. Old weed growth should not be removed before application. Do not make surface application to areas which have been cultivated during the fall or summer prior to application.

**INCORPORATED TREATMENT:** Apply and incorporate immediately after application in late fall or very early spring before May 1.

**ANNUAL WEED CONTROL**

This product is recommended for control of annual grassy and broadleaved weeds and certain perennial weeds such as: ANNUAL BLUEGRASS (POA ANNUA), BLUEGRASS, BULL THISTLE, CAMPHORWEED, CARPETWEED, CHICKWEED, CITRON MELON, COFFEEWEED, CRABGRASS, CUDWEED, DANDELION, DOGFENNEL, EVENINGPRIMROSE, FIDDLENECK, FLORIDA PURSLANE (PURSLEY), FOXTAIL, GISEKIA, GROUNDSEL, HENBIT, HORSETAIL, JERUSALEMOAK GOOSEFOOT, KNOTWEED, LAMBSQUARTERS, MAYPOPS, MILKWEED VINE (MORRENIA ODORATA), MINERS LETTUCE, NATALGRASS, OLD WITCHGRASS, PEPPERGRASS, PINEAPPLEWEED, PLANTAIN, PURSLANE, RAGWEED, RED DEADNETTLE, REDROOT PIGWEED, ROSARYPEA, RUSSIAN THISTLE, SHEPHERDSPURSE, SMARTWEED, SPANISHNEEDLES, SPURGE, TEAWEED, TEXAS PANICUM (HURRAHGRASS), WILD BARLEY, WILD MUSTARD, WILD RADISH, AND YELLOW WOODSORREL.

Apply either in early spring before seeds or annual weeds germinate or after cultivation has removed the growing weeds. Distribute granules uniformly over the soil surface. If treatment is followed by application of 1/2 to 1 inch of water by overhead irrigation the lower rates are recommended. Shallow incorporation immediately after application is recommended in southern areas. In other areas shallow incorporation or sprinkler irrigation is suggested when applications are made during periods of high temperatures.

**FRUIT IN ORCHARDS AND NURSERIES**

This product may be used on the following crops: do not allow livestock to graze treated orchards:

**BEST AVAILABLE COPY**

A. Apple. Blueberry. Cherry. Grape. Nectarine. Peach. Pear. Prune Plum: May be used on bearing, non-bearing and nursery stock. Do not apply within 4 weeks after transplanting. Do not make application within one month of harvest of stone fruits. Apply 100 to 150 pounds per acre for annual weed control based on area actually treated. For control of perennial weeds apply 150 pounds per acre. Follow General Directions above.

→ B. Blackberry and Raspberry: For control of annual weeds follow General Directions above except use only on established plantings and do not apply more than 100 pounds per acre. Do not apply during new shoot emergence.

WOODY ORNAMENTALS. (SHELTERBELTS) AND FOREST PLANTINGS ESTABLISHED IN NURSERIES

For control of annual weeds this product may be used on the following plants. Apply 100 to 150 pounds per acre, based on areas actually treated. Do not apply until 4 weeks after transplanting. Follow General Directions above.

For control of perennial weeds listed, apply 150 to 200 pounds per acre. Do not apply more than 150 pounds per acre on AZALEA, BOXWOOD, EUONYMUS, FORSYTHIA, HEATHER, HOLLY, IVY, LILAC, LEUCOTHOE. Do not apply more than 150 pounds per acre on plantings less than one year old. Follow General Directions listed above.

Amur Corktree	Filbert	Mountain Ash
Arbor Vitae	Flowering Almond	Nandine
Ash	Flowering Quince	Oak
Azalea <sup>1</sup>	Forsythia	Osmanthus
Barberry	Gardenia	(Holy-olive)
Beauty Bush	Goldenraintree	Pachistima
Birch	Hackberry	Pecan
Boxelder	Heather	Photinia
Boxwood	Holly (Ilex spp.) <sup>2</sup>	Pittosporum
Camellia	Honeysuckle	Popular
Caragana	Ivy (Hederahelix)	Privet
Cleyera japonica	Juniper	Pyracantha
Cotoneaster	Kinnikinnick	Rhododendron
Cottonwood	Laurel	Rockrose
Crabapple	Leucothoe	Rose
Deutzia	Lilac	Russian Olive
Dogwood	Linden	Spirea
~ Eastern Red Cedar	Locust	Squaw-carpet
Elm	Magnolia	Weigela
Elwood Cypress	Maple	Willow
English Walnut	Mock Orange	Yew (Taxus)
Euonymus		

1 Kurume, Mollis. Hardy hybrid types and hardy native species only.  
2 Except I. crenata, I. rotunda, I. vomitoria



### PRECAUTIONS

Do not use in seed beds, transplant or cutting beds or greenhouses. Do not apply until 6 months after rooting or cuttings in the field, or after planting rooted cuttings. Do not apply on fruit stock within 3 months prior to or following grafting or budding of root stocks or planting of new grafts. Do not apply more than the maximum recommended amount per growing season. Do not use on "St. Lucie fine sand", "Arzell fine sand" or other light sand soils. Do not graze livestock in treated areas.

### WEED CONTROL IN CRANBERRIES

Control of perennial and annual, broadleaved and grassy weeds (including rushes and sedges) in cranberries can be obtained by use of this product. The following weeds are controlled: Sensitive Fern, Bracken Fern, Royal Fern, Hair Cap Moss, Common Horsetail, Water Horsetail (pipes), Rush (Juncus spp.), Dodder, Bluejoint grass, Rattlesnake grass (Manna grass), Summer grass, Velvetgrass, Bent Grasses, Little Hairgrass, Crabgrass, Rice Cutgrass, Bunch grass, Muskrat grass, Nutsedge (Nutgrass), Short Wiregrass, Wideleaf grass, Stargrass, Woolgrass, Cotton grass, Needle grass, Oniongrass, Arrow-leaved Tear Thumb, Beggersticks, Knotweed, Loosestrife, Marsh St. Johnswort, Tideland clover, Ragweed, Sorrel, Wild Strawberry, Aster, Buckbean, Hawkweed, Western Lilaeopsis, Marsh Pea Plantain, Smartweed (Marshpepper, Pennsylvania, Spotted, Swamp and Water). Application should be made in early spring while perennial weeds are still dormant and annual weeds have not started to germinate or in late fall after crop has been harvested. Apply pre-bloom or post-harvest.

#### Spring Treatment

AREAS OTHER THAN PACIFIC NORTHWEST - Apply 100 pounds per acre. Do not make spring application if treatment was made the previous fall. See specific directions for spring treatment in the Pacific Northwest.

PACIFIC NORTHWEST - Two equal applications may be made in spring with an interval of 3 to 6 weeks between treatments. Total application should be 100 pounds per acre.

#### Fall Treatment

In all growing areas apply 100 to 150 pounds per acre in the fall after crop has been harvested. Do not make fall application if treatment was made the previous spring. Use high rate only on very weedy bogs, or on bogs abandoned because of weeds.

**BEST AVAILABLE COPY**

PRECAUTIONS

Distribute uniformly. Avoid over-application which may result from over-lapping during treatment. Temporary reddening of cranberry plants may occur especially with late spring application or when applied on sandy bogs. Do not apply after cranberry plants start to bloom as yields may be reduced. Do not use on young beds or newly sanded beds or on bogs prior to or immediately after mowing for vines.

OTHER USES

NUTSEDGE CONTROL: To control nutsedge (nutgrass) on mineral soils in areas on which containers of nursery stock are to be grown or held and also for control in non-crop areas: apply 250 to 500 pounds per acre and incorporate immediately to a depth of 4 to 6 inches. Seasonal control of annual weeds will also be obtained. Containers or plants listed above may be placed on treated soil immediately. Do not transplant into treated soil.

UNDER ASPHALT: For general weed control under asphalt (in areas such as roadways, parking lots, recreational areas), after final grade is achieved, apply 250 to 300 pounds per acre (5 3/4 to 7 pounds per 1,000 square feet). Treated area should be covered with asphalt as soon as possible. For re-surfacing work, existing weeds should be sprayed with a contact herbicide prior to application.

WARRANTY: The manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled uses when applied according to label directions. Because of widely varying use conditions, it is impossible to eliminate all risks even when label directions are followed. Therefore, the manufacturer makes no other implied or expressed warranty nor is any agent of the manufacturer allowed to do so. Upon purchase of this product, the buyer assumes all risks associated with use of this product. In the event of damage resulting from a breach of warranty the buyer agrees to accept a refund of the purchase price of the product as full discharge of the manufacturer's liability.

