GORDON'S PROFESSIONAL TURE PRODUCTS

TEREMEC

S P

TURE FUNGICIDE

ACCEPTED SEP 3 1982 Under the Federal Insecticide. Fungicide, and Rodenticide Act. as amended, for the posticide registered under 2017-69.2 EPA Reg. 112.

CONTROLS: Snow Mold & Pythium Blight

ACTIVE INGREDIENT:

Chloroneb (1,4-dichloro-2,5-dimethoxybenzene)	65%
INERT INGREDIENTS	35%
JATOT	100%

U.S. Patent No. 3,265,564.

KEEP OUT OF REACH OF CHILDREN

CAUTION <

- 14 pt. bold - 18 pt. bold

See back of bag for additional precautionary statements.

NET WEIGHT THREE POUNDS



STOP! READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND FOLIOW
DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: May irritate eyes, nose, throat, and skin, Avoid breathing dust or spray mist. Avoid contact with eyes, skin, or clothing. Do not graze or feed clippings from treated areas to livestock.

Statement of Practical Treatment

If Swallowed: Induce vomiting by drinking water and touching back of throat with finger.

if Inhaled: Remove to fresh air.

If In Eyes: Rinse with water for 15 minutes.

If On Skin: Wash with soap and water.

Environmental Hazards

Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL L 14 pt. Bold

<u>Prohibitions:</u> Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Do not store near other pesticides or seeds.

<u>Pesticide Disposal:</u> Pesticide or rinse water that cannot be used according to label instructions must be disposed of acfording to applicable federal, state or local procedures.

Container Disposal: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue ... into application equipment. Then, dispose of bag in a sanitary landfill or by incineration if allowed by state and local authorities.

۲

TURF DISEASES

Snow Mold - For the control of snow mold (Typhula), apply 6 to 9 ozs. of TEREMEC 3P per 1000 sq. ft. in sufficient water to give complete coverage, usually 3 to 5 gallons. Apply as a spray as late in the fall as possible prior to snowfall.

Pythium Blight - For control of Pythium blight on established turf, apply TEREMEC SP as a preventative at a rate of 4 ozs. per 100 sq. ft. in 3 to 5 gallons of water. Apply during periods of high temperature and humidity favorable to the development of disease. Repeat in 5 to 7 days if conditions

For the control of Pythium blight on new seedings, apply TEREMEC SP at the rate of 4 ozs. per 1000 sq. ft. following emergence. If conditions remain favorable to the development of disease, repeat in 7 to 10 days.

WARRANTY:

The manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled uses when applied according to label directions. Because of widely varying use conditions, it is impossible to eliminate all risks even when label directions are followed. Therefore, the manufacturer makes no other implied or express warranty nor is any agent of the manufacturer allowed to do so. Upon purchase of this product, the buyer assumes all risks associated with use of this product. In the event of damage resulting from a breach of warranty, the buyer agrees to accept a refund of the purchase price of the product as full discharge of the manufacturer's liability.

646/882

EPA REG. NO. 2217-EPA EST. NO. 2217-KS-1

