



STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: May irritate eyes, nose, throat, and skin. Avoid breathing dust or spray mist. Avoid contact with eyes, skin, or clothing. Do not graze or feed clippings from treated areas to livestock.

Statement of Practical Treatment

If Swallowed: Induce vomiting by drinking water and touching back of throat with finger.

If Inhaled: Remove to fresh air.

If In Eyes: Rinse with water for 15 minutes.

If On Skin: Wash with soap and water.

Environmental Hazards

Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

STORAGE: Store in original container in a locked storage area inaccessible to children or pets.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

TURF DISEASES

Snow Mold — For the control of snow mold (Typhula), apply 8 to 9 ozs. of TEREMEC SP per 1,000 sq. ft. in sufficient water to give complete coverage, usually 3 to 5 gallons. Apply as a spray as late in the fall as possible prior to snowfall.

Pythium Blight — For control of pythium blight on established turf, apply TEREMEC SP as a preventative at a rate of 4 ozs. per 1,000 sq. ft. in 3 to 5 gallons of water. Apply during periods of high temperature and humidity favorable to the development of disease. Repeat in 5 to 7 days if conditions persist.

For the control of Pythium blight on new seedings, apply TEREMEC SP at the rate of 4 ozs. per 1,000 sq. ft. following emergence. If conditions remain favorable to the development of disease, repeat in 7 to 10 days.

Sclerotium Blight — For control of Sclerotium Blight on established turf, apply TEREMEC SP as a preventative at a rate of 4 ozs. per 1,000 sq. ft. in 3 to 5 gallons of water. Apply during periods of high temperature and humidity favorable to the development of disease. Repeat in 5 to 7 days if conditions persist.

For the control of Sclerotium Blight on new seedings, apply TEREMEC SP at the rate of 4 ozs. per 1,000 sq. ft. following emergence. If conditions remain favorable to the development of disease, repeat in 7 to 10 days.

Brown Patch (Rhizoctonia Blight) — For warm season lawns and turfs (such as zoysiagrass, St. Augustine, bermudagrass and similar grasses) apply in the fall or spring at the first signs of infection. Mix 5 ozs. TEREMEC SP in 10 to 15 gallons water to thoroughly cover 1,000 sq. ft. Make a second application in 3 to 4 weeks if the disease is severe or reappears.

For cool season lawns and turfs (such as creeping bentgrass, annual bluegrass, fescue and similar grasses) apply at first signs of infection. Mix 3 to 4 ozs. TEREMEC SP in 3 to 5 gallons of water to thoroughly cover 1,000 sq. ft. Repeat every week for 10 days during warm, damp weather.

NOTICE: Do not graze or feed clippings from treated areas to livestock.

WARRANTY:

The manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled uses when applied according to label directions. Because of widely varying use conditions, it is impossible to eliminate all risks even when label directions are followed. Therefore the manufacturer makes no other implied or explicit warranty nor is any agent of the manufacturer allowed to do so. Upon purchase of this product, the buyer assumes all risks associated with use of this product. In the event of damages resulting from breach of warranty, the buyer agrees to accept a refund of the purchase price of the product as full discharge of the manufacturer's liability.

646-887
EPA REG. NO. 2217-692
EPA EST. NO. 42761 M-1

GORDON'S 

PROFESSIONAL TURF PRODUCTS

TEREMEC
SNOW MOLD
PYTHIUM BLIGHT
SP

TURF FUNGICIDE

CONTROLS: SNOW MOLD, PYTHIUM BLIGHT, SCLEROTIUM BLIGHT, & BROWN PATCH.

ACTIVE INGREDIENT:

Chloroneb (1,4-dichloro-2,5-dimethoxybenzene) 65%

INERT INGREDIENTS 35%

TOTAL 100%

U.S. Patent No. 3,265,564

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back of bag for additional precautionary statements.

ACCEPTED

SEP 15 1987

U.S. Department of Agriculture
Federal Insecticide, Fungicide, and Rodenticide Act
EPA Reg. No. 2217-692

NET WEIGHT THREE POUNDS

BEST AVAILABLE COPY

Mfd. by



pbi / Gordon
CORPORATION
KANSAS CITY, KANSAS 64110