

TEREMEC^(R) SP TURF FUNGICIDE

CONTROLS SNOW MOLD, BROWN PATCH, PYTHIUM AND SCLEROTIUM BLIGHT.

ACTIVE INGREDIENT:

Chloroneb (1,4-dichloro-2,5-dimethoxybenzene)	65%
INERT INGREDIENTS	35%
TOTAL	100%

Teremec^(R) is a registered trademark of PBI/Gordon Corporation
US Patent No. 3,265,564

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back of bag for Precautionary Statements and Statement of Practical Treatment.

NET WEIGHT THREE POUNDS

646/392 APXXXX92
EPA REG NO 2217-692
EPA EST NO 42761-MS-1

MFD. BY PBI/GORDON CORPORATION
KANSAS CITY, KANSAS 66118

ACCEPTED
APR 30 1992

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg No. 2217-692

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: May irritate eyes, nose, throat, and skin. Avoid breathing dust or spray mist. Avoid contact with eyes, skin, or clothing. Do not graze or feed clippings from treated areas to livestock.

Statement of Practical Treatment

IF SWALLOWED: Induce vomiting by drinking water and touching back of throat with finger.

IF INHALED: Remove to fresh air.

IF IN EYES: Rinse with water for 15 minutes.

IF ON SKIN: Wash with soap and water.

Environmental Hazards: Do not apply directly to any body of water. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

STORAGE: Store in original container in a locked storage area inaccessible to children or pets.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

- TURF DISEASES -

Do not apply this product through any type of irrigation system.

SNOW MOLD - For the control of Snow Mold (*Typhula spp.*), apply 6 to 9 ounces of TEREMEC SP per 1,000 square feet in sufficient water to give complete coverage, usually 3 to 5 gallons. Apply as a spray as late in the fall as possible prior to snowfall.

PYTHIUM BLIGHT - For control of pythium blight on established turf, apply TEREMEC SP as a preventative at a rate of 4 ounces per 1,000 square feet in 3 to 5 gallons of water. Apply during periods of high temperature and humidity favorable to the development of disease. Repeat in 5 to 7 days if conditions persist.

For the control of pythium blight on new seedlings, apply TEREMEC SP at the rate of 4 ounces per 1,000 square feet following emergence. If conditions remain favorable to the development of disease, repeat in 7 to 10 days.

SCLEROTIUM BLIGHT - For control of sclerotium blight on established turf, apply TEREMEC SP as a preventative treatment at a rate of 4 ounces per 1,000 square feet in 3 to 5 gallons of water. Apply during periods of high temperature and humidity favorable to the development of disease. Repeat in 5 to 7 days if conditions persist.

For the control of sclerotium blight on new seedlings, apply TEREMEC SP at the rate of 4 ounces per 1,000 square feet following emergence. If conditions remain favorable to the development of disease, repeat in 7 to 10 days.



BROWN PATCH (RHIZOCTONIA BLIGHT) - For warm season lawns and turfs (such as zoysiagrass, St. Augustinegrass, bermudagrass and similar grasses) apply in the fall or spring at the first signs of infection. Mix 5 ounces TEREMEC SP in 10 to 15 gallons of water to thoroughly cover 1,000 square feet. Make a second application in 3 to 4 weeks if the disease is severe or reappears.

For cool season lawns and turfs (such as creeping bentgrass, annual bluegrass, fescue and similar grasses) apply at first signs of infections. Mix 3 to 4 ounces TEREMEC SP in 3 to 5 gallons of water to thoroughly cover 1,000 square feet. Reapply every week to 10 days during warm, damp weather.

NOTICE: Do not graze or feed clippings from treated areas to livestock.

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.